

# Induced Mutations for Food and Energy Security: Challenge of Inducing Unique Mutants for New Cultivars and Molecular Research

Hitoshi NAKAGAWA<sup>\*1</sup> and Hiroshi KATO

## Abstract

Following the construction of the Gamma Field at the Institute of Radiation Breeding (IRB) in 1960, mutation breeding was accelerated in Japan. The facility is used to artificially induce mutations with a higher radiation dose (up to 2 Gy/day, that is *ca.* 300,000 times that of natural background) at a higher frequency than occurs in nature. Since the unit became operational, the number of mutant cultivars generated in Japan increased until 2000–2010 and has since decreased. There have been 295 direct-use mutant cultivars representing 70 species generated through irradiation utilizing gamma-rays, X-rays, ion beams and chemical mutagenesis and *in vitro* culture. Each cultivar has been registered and released in Japan, with approximately 79% of these induced by radiation. There have been 335 indirect-use mutant cultivars, including 298 rice, of which 150 cultivars (50.3%) were derived from the semi-dwarf mutant cv. “Reimei” or its offspring. The economic impact of these mutant cultivars, primarily of rice and soybean, is very large. Some useful mutations are discussed for rice, such as low digestible-protein content, low amylose content, giant embryo and non-shattering. Useful mutations in soybean such as radio-sensitivity, fatty acid composition, lipoxygenase lacking, glycinin rich and super-nodulation have been identified. A similar series of advantageous mutations have been found in Japanese pear and other crops through various screening methods. The achievements of biological researches such as characterization and determination of deletion size generated by gamma-rays, the effect of deletion size and the location are identified. Similarly, genetic studies generated through the use of gamma-ray induced mutations, such as phytochrome research, aluminum tolerance and epicuticular wax have also been conducted in Japan. A unique mutation induction technique for outcrossing Italian ryegrass is also explained. Mutation breeding is a very interesting and useful technology for isolating genes and for elucidating gene functions and metabolic pathways in various crops. Records show that mutation induction is a very useful conventional breeding tool for developing superior cultivars. The IRB is well equipped with appropriate facilities and equipment that will contribute to future mutation breeding developments and it will be a contributor in solving various genomic, proteomic and metabolic problems.

**Keyword:** genetic analysis, gamma-ray irradiation, Gamma Field, mutation breeding, mutant cultivar

---

Accepted on March 22, 2017

\*<sup>1</sup> Current affiliation: Central Research Laboratory, Hamamatsu Photonics K.K., 5000 Hirakuchi, Hamakita-ku, Hamamatsu, Shizuoka, 434-8601

# 食料・エネルギー安全保障に貢献する誘発突然変異 ：品種育成や分子遺伝学的研究のための 新たな突然変異体誘発への挑戦

中川 仁<sup>\*1</sup>・加藤 浩

## 抄 録

1960年、放射線育種場にガンマーフィールドが建設されて以来、我が国では突然変異育種が加速された。ガンマーフィールドは人工的に突然変異の誘発率を高める施設であり、線源から最も近い地点では1日に2グレイ (Gy) (自然界の30万倍：1日で1,000年分) 照射できる。我が国の突然変異品種登録数は、2000-2010年がピークとなり、現在は減少傾向にある。これまでの、ガンマ線、X線、イオンビーム、化学誘発剤、培養変異を利用した突然変異直接利用品種数は70作物、295にのぼり、その79%が放射線による突然変異である。直接利用品種との交配による間接利用品種数は335であり、その大半はイネ (298品種) であり、そのうちの150品種 (50.3%) は「レイメイ」に由来する。突然変異品種の経済効果は、イネとダイズ品種の貢献が大きい。有用な突然変異としてイネの低易消化性タンパク質、低アミロース含量、巨大胚および難脱粒性、ダイズの放射線感受性、脂肪酸組成改変、リポキシゲナーゼ欠失、高グリシニンおよび根粒超着生などを記述した。また、生物学的研究の成功例として、ガンマ線が誘発する突然変異と欠失の大きさ、欠失の位置と大きさの差による突然変異形質の変化が解明された。実験材料としての成功例は、フィトクローム研究、アルミニウム耐性および無ワックス特性の解明である。また、放射線育種場で開発された他殖性のイタライアンライグラスの変異作出技術についても紹介した。突然変異育種は遺伝子単離や遺伝子機能や代謝経路の解明において魅力的かつ有用なテクノロジーである。突然変異品種数が示すように、突然変異誘発技術は優れた品種を育成するための簡易な従来育種ツールである。アジアでは韓国とマレーシアにガンマ線照射施設が建設され、突然変異育種が推進されている。放射線育種場はガンマーフィールドなどの有用な施設を有し、今後も突然変異育種の発展と生物学的問題解決に貢献していくことが期待される。

キーワード：遺伝解析、ガンマ線照射、ガンマーフィールド、  
突然変異育種、突然変異品種

---

平成28年11月17日受付 平成29年3月6日受理

\*1 現 浜松ホトニクス株式会社 中央研究所

## Content

- I Introduction
- II Achievements from biological research on mutations induced by gamma-ray irradiation
  - 1) Deletion size generated by gamma-ray irradiation
  - 2) Different size and location of deletion generates different phenotypes
- III Useful mutations from various screening methods
  - 1) Low digestible-protein content
  - 2) Glutinous rice (low amylose content)
  - 3) Giant embryo
  - 4) Non-shattering
  - 5) Radio-sensitivity
  - 6) Fatty acid composition
  - 7) Lipoxygenase-lacking soybean
  - 8) Glycinin-rich soybean
  - 9) Super-nodulation
  - 10) Japanese pear and apple resistant to *Alternaria* disease
  - 11) Others
- IV Genetic studies of the useful mutations induced by acute or chronic gamma-ray irradiation
  - 1) Phytochrome
  - 2) Aluminum tolerance
  - 3) Wax-free mutation of sorghum
  - 4) Chronic gamma-ray irradiation facilities in Asia
- V Mutation breeding of outcrossing crops
- VI Mutation breeding and cultivars released in Japan
  - 1) The number of cultivars developed by mutation breeding
  - 2) The economic impact of mutant cultivars in Japan
- VII Conclusions
- Acknowledgements
- Literature cited

## I Introduction

After the construction of the Gamma Field, considered the world's largest radiation field (100 m radius with an 88.8 TBq  $^{60}\text{Co}$  source at the center; Fig. 1), at the Institute of Radiation Breeding (IRB) in Ohmiya-machi (now Hitachi-Ohmiya), Ibaraki, Japan in 1960, mutation breeding was accelerated through cooperative research with national and prefectural breeding laboratories, private companies and universities in Japan (Yamaguchi 2001).



Fig. 1. Gamma Field of IRB

In the New York Times (Broad 2007), Dr. P. J. L. Lagoda of the Joint FAO/IAEA was quoted to say, “Spontaneous mutations are the motor of evolution. We are mimicking nature in this. We’re concentrating time and space for the breeder so he can do the job in his lifetime. We concentrate how often mutants appear – going through 10,000 to one million – to select just the right one”.

The concept and objectives of the IRB’s Gamma Field has the same goals for the plant breeder. The facility has an irradiation tower installed with an 88.8 TBq  $^{60}\text{Co}$  at the center of a circular field with a radius of 100 m (Nakagawa 2010), and used to artificially induce mutations at a higher frequency than that occurs in nature. The radiation dose at the nearest point of the field (10 m from the center: *ca.* 2 Gy/day) is estimated to be about 300,000 times that of normal and natural background radiation when it is operated for 8 hours per day (Fig. 2). At the farthest point (100 m from the center: *ca.* 0.01 Gy/day), the radiation dosage is about 2,000 times the normal background radiation. This means that plants growing at the nearest point to the gamma-ray source are being treated to a 1,000 years of accumulated normal background rates of radiation per day. Although we do not know all the genes or mechanisms of mutations, radiation breeding has produced many useful mutant cultivars and contributed greatly to the farmers and industries of Japan and all over the world. In 1991, the Ministry of Agriculture, Forestry and Fisheries (MAFF) of Japan launched the Rice Genome Research Program, with the aim of fully decoding the rice genome in three phases over a 21-year period. With the cooperation of 10 participating countries (Sasaki and Burr 1998), the genome sequencing of the 12 rice chromosomes was completed in 2005 (International Rice Genome Sequence Project 2005). Following this achievement, molecular genetic studies based on the results of the genome sequencing project became the most powerful tool for selecting mutants of certain characteristics in rice. This is anticipated to revolutionize mutation breeding success in rice, and be applicable to a number of other important crop species.

In this report, the mutant cultivars developed mainly by gamma-ray irradiation in Japan are discussed. In addition, their economic impacts in Japan, as well as molecular studies performed to contribute to food security

and elucidate the mutation at the DNA level, are described.

Part of this report was presented in the 53<sup>rd</sup> Gamma Field Symposium held in Mito, Ibaraki, Japan in 2014.




Fig. 2. Irradiation tower located in the center of the Gamma Field and its irradiation field.

## II Achievements from biological research on mutations induced by gamma-ray irradiation

### 1) Deletion size generated by gamma-ray irradiation

Naito *et al.* (2005) studied the deletion sizes of transmissible and non-transmissible mutations induced with gamma-ray and carbon ion beam irradiation by utilizing sophisticated pollen-irradiation methods in *Arabidopsis*. Many mutants induced with these ionizing irradiations possess extremely large deletions (more than 6 Mbp) and many also have small deletions (1 or 4 bp), which are normally transmissible. The larger 6-Mbp deletions were found to not be transmissible to the next generation.

In rice, the same trends were observed (Table 1). Morita *et al.* (2009) researched the frequency of transmission of different mutations possessing different deletion sizes obtained with gamma-ray irradiation. Among 24 gamma-ray induced mutants, three exhibited 1-bp substitution, 15 exhibited a small deletion, four exhibited large deletions and two exhibited inversions. Among 15 mutants with small deletions, six mutations including *cao* (*chlorophyllide-a oxygenase*), *cps* (*entcopalyl diphosphate synthase*), *ga3os* (*GA3-beta-hydroxylase*), *gid* (*GA-insensitive dwarf*), *gluA1* (*glutelin A1*) and *gluA2* (*glutelin A2*) exhibited 1-bp deletion; and nine mutations including *cao-g2* and *ga3ox-g2*, *kao-g1*, *kao-g2*, *pla1-g1* and *-g2* (*Plastochron1*), and *wx-g1*, *-g2*, and *-g3* exhibited deletion of 2–16 bp. In contrast, four mutations had large deletions including one *gid2-g1*, one *gluB4/5* (*glu1: Glutelin B4/5*), one *glb1* ( *$\alpha$ -globulin*) and one *wx-g4* mutant with 42.2, 129.7, 62.8 and 9.4 kb, respectively. As a result, gamma-ray induced mutations transmittable to the next generation are primarily classified into four groups: (1) those with a base substitution; (2) those with small deletions (1–16 bp);

**Table 1.** I Gamma ray irradiation-induced deletions, base substitutions, and inversions and the size of mutation (Morita *et al.* 2009).

Mutation Type	Gene	Size (bp)
Small Deletion	<i>CAO (cao-g1)</i>	1
	<i>CAO (cao-g2)</i>	3
	<i>CPS (cps-g1)</i>	1
	<i>GA3ox (ga3ox-g1)</i>	1
	<i>GA3ox (ga3ox-g2)</i>	3
	<i>GID1 (gid1-g1)</i>	1
	<i>GluA1 (gluA1-g1)</i>	1
	<i>GluA2 (gluA2-g1)</i>	1
	<i>KAO (kao-g1)</i>	4
	<i>KAO (kao-g2)</i>	16
	<i>PLA1 (pla1-g1)</i>	5
	<i>PLA2 (pla2-g1)</i>	5
	<i>Wx (wx-g1)</i>	2
	<i>Wx (wx-g2)</i>	5
	<i>Wx (wx-g3)</i>	6
Large Deletion	<i>GID2 (gid2-g1)</i>	42, 200
	<i>Glb (glb1)</i>	62, 800
	<i>GluB4/B5 (glu1)</i>	129, 700
	<i>Wx (wx-g4)</i>	9, 400
Base Substitution	<i>GluA2 (gluA2-g2)</i>	1
	<i>PLA1 (pla1-g2)</i>	1
	<i>Wx (wx-g5)</i>	1
Inversion	<i>Wx</i>	1, 284, 800
	<i>PLA2</i>	3, 208, 500

*CAO*: chlorophyll b deficiency; *CPS*, *KAO* and *GA3ox*: gibberellin deficiency; *GID*: gibberellin insensitivity; *GluA* and *GluB*: glutelin deficiency; *Glb*: alpha-globulin deficiency; *PLA*: shortened plastochron; *Wx*: glutinous endosperm

(3) those with extremely large deletions; and (4) those with a large inversion of over 1 Mbp with small deletions. The mechanisms of generating extremely large deletions and inversions may be similar—the former mutation areas were missing and the latter mutation areas were reversely placed again. It is not known how difficult it may be to generate mutants with medium-sized deletions (1.0–5.0 kb) through gamma-ray irradiation, although some reports mentioned that neutrons with high linear energy transfer (LET) can induce deletions ranging between 300 bp–12 kb (Sun *et al.* 1992; Li *et al.* 2001; Nagano *et al.* 2008). However, it is interesting that inversions are not considered rare events following gamma-ray irradiation, as is the case for sorghum (Mizuno *et al.* 2013)—this is explained in the following section.


## 2) Different size and location of deletion generates different kinds of phenotypes

In the course of plant evolution, genes are often duplicated in tandem, resulting in functional redundancy. The analysis of function of these genes by developing double mutants may be difficult because they are located near each other and are tightly linked. Glutelin is a major digestible seed storage protein encoded by a multigene family. Mutants of tandem-duplicated glutelin genes were investigated for their genotypes and phenotypes. They represent a reversely repeated two-loci event (Fig. 3), with both regions coding for mRNA of glutelin production. Various mutants with low glutelin contents have been isolated using SDS-PAGE (polyacrylamide gel electrophoresis) (Iida *et al.* 1993, 1997). The mechanisms of low glutelin content in the mutants that have been studied suggest that the size and position of deletions generate different characteristics of mutations (Fig.


4). Some act as dominant or recessive genes, and these relationships between genotypes and phenotypes are provided as examples below.

*Low glutelin content1 (Lgc-1)* is a dominant mutation that reduces glutelin content in rice grain. Kusaba *et al.* (2003) reported that in *Lgc-1* homozygotes contain a 3.5-kb deletion between two highly similar glutelin genes that forms a tail-to-tail inverted repeat, that could produce a double-stranded RNA molecule, a possible potent inducer of RNA silencing (Fig. 3). As a result of this inverted repeat, glutelin synthesis is suppressed and the glutelin content is lowered. This was the first report that showed the mechanism of a mutation was RNA interference (RNAi) in plants. The *Lgc-1* provides an interesting example of RNA silencing among genes that exhibit various levels of similarity to a gene-induced RNA silencing.

The “*glu1*” is a gamma-ray-induced rice mutant, which lacks an acidic subunit of glutelin. Morita *et al.* (2007) elucidated that the *glu1* gene of the “*glu1*” mutant harbors a 129.7-kb deletion involving two highly similar, tandem-repeated glutelin genes, *GluB5* and *GluB4*. The deletion eliminates the entire *GluB5* and *GluB4* genes except for half of the first exon of *GluB5*. As a result, the phenotype of the *glu1* gene completely lacks the acidic subunit of glutelin and acts as a recessive gene for low glutelin content in rice grains (Fig. 4).



**Fig. 3.** Mechanism of low glutelin in LGC-1. A deletion containing the transcription termination signal between *GluB4* and *GluB5* causes generation of hairpin RNA with dsDNA region, which induces *GluB* mRNA degradation via RNA interference (based on Kusaba *et al.* 2003). (Courtesy of Prof. M. Kusaba, Hiroshima University)



**Fig. 4.** Comparison of phenotype, mode of inheritance and mechanism of mutation character between *glu1* and *LGCI* mutation with different sizes and place of deletion in the same region of two loci, *GluB4* and *GluB5*; *glu1*<sup>1</sup>: Morita *et al.* (2007); *Lgc1*<sup>2</sup>: Kusaba *et al.* (2003). (Courtesy of Dr. R. Morita, RIKEN)



The above examples illustrate that the position and size of deletions at the same locus can dramatically alter the phenotype of mutant gene expression through the process of transcription and translation. The *glu1*, which has a large 129.7-kb deletion, acts as a recessive gene, while the *LGC1*, which has 3.5-kb deletion that probably includes a terminal signal of the transcript region, acts as a dominant gene.


Furthermore, the *GluB5* and the *GluB4* have the same amino acid sequence in their acidic subunit, suggesting that only the mutation involving both *GluB5* and *GluB4* generates the resultant phenotype. This is the reason for the lack of the glutelin acidic subunit deleted in the “*glu1*” mutant.

Sequenced plant genomes exhibit that more than 14% of their genes are highly similar, tandem-repeated genes (Arabidopsis Genome Initiative 2000; International Rice Genome Sequence Project 2005). This finding suggests that gamma-rays can be an effective mutagen to generate knockout mutants of both loci and to use for analysis of tandem repeat and functionally redundant genes.

### III Useful mutations from various screening methods

#### 1) Low digestible-protein content

Although rice is not a high-protein grain crop, the protein content is *ca.* 7% when white rice is cooked. A mutant line with a low glutelin content was obtained from the ethyleneimine (EI) treatment of cv. “Nihon-masari”. The cv. “LGC-1” was developed from backcrossing this mutant with the original cv. “Nihon-masari” to eliminate undesirable characteristics, such as semi-sterility and semi-dwarfing (Iida *et al.* 1993). The seed protein of cv. “LGC-1” is mainly composed of a decreased amount of digestible glutelin induced by a dominant mutation that also increases the amount of indigestible prolamine. The characterization of this mutation and the mechanism were mentioned in the previous section. This construction of protein is disadvantageous for human digestion of rice grains, although the total amount of protein is similar to the original cultivar. As a result, cv. “LGC-1” is useful as a low-protein rice, and some clinical trials indicate that the cooked rice of this variety is a



**Fig. 5.** Development of new cultivars, cv. “LGC-Jun” and cv. “LGC-Katsu” with lower easy-to-digest protein content by the hybridization of a low-glutelin mutant cv. “LGC1” and an  $\alpha$ -globulin deficient mutant (named “89WPKG30–433”) of cv. “Koshihikari” (SDS-PAGE data) (based on Nishimura *et al.* 2005). (Courtesy of Prof. M. Nishimura, Niigata University)

useful and effective daily food for patients with kidney disease (Mochizuki and Hara 2000). The defect of cv. “LGC-1” is its eating quality and the presence of other loci that control the biosynthesis of digestible protein, such as globulin. Therefore, Nishimura *et al.* (2005) induced a mutant line “89WPKG30-433” that exhibited a deficiency in globulin through gamma-ray irradiation from the leading Japanese cv. “Koshihikari”, which is famous for a good eating quality. The “89WPKG30-433” was hybridized with cv. “LGC-1” and cvs. “LGC-Katsu” and “LGC-Jun” were selected from the hybrids (Fig. 5). The glutelin content of these two cultivars is as low as for cv. “LGC-1” and the globulin content is zero. The total digestible-protein content tested is about 30% of ordinary rice. As the eating quality is highly improved and digestible-protein content is lower than for cv. “LGC-1”, these two cultivars will greatly help in the dietary management of proteins in cases of chronic renal failure.

## 2) Glutinous rice (low amylose content)

Stickiness of cooked rice is one of the most important characteristics of rice cultivars, which are diverse and unique among peoples of the world. In general, people in Japan as well as Korea, northern Thailand, Myanmar and southern China prefer sticky rice. Amylose contents are closely related to this character and range from 0 (waxy: glutinous) to higher than 20, especially in indica-type rice. In Japan, glutinous rice has a special utilization for “okowa” and “mochi” production for festivals and celebrations, as well as non-glutinous popular cultivars used for daily cooking, which exhibit ca. 17% amylose content. This waxy locus (*Wx*) was mapped on chromosome 6 of rice (Iwata and Omura 1971) and knockout of *Wx* makes non-glutinous ordinary rice completely glutinous (*wx*). The waxy genes were identified to encode granule-bound starch synthesis, which is performed by a key enzyme in amylose synthesis of plants (Nelson and Pan 1995). In Japan, glutinous cv. “Miyuki-mochi” (Toda 1982) was induced from non-glutinous cv. “Toyonishiki”, glutinous cv. “Fujimi-mochi” from non-glutinous cv. “Aki-chikara”, and glutinous cv. “Odoroki-mochi” (Imbe *et al.* 2004) from non-glutinous cv. “Takanari” through gamma-ray irradiation in the IRB Gamma Field.

There is another type of endosperm starch mutation termed “dull”, whose amylose content is not zero as found with the waxy (*wx*) mutation in rice. The dull mutation has ca. 10% amylose content, which is lower than the ca. 20% of non-waxy (*Wx*) rice, and exhibits partial stickiness when cooked. Genetic analysis of dull mutants induced by <sup>32</sup>P (beta-ray radiation) showed that the mutations were controlled by a single recessive gene which is non-allelic to the *wx* alleles (Okuno *et al.* 1983). One of the most popular dull cultivars is cv. “Milky Queen” induced by chemical mutagen (MNU: N-methyl-N-nitrosourea) treatment of the most popular cv. “Koshihikari” with amylose content of 9–12% (Ise *et al.* 2001). This dull phenotype is caused by *wx* locus mutation (*Wx-mq*) (Sato *et al.* 2002).

## 3) Giant embryo

The rice embryo contains a high level of proteins, fats, vitamins and some compounds good for human health. As the embryo grows, gamma-aminobutyric acid (GABA) accumulates following transformation from glutamic acid caused by soaking rice in water (Saikusa *et al.* 1994a, 1994b). Defatted rice embryos enriched with GABA are useful as a functional food for controlling blood pressure and sedative qualities related to sleeplessness and autonomic disorder (Okada *et al.* 2000). The GABA-accumulated brown rice is already on the market as a health food based on an ordinary rice variety. The giant embryo lines “EM40” and “GM15-34”, each possessing giant embryo by treating cv. “Kinmaze” with MNU (Satoh 1981), useful for this purpose are anticipated. The embryo volume of these lines is 3–4 times that of ordinary rice cultivars. The giant embryo trait of the “EM40” is known to be controlled by one recessive gene (*ge*) located on chromosome 7 (Satoh and Iwata

1990).

The utilization of the giant embryo mutants was previously limited to the development of rice bran oil. However, the nutritional value of cooked brown rice is beginning to attract the attention of consumers interested in healthy food. The first giant embryo mutant cv. “Haiminori” was developed by the hybridization between the “EM40” and cv. “Akenohoshi”, and selection from the hybrid population in 1990 (Nemoto *et al.* 2001). The embryo volume of cv. “Haiminori” is 3–4 times that of ordinary rice. After soaking in water for four hours, the amount of accumulated GABA in cv. “Haiminori” is about four times that of the traditional cv. “Nipponbare”. The cv. “Haiminori” is utilized in the commercial production of rice cakes. Common defects of giant embryo cultivars are lower germination and seedling establishment rates, sometimes only 40–50 % of that of cv. “Nipponbare”.

The *GIANT EMBRYO (GE)* gene has been identified as essential for controlling the size balance in rice, and the function of *GE*, which encodes CYP78A13, is predominantly expressed in the interfacing tissues of the both embryo and endosperm and controlling cell size in the embryo and cell death in the endosperm (Nagasawa *et al.* 2013). Development of giant embryo lines with good seedling establishment will be a continuing objective of the breeding.

#### 4) Non-shattering

Much attention has been paid recently to indica-type high-yielding genetic resources for improving biomass productivity as forage rice. Seed shattering is one of the most important characteristics of indica-type rice that requires improvement. Mutation in the dominant *qSH1* gene in domesticated rice eliminates the abscission layer and results in non-shattering seeds (Konishi *et al.* 2006). Thus, knockout of this dominant gene generates a non-shattering rice. Kato *et al.* (2006) attempted to induce a mutation at this locus in cv. “Mohretsu”. The cv. “Mohretsu” is a mutant forage rice cultivar induced through NMU treatment and through gamma-ray irradiation of seeds in IRB, National Institute of Agrobiological Sciences (NIAS), which was eventually developed as the direct-use, non-shattering mutant cv. “Minami-yutaka” for use as rice silage. Sakai *et al.* (2013) attempted to induce mutation at the same locus in the high-yielding traditional cv. “Taporuri” introduced from Taiwan through gamma-ray irradiation of seeds in IRB, and successfully released the direct-use non-shattering mutant cv. “Ruri-aoba” as a rice silage crop. The loci or genes controlling non-shattering in these newly induced mutant cultivars have not been identified. However, through genetic analyses of rice cultivars and wild relatives, as well as chromosomal segment substitution lines and an induced shattering mutant line derived from gamma-ray irradiation (Zhou *et al.* 2012), several quantitative trait loci associated with seed shattering have been identified. These includes *SH4* which promotes hydrolyzing of an abscission zone cells during the abscission process (Li *et al.* 2006), *OsCPL1 (Oryza sativa CTD phosphatase-like 1)* that enhances the development of the abscission layer during panicle development (Ji *et al.* 2010) and *SHAT1 (Shattering Abortion1)*, the AP2 domain-containing transcription factor gene responsible for abscission zone development (Zhou *et al.* 2012).

#### 5) Radio-sensitivity

Takagi (1969) identified two major genes that control radio-sensitivity in some soybean cultivars. When the 50% reduction rate ( $RD_{50}$ ) of root length was associated with exposure to acute irradiation of seeds or chronic irradiation of plants for the entire growth period, radio-sensitivity of soybean was identified. The cv. “Shinmejiro” is more than twice as high for radio-sensitivity as the resistant cv. “Tachisuzunari”. The differences in radio-sensitivity between the cultivars to chronic irradiation in the Gamma Field are controlled by a single recessive

*rs1* gene. A second recessive gene, *rs2*, was discovered in cv. “Goishi-shirobana”, whose activity is only expressed following acute seed radiation.

## 6) Fatty acid composition

Soybean is the most widely used source of edible oil for human consumption. Every oil crop produces unique oil with a specific fatty acid composition. For example, linolenic acid content is high and generally 8.0% in soybean, and is not found in oils from maize, traditional sunflower and traditional safflower. There is an extremely high diversity of fatty acid compositions across the oil crops. The biosynthesis of fatty acids in oil crops is summarized in Fig. 6. The biosynthesis for fatty acid composition, common to all plant species, involves a carbon elongation process (palmitic acid (16:0) → stearic acid (18:0)) and unsaturated reaction process (stearic acid (18:0) → oleic acid (18:1) → linoleic acid (18:2) → linolenic acid (18:3)). Another biosynthesis is a carbon elongation reaction unique to traditional rapeseed of oleic acid (18:1) → eicosenoic acid (20:1) → erucic acid (22:1). The particular fatty acid contents of these cultivars or species result from differences in activity or reactivity of their enzymes and the number of genes encoding each enzyme involved in the various steps related to carbon elongation and unsaturated reaction of their fatty acids.

The incorporation of target genes, known to modify the fatty acid components of oil crops were initially identified in natural populations of specific species, and then incorporated by various breeding techniques. White *et al.* (1961) previously identified inter-varietal variation of fatty acid composition in 251 soybean cultivars and reported that contents of linoleic and linolenic acids were controlled by polygenes and exhibited heritability typically associated with quantitative traits because the investigated hybrid population exhibited continuous variation across their fatty acid compositions. However, Takagi *et al.* (1986) reported that the expression and levels of oleic and linolenic acids were controlled by two and one gene, respectively. Recent research suggests that the effects of each gene on fatty acid content are small and indicates quantitative trait expression and larger




Fig. 6. Genes responsible for different fatty acid biosynthesis (based on Takagi and Anai 2006; Nakagawa *et al.*, 2011). (Courtesy of Prof. Y. Takagi, Saga University).

environmental effects.

Takagi and his colleagues identified 46 mutant lines from a 12,266 M<sub>2</sub> population generated by X-ray application to cv. “Bay”. Fatty acid compositional changes of the mutant lines are listed in Table 2. The mutant lines include “J3” with low palmitic acid content (6.1%) (Takagi *et al.* 1995), “J10” with high palmitic acid content (17.2%) (Takagi *et al.* 1995), “M25” with high stearic acid content (21.2%) (Rahman *et al.* 1995), “M23” with high oleic acid content (48.6%) (Rahman *et al.* 1994), “M5” with low linolenic-acid content (4.9%) (Takagi *et al.* 1990) and “B739” with high linolenic acid content (18.4%) (Takagi *et al.* 1989). In addition, lines with wider fatty acid composition variability were obtained from the hybridization between these mutants.

The 46 individual mutant populations generated from cv. “Bay” exhibit a wider range of fatty acid composition than the 99 landraces developed from naturally occurring variation (Table 3) (Takagi and Rahman 1995). This induction of fatty acid variability confirms that artificial mutation is useful for enhancing fatty acid diversity of soybean.

**Table 2.** Fatty acid composition (% of total oil) of different soybean mutants and their genotypes (Nakagawa *et al.* 2011).

Mutant Line	Fatty Acid Composition (%)					Genotype
	Palmitic Acid (16:0)	Stearic Acid (18:0)	Oleic Acid (18:1)	Linoleic Acid (18:2)	Linolenic Acid (18:3)	
J3	6.1	3.4	26.5	55.4	8.6	<i>sop1</i>
C1726	8.5	3.9	22.1	56.0	9.6	<i>fap1</i>
LPKKC-3	4.4	3.2	27.1	55.5	9.8	<i>sop1, fap1</i>
KK7	14.2	5.0	23.3	49.5	7.9	<i>fapx</i>
J10	17.2	5.0	19.7	48.4	9.7	<i>fap2</i>
HPKKJ10	21.6	5.7	18.0	45.6	9.0	<i>fapx, fap2</i>
KK-2	10.3	7.2	23.1	51.6	7.9	<i>st<sub>1</sub></i>
M25	9.5	21.2	15.6	44.0	9.8	<i>st<sub>2</sub></i>
M25KK2	9.4	31.4	12.4	38.4	8.4	<i>st<sub>1</sub>, st<sub>2</sub></i>
M11	10.3	4.3	35.8	41.4	8.2	<i>ol<sup>a</sup></i>
M23	8.9	4.4	48.6	29.5	8.5	<i>ol</i>
M24	10.0	5.3	27.3	51.5	5.9	<i>fanx<sup>a</sup></i>
M5	9.8	4.9	25.5	55.3	4.9	<i>fan</i>
LOLL	10.3	4.5	27.7	54.7	2.9	<i>fan, fanx<sup>a</sup></i>
B739	7.9	4.3	14.3	55.1	18.4	<i>lin<sup>h</sup></i>
cv. Bay	10.6	4.2	25.2	51.7	8.3	

**Table 3.** Mean and range of fatty acid composition (% of total oil) in 46 different soybean mutants and 99 cultivars developed by using natural resources (Takagi and Rahman 1995).

	Fatty Acid Composition (%)				
	Palmitic Acid (16:0)	Stearic Acid (18:0)	Oleic Acid (18:1)	Linoleic Acid (18:2)	Linolenic Acid (18:3)
<b>Mutants</b>					
Mean	10.6	5.2	26.5	49.8	7.9
Range	6.3-16.7	2.9-16.1	17.5-48.2	32.2-60.5	4.6-12.6
<b>Cultivars</b>					
Mean	12.1	2.9	25.1	51.9	8.0
Range	10.0-15.5	1.6-4.1	14.2-44.3	36.6-61.1	5.2-12.6

As mentioned above, Takagi and his colleagues isolated a significant number of mutant lines exhibiting varying fatty acid composition through use of X-rays. Hybridization of these mutant lines, selection, characterization and experimentation can attain the pyramiding of mutant genes exhibiting complementary gene action in order to modify expression of various fatty acids. Following the hybridization and examination of the subsequent generations, the inheritance of genes conditioning low and high palmitic, high stearic, high oleic, and low and high linolenic acids were elucidated.

Figure 6 illustrates the fatty acid biosynthesis pathway of soybean seed oil. Palmitoyl-ACP (16:0) is synthesized from malonyl-ACP (4:0) through the condensation (fatty acid elongation) cycle. Three types of  $\beta$ -ketoacyl-synthases ( $\beta$ KASs) are involved in this condensation cycle. Palmitoyl-ACP is synthesized by  $\beta$ KASI and  $\beta$ KASIII, and then stearoyl-ACP (18:0) is synthesized by  $\beta$ KASII. Following this process, stearoyl-ACP is converted into oleoyl-ACP by stearoyl-ACP desaturase. Acyl-ACP thioesterase catalyzes the acyl-ACPs, and palmitic, stearic and oleic acids are produced with these reactions. Then oleic acid is converted to linolenic acid through linoleic acid by two distinct microsomal fatty acid desaturases, omega-6 and omega-3. These fatty acid biosynthesis enzymes are encoded by *Gm $\beta$ KASII*, *GmSte*, *GmFAD2* and the *GmFAD3* gene family, respectively. As discussed above, genetic analysis of fatty acid content in soybean mutants suggests that several genes are involved in each step of fatty acid synthesis pathway.

Byrum *et al.* (1997) reported that one of the low linolenic-acid mutants, “A5”, carries a deletion in gene *GmFAD3*. Takagi’s group also identified a large deletion in gene *GmFAD3* of “J18” (Yamashita *et al.* 1998), *GmFAD2* of “M23” (Kinoshita *et al.* 1998) and *GmSte* of “KK2” (Rahman *et al.* 1998) with Southern-blot analysis and also confirmed that these deletions co-segregated with their fatty acid content phenotypes. However, the vast majority of fatty acid mutants of soybean have not been characterized in detail. A superior understanding of the relationships between the corresponding gene and the phenotype of individual mutants will be valuable for soybean oil breeding programs.

Takagi’s group determined the corresponding genes of three low linolenic-acid mutants (“M24”, “M5” and “J18”) and divided them into two groups: “M24” had a mutation on gene *GmFAD3-1a*, and “M24” and “J18” had a respective mutation on *GmFAD3-1b* (Anai *et al.* 2005). The molecular status of these mutated genes were as follows: a 1-bp deletion in the open reading frame (ORF) of gene *GmFAD3-1a*, a 19-bp deletion in the ORF of *GmFAD3-1b* and a large deletion in the region containing *GmFAD3-1b*, respectively. The mutant of *GmFAD3-2a* has not yet been obtained from Takagi’s mutant collection, but this mutant represents an important target for decreasing linolenic acid content in soybean. Because gene *GmFAD3-2a* is expressed in developing seeds and its product exhibits the correct enzymatic activity in yeast cells (Anai *et al.* 2005), it is anticipated that further study of the *GmFAD3-2a* mutant line may result in development of a near-zero linolenic acid soybean cultivar.

Allele-specific genotypic selection, through use of molecular markers related to these gene families, will be superior to phenotypic selection using gas-liquid chromatography. Multiple sources of alleles for each candidate gene isoform can provide further benefits in the breeding of germplasm with superior fatty acid composition and minimize fixation of alleles linked to target genes (Takagi *et al.* 1998; Anai *et al.* 2005; Takagi and Anai 2006; Nakagawa *et al.* 2011a).

## 7) Lipoxygenase-lacking soybean

Lipoxygenase is an enzyme that generates a beany-flavor within soybean and its processed products. When soybean meal is mixed with water, lipoxygenases oxidize the abundant unsaturated fatty acids in the soybean seed and produce aldehydes such as n-hexanal (Fig. 7). Heat treatment is generally used to reduce this




Fig. 7. The reaction pathway of the seed lipoxygenases (based on Arai *et al.* 1970). (Courtesy of Dr. M. Hajika, NARO)

flavor in soybean products; however, once produced, heat treatment will not completely eliminate this flavor and this results in soybean products with poor taste quality. Trials have been performed to determine a better method to remove these undesirable flavors from soybean products in the food processing industry.

Soybean seed has three lipoxygenases: L-1, L-2 and L-3 (Arai *et al.* 1970). In the 1980s, mutant cultivars lacking lipoxygenase were isolated from germplasm (Hildebrand and Hymowitz 1981; Kitamura *et al.* 1983, 1985) and both L-1 and L-3 lacking and L-2 and L-3 lacking lines were bred by hybridizations between the two types. However, neither L-1 and L-3 lacking plants nor plants lacking L-1, L-2 and L-3 were obtained through breeding and hybridization. Therefore, mutation breeding was conducted to induce soybean mutants lacking L-1, L-2 and L-3 (Kitamura *et al.* 1985). Expecting not only the mutations but also chromosome crossing-over, 100–150 g of  $F_2$  seeds derived from the cross between cv. “Kankei 2” (later named cv. “Kanto No. 102”, which lacks L-1 and L-3), and cv. “Kankei 1” (later registered as cv. “Yume-yutaka”, which lacks L-2 and L-3) were irradiated at the IRB in 1989.

The irradiated  $F_2$  seeds ( $M_1$ ) were planted to establish the  $M_1$  population and obtain  $M_2$  seeds. Harvested  $M_2$  seeds were analyzed for protein composition using SDS-PAGE. The  $F_2$  seeds derived from a cross between a line without L-1 and L-3 and a line without L-2 and L-3 were irradiated with gamma-rays. After surveying 1,813  $M_3$  seeds using SDS/PAGE, one mutant seed lacking L-1, L-2, and L-3 was selected (Hajika *et al.* 1991; Nakagawa *et al.* 2011a). This lipoxygenase-lacking line was accelerated to obtain the  $M_6$  generation in a field in summer and a greenhouse in winter. From the  $M_7$  generation, field tests were conducted to evaluate yield and agronomic characteristics. In 1995, one selected line was released as cv. “Ichi-hime” (Hajika *et al.* 2002), and exhibits similar agronomic characteristics to cv. “Suzu-yutaka”, the recurrent parent of “Kankei 1” and “Kankei 2”. Several all-lipoxygenase-lacking soybean cultivars were developed by crossing between normal soybean cultivars and the all-lipoxygenase-lacking mutant lines. Several of these lines were sent to other countries and many all-lipoxygenases-lacking cultivars have since been developed.

## 8) Glycinin-rich soybean

The major components of soybean seed storage proteins are glycinin (11S globulin) and  $\beta$ -conglycinin

(7S globulin), which account for about 70% of all seed proteins. The amount of sulfur-containing amino acids (methionine and cysteine) in glycinin is 3–4 times that of  $\beta$ -conglycinin. Compared with  $\beta$ -conglycinin, glycinin exhibits better processing properties in texturized and filmed soy foods as well as in tofu gels. The  $\beta$ -conglycinin is composed of  $\alpha$ -,  $\alpha'$ - and  $\beta$ -subunits. The cv. “Keburi” characterized by the absence of the  $\alpha'$ -subunit, and cv. “Mo-shi-dou Gong 503” with low levels of  $\alpha$ - and  $\beta$ -subunits, were identified in Japanese soybean germplasm collections (Kitamura and Kaizuma 1981). Ogawa *et al.* (1989) developed some lines from these spontaneous soybean mutants and showed that it was possible to increase the amount of glycinin by decreasing the amount of  $\beta$ -conglycinin. The cv. “Kari-kei 434” was developed as one line characterized by the lack of the  $\alpha'$ -subunit and a marked decrease of the level of  $\alpha$ - and  $\beta$ -subunits of  $\beta$ -conglycinin. However, neither  $\alpha$ -less nor  $\beta$ -less cultivars were identified in the germplasm collection. Therefore, knockout of the  $\alpha$ -subunit gene was attempted by gamma-ray irradiation of seed of cv. “Kari-kei 434” at the IRB, and five mutated seeds, which exhibited loss of the  $\alpha$ -subunit and  $\alpha'$ -subunit, were identified using SDS-PAGE in the  $M_2$  and  $M_3$  populations. The deleted form of the  $\alpha$ -subunit was controlled by a recessive gene and crossing studies showed that existence or nonexistence of the  $\alpha$ -subunit was independent of the  $\alpha'$ -subunit (Takahashi *et al.* 1996). In 2001, a new mutant cv. “Yumeminori” was developed from this mutation and released as a glycinin-rich and hypoallergenic soybean (Fig. 8; Takahashi *et al.*, 2004; Nakagawa *et al.* 2011a). The cause for the hypoallergenic property in cv. “Yumeminori” is due to; (a) the  $\alpha$ -subunit is a major allergen in soybean (Ogawa *et al.* 1991); and (b) GlymBd30K, which represents one major allergen, was effectively removed from soybean protein isolates using chemical and physical methods (Samoto *et al.* 1996). Almost all growth habits of cv. “Yumeminori” were similar to cv. “Tachinagaha” except for the slightly lower seed weight and ca. 4% higher protein content. However, the processing properties of tofu curd were poorer than those for normal soybeans presumably caused by the high reactivity of glycinin proteins in cv. “Yumeminori”. To solve this problem, further studies are required. Recently, it was clearly shown that the deficiency of the  $\alpha$ -subunit in a mutant line “ $\alpha$ -null(1)” was caused by the termination codon resulting from inserting four bases into gene *CG-2* encoding the  $\alpha$ -subunit (Ishikawa *et al.* 2006).

Today, these glycinin-rich soybeans are utilized for soymilk products with decreased allergic reaction.




Fig. 8. Pedigree chart of glycinin rich mutant cv. “Yumeminori” (based on Takahashi *et al.* 2004).  
(Courtesy of Dr. M. Hajika, NARO)

## 9) Super-nodulation

Super-nodulation is a character that generates a large number of root nodules in leguminous plants. Super-nodulating mutants of soybean may provide a new genetic resource for improving soybean productivity by higher nitrogen-fixation as well as for elucidating mechanism of rhizobium-plant interactions. A super-nodulating mutant “En6500” was isolated from a 2,800  $M_2$  population generated by treating cv. “Enrei” with 0.5% of




ethyl methanesulfonate (EMS) (Akao and Kouchi 1992). The “En6500” had an increased number of nodules, inherited as a Mendelian recessive trait (Kokubun and Akao 1994); however, its growth and yield performance were inferior to that of cv. “Enrei”. The cause of this reduced yield performance is presumably from the high consumption of carbohydrates by nodules and a decrease of total nitrogen assimilation through the reduction of nitrate absorption (Takahashi, *et al.*, 1995). Due to the inferior performance of this new germplasm, a second attempt utilizing traditional backcross breeding to cv. “Enrei” was initiated and the super-nodulating cv. “Kanto 100” was successfully selected at the National Institute of Crop Science (NICS), National Agriculture and Food Research Organization (NARO). The cv. “Kanto 100” exhibits super-nodulation and the yield is similar to that of cv. “Enrei” (Takahashi *et al.* 2003). This super-nodulating mutation has potential impact on the development of high efficiency, nitrogen-fixing soybeans as well as for elucidation of rhizobium-plant interaction.

#### 10) Japanese pear and apple resistant to *Alternaria* disease

A popular cultivar of Japanese pear (*Pyrus serotina* Rehd. var. *culta* Rehd.), cv. “Nijisseiki”, was a leading cultivar that occupied 28% of the total cultivated area of Japanese pear in 1990 in Japan. However, the cultivar is highly susceptible to the black spot disease, *Alternaria alternata* (Fr.) Keissler (= *Alternaria kikuchiana* Tanaka), one of the most serious diseases of pear (Nishimura *et al.* 1978). Growers need to spray fungicides several times during the growing season to counter the disease. To induce mutations resistant to the disease, small plants of the cv. “Nijisseiki” were planted every 4 meters within 37-63 m from the <sup>60</sup>Co source in 1962 and chronic gamma-ray irradiation was applied ( $30 \times 10^{-2}$  to  $4 \times 10^{-2}$  Gy/day) in the Gamma Field (Sanada *et al.* 1993). In 1981, nearly 20 years after the planting, a twig bearing green leaves exhibiting no symptom of the disease while all the other twigs were infected by the fungus and most leaves had many black spots. The resistant twig was found on a plant planted 53 m from the radiation source. As it was ascertained that there was no difference in other agronomic characteristics between the mutant and the original cultivar except for resistance to black spot disease, it was registered and released in 1991 with the name cv. “Gold Nijisseiki” (Sanada *et al.* 1993). It was registered under the same name in Australia in 2004 (Certificate Number 2533).

Dr. Sanada, one of the breeders of this cultivar mentioned, “The situation of mutation breeding on Japanese pear has been severely criticized because there have been no successful results”. Although it took them nearly 20 years to identify a useful mutation and 30 years for its registration, the release of cv. “Gold Nijisseiki” became a monumental achievement for the IRB’s Gamma Field.

At the same time, an easy and effective method for screening of resistance to the fungus was developed by treating leaf disks (punched out by 7 mm in diameter) of each branch by the toxin produced by the fungus in the culture (Sanada 1988). After that, Nakashima *et al.* (1982, 1985) isolated and identified the chemical structure of the toxin (“AK-toxin”) produced by the fungus of black spot disease. As a consequence, the breeding group of the IRB entered into a cooperative research program with the chemistry group in Kyoto University and established the further use of this approach. When the punched leaf disks are placed on filter paper in a Petri dish soaked with AK-toxin obtained either from the extracts of the fungal body or artificial synthesis, and kept for 2 days at 25°C, leaves of susceptible cultivars such as cv. “Nijisseiki” turn black and leaves of resistant cultivars such as cv. “Chojuyou” stay green (Fig. 9). The intact leaves and old leaves of cv. “Gold Nijisseiki” are completely resistant, but interestingly the cut young leaves are partially susceptible, probably because only leaf surface tissues (the L1 layer) became resistant but inner tissues still keep susceptible genotype. As a result, leaves of cv. “Gold Nijisseiki” is really resistant to the disease in the field. Following its development, two new mutant cvs. “Osa-Gold (Masuda *et al.* 1997, 1998)” and “Kotobuki Shinsui (Kitagawa *et al.* 1999)” were



**Fig. 9.** Resistance bioassay for black spot disease using the AK-toxin obtained from the culture of the fungus (based on Nakagawa 2009). Upper to lower leaf disks (1–5) indicate 1 (young) to 5 (older) leaf; cv. “Chojyurou”: highly resistant; cv. “Nijisseiki”: highly susceptible; cv. “Gold Nijisseiki”: resistant

developed in a short period using this unique screening method. The economic effect of this research has been great.

Furthermore, the same bioassay method can also be utilized for the selection of plant resistant to *Alternaria* blotch disease using the AM-toxin (Ueno *et al.* 1977) produced by the fungus *Alternaria mali* that attacks apple. A new mutant apple cv. “Houiku Indo” was selected from gamma-ray irradiated micro-propagated shoots of susceptible cv. “Indo” (total dose: 80 Gy) and found to be resistant to this disease (Yoshioka *et al.* 2001).

This research suggests that breeding of fruit trees requires patience and that development of simple, efficient and precise screening methods is very important for mutation breeding.

## 11) Others

Many other interesting mutations have been identified using various screening methods in Japan. Additional information about unique mutations, screening methods, and released cultivars is given in the Appendices of this report. Gamma Field Symposia (a series of proceedings; <http://www.nias.affrc.go.jp/eng/public/index2.html>) and Radiation Breeding Technologies (*ibid.*) also provide a wide range of information.

## IV Genetic studies of the useful mutations induced by acute or chronic gamma-ray irradiation

Spontaneous and induced-mutation resources have played important roles not only for mutation breeding but also in genetic studies and the elucidation of gene functions. Some examples are mentioned here.

### 1) Phytochrome

Takano *et al.* (2005) isolated *phytochrome B* (*phyB*) and *phy C* mutants from rice and produced all

combinations of double mutants. Seedlings of *phy B* and *phyB phyC* mutants exhibited a partial loss of sensitivity to continuous red light but still showed significant de-etiolation responses. The response to red light was completely lost in *phyA phyB* double mutants. These results indicate that *phyA* and *phyB* act in a highly redundant manner to control de-etiolation under red light. They also found that mutations in either the *phyB* or *phyC* locus cause moderate early flowering under a long-day photoperiod, but monogenic *phyA* mutations had little effect on flowering time. However, the *phyA* mutation in combination with the *phyB* or *phyC* mutation caused dramatically early flowering. The *phyB* mutants were generated by chronic gamma-ray irradiation with dose rates of 3-6 Gy/day (Takano *et al.* 2005).

## 2) Aluminum tolerance

Ma *et al.* (2005) isolated a mutant with high sensitivity to aluminum (Al) concentration from cv. “Koshihikari”, an Al-resistant japonica rice (Wu *et al.* 1997). The mutant was induced through chronic gamma-ray irradiation and exhibited the same phenotype as the wild-type; however, without Al tolerance. That is, M<sub>1</sub> plants were irradiated in the Gamma Field from 7 days before heading to 2 days after heading under 20 Gy/day for 8 days. The root elongation of the mutant was highly inhibited in the presence of 10 µM Al. The mutant also exhibited poorer root growth in acid soil. Genetic analysis showed that the high sensitivity to Al is controlled by a single recessive gene, mapped to the long arm of chromosome 6.


## 3) Wax-free mutation of sorghum

Sorghum (*Sorghum bicolor* (L.) Moench.) produces and deposits bloom or epicuticular wax on the surface of stems and leaves (Fig. 10 left). The deposition of this wax is a dominant trait (*Bm*) with variation under different environments (Jordan *et al.* 1983), and bloomless (*bm*) mutants have no bloom or wax on the surface of plant parts. This wax is thought to make an important contribution to tolerance to abiotic stresses such as drought through reduced cuticular transpiration (Blum 1975) and protection from ultraviolet (UV) light in the semiarid tropics (Jordan *et al.* 1983). Reducing cuticular wax and cuticle deposition is also known to increase susceptibility to the fungal disease northern corn leaf blight caused by *Exserohilum turcicum* (Jenks *et al.* 1994). However, *bm* sorghum tends to increase resistance to biotic stresses such as resistance to greenbug attacks (Peiretti *et al.* 1980; Nakagawa *et al.* 2011b), which is achieved through nonpreference (Weibel and Starks 1986), and sheath blight (Kasuga *et al.* 2001).

In Japan, sweet sorghum has recently become a candidate crop for bioethanol production by utilizing the sugars produced and stored in stems and lignocellulose of stems and leaves through fermentation (Nakagawa *et al.* 2013). The development of sorghum cultivars resistant to biotic-stresses will have advantages for low-input without pesticides and fungicide and sustainable biofuel production. In 2006, groups of 1,000 seeds of sweet sorghum cv. “Italian” were irradiated with different doses of gamma-rays (0, 100, 200, 400, 500, 800, and 1,000 Gy for 20 hours) in the Gamma Room at the IRB, NIAS. After germination tests of the M<sub>1</sub> seeds, the seedlings were transplanted in the field and self-pollinated M<sub>2</sub> seeds were obtained by bagging of M<sub>1</sub> inflorescences before heading. In 2008, six seeds from each M<sub>2</sub> line were planted in the field plots for the evaluation. Two *bm* mutant plants without visible epicuticular wax (bloom) on the stems (Fig. 10) were identified from the M<sub>2</sub> population irradiated with 400 Gy of gamma-rays. The two self-pollinated lines were planted in the field and identified as pure lines with no variations. At the same time, these two lines were hybridized with the original cv. “Italian” plants, and the F<sub>1</sub> seeds and self-pollinated F<sub>2</sub> seeds were obtained. In 2010, 96 F<sub>2</sub> seeds from each hybridization were planted in the field for genetic and molecular analyses. The F<sub>2</sub> population segregated for individuals with




**Fig. 10.** Wild type (left) and *bm* mutant (right) of sorghum cv. “Italian”.



**Fig. 11.** Genomic inversion in the *bm* mutant. An inversion of *ca.* 1.4 Mbp occurred on the chromosome 6 (based on Mizuno *et al.* 2013). Arrows indicate cloned sequence from the wild-type (WT) and the *bm* mutant genome. At the junction, four bases (TACT) or two bases (AA) were deleted in the *bm* mutant genome. Because of this inversion, the upstream region of the Sb06g023280/WBC11 gene (black arrow) was exchanged with the downstream sequence. (Courtesy of Dr. H. Mizuno, NARO)

and without epicuticular wax at a frequency of 74:22 ( $\chi^2 = 0.1277$ ;  $P = 0.7209$  for a 3:1 segregation ratio, Chi square test) suggesting that the *bm* phenotype was controlled by a single recessive nuclear gene (Nakagawa *et al.* 2011c; Mizuno *et al.* 2013). The leaf-sheath of  $F_2$  plants with the wild-type phenotype and those with the *bm* phenotype were subjected to RNA-seq analysis. In addition, the sorghum *bm* mutant and wild-type plants were examined using scanning electron microscopy, which showed that the wax was not located on the leaf surface of plants but deposited inside of the cells (Mizuno *et al.* 2013).

Total RNA of both wild-type and *bm* mutant were extracted from leaves, and converted to cDNA for massive parallel sequencing in an Illumina Genome Analyzer and differentially expressed genes were identified (Mizuno *et al.* 2013). Of the 31 downregulated genes, one gene was similar to the ABC transporter responsible for wax secretion in *Arabidopsis* (Bird *et al.* 2007; Panikashvili *et al.* 2007; Ukitsu *et al.* 2007). The induced *bm* mutant was identified to carry a 1.4-Mb genomic inversion proximal to the promoter region of Sb06g023280, which is the candidate gene of the *bm* mutant, with small deletions at both ends (Fig. 11). Using genome PCR, six *bm* mutant-phenotype progeny of the F<sub>2</sub> population were found to carry the same inversion. The analysis proved that the inversion involving the Sb06g023280 gene inhibited wax secretion in the bloomless sorghum, although the epicuticular wax was synthesized inside the cells.



**Fig. 12.** Gamma Phytotron located at Korean Atomic Energy Research Institute (KAERI) in Jeongup, Jeonbuk Province, Republic of Korea. (Courtesy of Dr. Si-Yong Kang, KAERI), A: A building of the Gamma Phytotron; B: Inside the Gamma Phytotron

#### 4) Chronic gamma-ray irradiation facilities in Asia

Two facilities for chronic gamma-ray irradiation were recently established in Asia. The Gamma Phytotron (Fig. 12) was established at the Advanced Radiation Technology Institute, Korea Atomic Energy Research Institute in Jeongeup, Jeollabuk-do, Rep. of Korea in 2005. The irradiation source is  $^{60}\text{Co}$  with the radioactivity strength of about 400 curies (Kang et al. 2010). The Gamma Greenhouse (Fig. 13) was established at the Malaysian Nuclear Agency, Ministry of Science, Technology and Innovation in Bangi, Selangor, Malaysia



**Fig. 13.** Gamma Greenhouse located at Malaysia Nuclear Agency, Ministry of Science, Technology and Innovation, Bangi, Kajang, Selangor, Malaysia. (Courtesy of Dr. Rusli bin Ibrahim, Malaysian Nuclear Agency). A: A building of the Gamma Greenhouse; B: Inside the Gamma Greenhouse

in 2008. The Gamma Greenhouse is a circular greenhouse with a radius of 15 m, installed with  $^{137}\text{Cs}$  with the radioactivity strength of about 800 curies at the center (Ibrahim 2010). As with the Gamma Field, both facilities are focused on the induction of mutation by chronic gamma-ray irradiation in growing plants of important crop species. As described above, chronic irradiation is a useful tool for the generation of mutant genome resources that have application for molecular analysis as well as conventional breeding.

## V Mutation breeding of outcrossing crops

Mutation breeding has been primarily performed in seed propagated, self-pollinated species. Although several methods have been widely used for screening of mutants in self-pollinated species by the single-seed descent approach (Stadler 1930; Nybom 1954) and by one-plant-one-grain method (Yoshida 1962), these methods have not been applied to cross-pollinated species. Ukai (1990) developed a new and efficient method - the “crossing-within-spike-progenies method”- for obtaining mutants of cross-pollinated species in a temperate forage grass, Italian ryegrass (*Lolium multiflorum* Lam.). This method composes (1) taking seeds separately from each spike from a population of plants following gamma-ray irradiation, (2) sowing the seeds in a hill plot as a spike-progeny, (3) isolating each hill from others at the time of flowering and allowing the open-pollination of plants within hills and (4) taking seeds from each of the hills and sowing the seeds in hill progenies for the screening of mutants. This procedure is repeated each year. Using 300 Gy of gamma-ray irradiation for the seed, the frequency of chlorophyll mutations was approximately 70.6% per hill progeny and 1.87% per plant; in comparison, open-pollinated populations exhibited only 10% per progeny and 0.12% per plant. This method has application in other wind- or insect-pollinated outcrossing crop species.

## VI Mutation breeding and cultivars released in Japan

In a 2015 search regarding the number of induced-mutation cultivars in the IAEA database (<http://mvgs.iaea.org/AboutMutantVarieties.aspx>), China had the most described induced-mutation cultivars with 810, Japan was second with 481 and India was third with 330 (including those with doubled chromosome numbers through colchicine treatment). The total number of mutant cultivars, including direct-use mutant cultivars and indirect-use cultivars exceed these totals because all mutant cultivars have not yet been listed by breeders. A selection of mutant cultivars developed in Japan, including their economic impact of these cultivars, and their characteristics are reviewed here.

### 1) The number of cultivars developed by mutation breeding

The numbers of direct-use and indirect-use (hybrid) mutant cultivars registered in Japan in each 5-year period from 1960 to 2015 are shown in Fig. 14. The number of registered direct-use cultivars rapidly increased until 2000 when 65 cultivars were registered in five years (13 cultivars per year). This number has since fallen, with 49 cultivars registered during 2001-2005, 54 cultivars during 2006-2010 (*ca.* 10 cultivars per year) and 27 cultivars during 2011-2015. The number of indirect-use cultivars primarily generated in rice steadily increased with 79 during 2001-2005 and 80 during 2006-2010 but decreased to 58 during 2011-2015. This reduction can




Fig. 14. Number of cultivars developed by mutation breeding in each five years from 1961–2015. The total number of direct use cultivars is 295 and that of indirect use cultivars is 334.




Fig. 15. Numbers and percentages of total 295 cultivars developed by mutation breeding using various methods in Japan (2016). Chemical mutagen does not include colchicine.



be turned around if agronomically useful, direct-use mutant cultivars, such as cv. “Reimei”, possessing the *sd1* dwarf gene for rice are induced and utilized by rice breeders. It is presumed that such breeders will gravitate toward this and other cultivars carrying advantageous unique genes.

There have been 295 direct-use mutant cultivars comprising 70 species generated through irradiation utilizing gamma-rays, X-rays and ion beams, chemical mutagenesis and *in vitro* culture (somaclonal variation), registered and released in Japan (Fig. 15). Approximately 79% of these were induced by radiation, including gamma-rays (59.3%), ion beams (9.2%), X-rays (8.8%) and other radiation sources such as UV radiation or those providing no information of radiation source (1.4%). Those induced by somaclonal variation and chemical mutagens (not including those with doubled chromosome numbers through colchicine treatment), are 14.6 and 6.8%, respectively. Recently, the development of mutant cultivars of flower and food crops, generated by ion beam irradiation has been a growing area of mutation induction in Japan.

Table 4 shows the number of registered direct-use mutant cultivars of some crops developed by radiation, gamma-rays, and those irradiated at the IRB, NIAS. These include 43 mutant cultivars of rice (*Oryza sativa* L.), 17 of soybean (*Glycine max* (L.) Merrill), four of wheat (*Triticum aestivum* L.), four of barley (*Hordeum vulgare* L.), five of barnyard millet (*Echinochloa esculenta* (A. Braun) H. Scholz: glutinous mutants for

**Table 4.** Number of registered direct-use mutant cultivars developed by radiation, gamma rays, and those irradiated in the Institute of Radiation Breeding (IRB), NIAS (2016).

	Cultivar <sup>1</sup>	Radiation	Gamma rays	IRB <sup>2</sup>
70 Crops	295	228	176	127
Rice	43	21	20	19
Wheat	4	2	2	0
Barley	4	4	3	0
Soybean	17	16	15	9
Adzuki bean	1	1	1	1
Broad bean	1	1	1	0
Barnyard millet	5	5	4	4
Job's tear	2	2	2	2
Buckwheat	3	3	3	3
Tartary buckwheat	3	2	1	1
Taro	3	1	0	0
Potato	2	0	0	0
Sweet potato	1	1	0	0
Sugarcane	1	1	1	1
Burdock	5	5	5	4
Apple	2	2	2	2
Japanese pear	3	3	3	3
Peach	2	2	2	2
Loquat	1	1	1	1
Enoki mushroom	2	2	2	2
Chrysanthemum	60	55	40	38
Rose	10	7	7	6
Carnation	15	11	3	2
Sea pink ( <i>Limonium</i> )	6	6	6	0
Cytisus	8	8	8	8
Statice	6	6	6	0
Begonia	6	6	6	0
Lily	4	4	4	0
Clematis	2	2	2	2
Petunia	5	5	0	0
Margaret	4	4	0	0
Others	64	39	26	17

1 : Total number of mutant cultivars developed by radiation (gamma-ray, X-ray, ultraviolet, ion beams and unwritten radiation), chemicals (excluding colchicine treatment), somaclonal variation: 2: Number of mutant cultivars irradiated in the Institute of Radiation Breeding (IRB).

health food use), 60 of chrysanthemum (*Chrysanthemum*) and 10 of rose (*Rosa*). Among them, 126 cultivars (ca. 72.1% of gamma-ray induced cultivars) have been generated through gamma-ray irradiation in the Gamma Field, the Gamma Room and the Gamma Greenhouse of the IRB. This high percentage of gamma-ray-irradiated mutants indicates that mutation breeding via gamma-ray irradiation is an effective and highly successful approach for generation of commercial cultivars. Detailed data of developed cultivars are listed in Appendix 1, that includes cultivars induced through colchicine treatment.

The first mutant rice cultivar developed using gamma-ray irradiation was cv. “Reimei”, which means “dawn” in Japanese. Its development illustrated the potential of gamma-rays for breeding improvements in Japan. The cv. “Reimei”, registered in 1966 (Futsuhara 1968), was a successful case of an irradiation-induced semi-dwarf mutant. This cultivar exhibits a mutation of the *SD1* locus (Ashikari *et al.* 2002), which is same as the mutation of a miracle rice, cv. “IR8”, through cv. “Dee-geo-woo-gen” and later contributed to the “Green Revolution” of rice, and shows a culm 15 cm shorter than the original cv. “Fujiminori”. The semi-dwarf character is associated with the high-yielding ability and recorded the highest yield in Japan in 1967 (Futsuhara 1968)

In Japan, the total number of indirect-use mutant cultivars is 335, which includes 298 rice, 15 soybean, eight barley, six wheat, three tomato, one eggplant, one Japanese lawngrass (*Zoysia japonica* Steud), two mat rush (*Juncus effusus* L. var. *decipens* Buchen.) and one Job’s tear (*Coix lacryma-jobi* L. var. *ma-yuen* Stapf) in 2016 (Table 5). Detailed data of developed cultivars are listed in Appendix 2. Interestingly, among the 298 indirect-use mutant rice cultivars in 2016, 150 cultivars (50.3%) exhibited lodging resistance characteristics in the semi-dwarf cultivars derived from cv. “Reimei” or its offspring. This demonstrates that agronomically useful mutations can be efficiently and intensely utilized as parental lines to develop new cultivars with the same characteristic, and over the years the mutant gene multiplies itself in the farmers’ fields.

**Table 5.** Number of indirect-use mutant cultivars in Japan (2016).

Rice	Wheat	Barley	Soybean	Tomato	Others	Total
298	6	8	15	3	5	335


## 2) The economic impact of mutant cultivars in Japan

The increase of mutant rice cultivars derived from mutants generated by gamma-rays and sown in farmers’ fields in Japan since 1960 is shown in Fig. 16. The cv. “Reimei” was first cultivated on 61,598 ha in 1968 (<http://ineweb.narcc.affrc.go.jp/>). The number of mutant cultivars has been increasing and 99 mutant cultivars (two direct-use and 97 indirect-use cultivars) were in cultivation in 2005 (Nakagawa 2008).


The total cultivated area of mutant cultivars mostly derived from gamma-ray irradiation during 1961-2005 is shown in Fig. 17. This number increased after cv. “Reimei” was released for cultivation in 1968. The peak use of induced-mutation cultivars was 250,000 ha in 1986 and slightly exceeded 200,000 ha during 1994-2005. In 2005, the total cultivated area of mutant cultivars was 210,692 ha, which was 12.4% of the 1,702,000 ha cultivated for paddy rice in Japan (Nakagawa 2008).

The total crude income of farmers selling the brown rice of mutant cultivars has also increased with the expanding cultivation area. The amount of total income was estimated to be approximately 250 billion Yen (2.34 billion USD) in 2005 (Fig. 18) (Nakagawa 2008).


The following 17 cultivars were derived from mutant cultivars induced through gamma-ray irradiation (Appendix 2) and were cultivated on more than 5,000 ha during 2001-2005: cv. “Kinuhikari” (263,223 ha), cv. “Haenuki” (219,734 ha), cv. “Tsugaru-roman” (106,423 ha), cv. “Yume-akari” (66,491 ha), cv. “Yume-



**Fig. 16.** Total number of mutant rice cultivars derived from mutants mostly generated by gamma-rays, cultivated in farmers' fields during 1960–2005 in Japan (Nakagawa 2008, 2009).



**Fig. 17.** Total cultivated areas of mutant rice cultivars derived from mutants mostly generated by gamma-rays, cultivated in farmers' fields during 1960–2005 in Japan (Nakagawa 2008, 2009).



**Fig. 18.** Total estimated crude income of farmers obtained from cultivation of mutant rice cultivars mostly induced by gamma-rays in Japan during 1996–2005 (Nakagawa 2008, 2009).

tsukushi” (58,893 ha), cv. “Aichi-no-kaori” (53,697 ha), cv. “Asahi-no-yume” (51,049 ha), cv. “Mutsuhomare” (46,959 ha), cv. “Dontokoi” (17,008 ha), cv. “Yume-shizuku” (14,076 ha), cv. “Mine-asahi” (10,698 ha), cv. “Yume-hitachi” (10,440 ha), cv. “Yume-minori” (9,957 ha), cv. Aki-geshiki” (7,510 ha), cv. “Aki-roman (7,450 ha)”, “Miyama-nishiki (7,242 ha)”, and “Tsukushi-roman (5,533 ha)”. The following five mutant cultivars have been cultivated on more than 100,000 ha of farmers’ fields: cv. “Akihikari” (1,410,810 ha), cv. “Reimei” (886,188 ha), cv. “Kinuhikari” (263,223 ha), cv. “Haenuki” (219,734 ha) and cv. “Tsugaru-roman” (106,423 ha). Among them, only cv. “Reimei” is a direct-use mutation cultivar and the others are indirect-use cultivars (Nakagawa 2008; Appendix 1, 2).

The latest data of the top three paddy rice cultivars in 47 prefectures of Japan in 2013 shows that 14 indirect-use mutation cultivars are listed in 180,233 ha of paddy field (11.03% of total paddy field) (Table 6). Among them, cv. “Kinuhikari”, which is descended from a gamma-ray induced semi-dwarf and lodging-tolerant mutant line, cv. “Hokuriku 100 Gou” (Samoto and Kanai 1975), from the most popular tall cv. “Koshihikari” and semi-dwarf “IR8” (Koga *et al* 1989), is the most popular mutation variety with a good taste and shorter stems and covered 48,187 ha in 11 prefectures in 2013. The cv. “Haenuki” descended from cv. “Reimei”, covered 63% of Yamagata Prefecture’s paddy fields (43,848 ha). Interestingly, cvs. “Yume-hitachi”, “Kinumusume”, “Ikuhikari”, “Yume-tsukushi”, “Genki-tsukushi”, “Yume-shizuku”, “Saga-biyori” and “Nikomaru” are descended from cv. “Kinuhikari”. Therefore, the total cultivated area of cv. “Kinuhikari” and its descendants was 95,103 ha or 5.8% of all the paddy field in Japan. The real cultivated areas of mutant cultivars exceed these values because the data include only the top three cultivars of prefectures and do not include all cultivars. This means the economic impacts of mutant cultivars of rice are huge and the roles of cvs. “Reimei”, “Kinuhikari” and “Mineasahi” are very important in Japan.

There have been 16 direct-use mutant cultivars of soybean registered in Japan (Table 4) since cvs. “Raiden” and “Raikou” were developed by gamma-ray irradiation in 1960. The improved characteristics were early- and late-maturity, yellow hilum, seed-coat color, short-stems, numbers of pods/stem, lipoxxygenase-free and low allergens (Nakagawa *et al.* 2011a). Among them, cv. “Mura-yutaka” with yellow hilum color mutation is

**Table 6.** Mutant rice cultivars listed in the top 3 in 47 prefectures in 2013.

Variety	Cultivated Area (ha)	No. of Pref.	Origin of Mutant Cultivar
<b>Kinuhikari</b>	48,187	11	Hokuriku No. 100
<b>Haenuki</b>	43,848	1	Reimei
<b>Koshi-ibuki</b>	20,400	1	Reimei
<b>Asahi-no-yume</b>	17,598	3	Mineasahi
<b>Yume-tsukushi</b>	16,120	1	Kinuhikari
<b>Yume-shizuku</b>	8,463	1	Kinuhikari
<b>Kinumusume</b>	6,192	1	Kinuhikari
<b>Sagabiyori</b>	6,006	1	Mineasahi; Kinuhikari
<b>Genki-tsukushi</b>	3,627	1	Kinuhikari
<b>Nikomaru</b>	3,266	1	Kinuhikari
<b>Aki-roman</b>	2,358	1	Mineasahi
<b>Yume-hitachi</b>	2,331	1	Kinuhikari
<b>Mie-no-yume</b>	927	1	Mineasahi
<b>Ikuhikari</b>	911	1	Kinuhikari
<b>Total</b>	180,233	23	
Total paddy field in Japan	1,632,000		
%	11.03		

preferred in Japan and was induced by X-rays at Saga University (Nakamura *et al.* 1991; Appendix 1). There are 15 indirect-use mutant soybean cultivars (Table 5), of which four, including cv. “Ryuhou” are descended from a mutant cv. “Raiden” with induced early-maturity characteristics (Appendix 2). Table 7 shows the cultivated areas and estimated farmers’ income for soybean mutant cultivars in 1997, 2001, 2005, 2006, 2011 and 2014 in Japan. Among the direct-use mutant cultivars, the cultivated area of cvs. “Mura-yutaka” was 1,173 ha (1,403 ha in 2011), “Kosuzu” 134 ha (194 ha in 2011) and “Akita-midori” 40 ha (0 ha in 2011) in 2014. Among the indirect-use mutant cultivars, the cultivated area of cv. “Ryuhou”, a descendant of the gamma-ray induced mutant cv. “Raiden”, was the third of all soybean cultivars planted in Japan with 10,548 ha in 2011 and fourth with 9,600 ha in 2014; cv. “Nanbu-shirome”, which was selected from hybridization between cvs. “Raiden” and “Kitami-nagaha”, had 1,472 ha (2,132 ha in 2011); and cv. “Suzu-sayaka” descended from a lipoxygenase-free mutant, and cv. “Suzu-kaori”, which was selected from hybridization between cv. “Kosuzu”, a gamma-ray induced direct-use mutant cultivar from cv. “Natto Kotsubu”, and “Karikou 778F5”, a descendant of cv. “Kosuzu” (Appendix 1, 2), had 61 ha (65 ha in 2011) and 70 ha (56 ha in 2011), in 2014, respectively. The cv. “Kyo-shirotanba”, selected from hybridization between gamma-ray induced direct-use mutant cv. “Murasaki-zukin” and local cv. “Tamadaikoku” (Appendix 1, 2), was cultivated in 4 ha of Kyoto Prefecture. Thus, the economic impact of mutant cultivars of soybean is huge in Japan. The total cultivated area of mutant cultivars in farmers’ fields was 14,399 ha (10.5% of total cultivated area of 136,700 ha of soybean in Japan) in 2011 and 12,614 ha (9.5% of total area of 131,900 ha in 2014), and total farmers’ crude income was estimated as 11.6 billion Yen (*ca.* 116 million USD) in 2011.

**Table 7.** Cultivated areas (ha) of soybean mutant cultivars and income of farmers from the production in 1997, 2001, 2005, 2006, 2011 and 2014 in Japan.

Cultivar Name	1997	2001	2005	2006	2011	2014
Raiden <sup>1</sup>	80	8				
Wase-suzunari <sup>1</sup>	120					
Mura-yutaka <sup>1</sup>	3507	5910	2466	2265	1403	1173
Kosuzu <sup>1</sup>	498	863	576	512	194	134
Ichi-hime <sup>1</sup>		35	130			
Akita-midori <sup>1</sup>		8	87	95		40
Nanbu-shirome <sup>2</sup>	1246	1550	1534	1365	2132	1472
Tomo-yutaka <sup>2</sup>	2					
Suzu-no-ne <sup>2</sup>	10	50				
Eru-star <sup>2</sup>			447	43		
Suzu-sayaka <sup>2</sup>			10	234	65	61
Suzu-kaori <sup>2</sup>				25	56	70
Ryuhou <sup>2</sup>	1150	7050	8033	7955	10548	9600
Tsuya-homare <sup>2</sup>					1	
Kinu-sayaka <sup>2</sup>						61
Kyo-shirotanba <sup>2</sup>						4
Total	6,613	15,474	13,283	12,494	14,399	12,614
Total cultivated areas of soybean in Japan	83,200	143,900	134,000	142,100	136,700	131,900
%	7.95	10.75	9.91	8.79	10.53	9.56
Income of farmers (Million USD)	ca. 20	ca. 59	ca. 52	-	22.6 (115.7 with subsidy)	

<sup>1</sup> Direct-use mutant cultivar

<sup>2</sup> Indirect-use mutant cultivar

## VII Conclusions

A. M. van Harten (1998) describes in the Preface of “Mutation Breeding – Theory and Practical Application”:

“An explanation for the decreasing interest in mutation breeding, at least in most “developed” countries, may be that during the past two decades attention has become more and more directed towards studying the possibilities offered to plant breeding by various new molecular technologies...As a result of these developments mutation breeding seems to have lost part of its previous attraction for young researchers. It is even not inconceivable that mutation breeding, as a discrete branch of plant breeding, may sink into oblivion and that, as a consequence, much valuable knowledge on this topic built up throughout the years, will be lost.”

The record has also shown that mutation induction is a very useful conventional breeding tool for developing superior cultivars. Today, site-directed mutagenesis *in vivo* or *in vitro* cell can be envisioned and many researchers are conducting programs in this direction.

New fields of science and technologies were developed on the basis of achievements through the application of traditional or classic methods. It is highly desirable that the IRB continues its work while incorporating new discoveries and technologies. The IRB is well equipped with appropriate facilities and accumulated know-hows that will contribute to the future mutation breeding developments and it will remain a viable cooperator in solving the problems and opportunities of world food security and production.

The list of direct-use mutant cultivars (Appendix 1) includes information of crop species, cultivar names, year of registration in Japan and the registration number, breeding methods, the institutes of treatment and development, improved characteristics, ID numbers listed in IAEA database and references. The list of indirect-use (hybrid) mutant cultivars (Appendix 2) includes the crop species, cultivar name, parental lines (descendants of mutant cultivars are written in red font with the lineage), year of registration in Japan and the registration number, the institute of development, improved characteristics of original mutant cultivars, ID numbers listed in IAEA database, and references. These lists include only mutant cultivars identified from the references and a database of the Plant Variety Protection office at MAFF, Japan ([http://www.hinsyu.maff.go.jp/en/en\\_top.html](http://www.hinsyu.maff.go.jp/en/en_top.html)). Therefore, if the cultivar breeder did not mention that it was a mutation cultivar induced through a certain method, we cannot identify it. We did not include cultivars from spontaneous mutations, which are mentioned just as a mutant plant or twig found in the original cultivar, in the lists. We anticipate that these lists will demonstrate the importance of mutation breeding in Japan.

## Acknowledgements

The authors would like to express their sincere thanks to Dr. Bryan Kindiger, USDA-ARS, Grazinglands Research Laboratory for his critical reading of the manuscript.

The authors thank Prof. Makoto Kusaba of Hiroshima University, Prof. Minoru Nishimura of Niigata University, Prof. Yutaka Takagi of Saga University, Dr. Ryohei Morita of Riken Nishina Center for Accelerator-based Science, and Drs. Hiroshi Mizuno and Makita Hajika of NICS, NARO, for permission to use figures in this

report.

The authors also express their sincere thanks to Dr. Rusli bin Ibrahim, Senior Research Officer, Malaysian Nuclear Agency, Ministry of Science, Technology and Innovation, and Dr. Si-Yong Kang, Principal Researcher of the Advanced Radiation Technology Institute, Korea Atomic Energy Research Institute for providing us the photos of their facilities.

Part of this work was supported by the Cabinet Office, Government of Japan, Cross-ministerial Strategic Innovation Promotion Program (SIP), “Technologies for Creating Next-generation Agriculture, Forestry and Fisheries” (funding agency: Bio-oriented Technology Research Advancement Institution, NARO).

## Literature cited

- Akao, S. and H. Kouchi (1992) A supernodulating mutant isolated from soybean cultivar Enrei. *Soil Sci. Plant Nutr.* 38: 183-187.
- Anai, T., T. Yamada, T. Kinoshita, S. M. Rahman and Y. Takagi (2005) Identification of corresponding genes for three low- $\alpha$ -linolenic acid mutants and elucidation of their contribution to fatty acid biosynthesis in soybean seed. *Plant Sci.* 168: 1615-1623.
- Arabidopsis Genome Initiative (2000) Analysis of the genome sequencing of the flowering plant *Arabidopsis thaliana*. *Nature* 408: 796-815.
- Arai, S., M. Noguchi, M. Kaji, H. Kato and M. Fujimaki (1970) n-Hexanal and some volatile alcohols. Their distribution in row soybean tissues and formation in crude soy protein concentrate by lipoxygenase. *Agric. Biol. Chem.* 34 (9): 1420-1423.
- Ashikari, M., A. Sasaki, M. Ueguchi-Tanaka, H. Itoh, A. Nishimura, S. Datta, K. Ishiyama, T. Saito, M. Kobayashi, G. S. Khush, H. Kitano and M. Matsuoka (2002) Loss-of-function of a rice gibberellin biosynthetic gene, *GA20 oxidase (GA20ox-2)*, led to the rice ‘Green Revolution’. *Breed. Sci.* 52: 143-150.
- Bird, D., F. Beisson, A. Brigham, J. Shin, S. Greer, R. Jetter, L. Kunst, X. W. Wu, A. Yephremov and L. Samuels (2007) Characterization of Arabidopsis ABCG11/WBC11, an ATP binding cassette (ABC) transporter that is required for cuticular lipid secretion. *Plant J.* 52 (3): 485-498.
- Blum, A. (1975) Effect of the *Bm* gene on epicuticular wax and water relations of *Sorghum bicolor* L. (Moench). *Isr. J. Bot.* 24: 50-51.
- Broad, W. J. (2007) Useful mutants, bred with radiation. *New York Times*, August 27, 2007: [http://www.nytimes.com/2007/08/28/science/28crop.html?rref=collection%2Fbyline%2Fwilliam-j.-broad&action=click&contentCollection=undefined&region=stream&module=stream\\_unit&version=search&contentPlacement=1&pgtype=collection](http://www.nytimes.com/2007/08/28/science/28crop.html?rref=collection%2Fbyline%2Fwilliam-j.-broad&action=click&contentCollection=undefined&region=stream&module=stream_unit&version=search&contentPlacement=1&pgtype=collection).
- Byrum, J. R., A. J. Kinney, K. L. Stecca, D. J. Grace and B. W. Diers (1997) Alteration of the omega-3 fatty acid desaturase gene is associated with reduced linolenic acid in the A5 soybean genotype. *Theor. Appl. Genet.* 94: 356-359.
- Futsuhara, Y. (1968) Breeding of a new rice variety Reimei by gamma-ray irradiation. *Gamma Field Symposia* 7: 87-109.
- Hajika, M., K. Igita and K. Kitamura (1991) A line lacking all the seed lipoxygenase isozymes in soybean (*Glycine max* (L.) Merrill) induced by gamma-ray irradiation. *Jpn. J. Breed.* 41: 507-509.

- Hajika, M., M. Takahashi, K. Igita, S. Sakai and Y. Nakazawa (2002) A new soybean variety 'Ichihime'. Bull. Natl. Agric. Res. Cent. Kyushu Okinawa Reg. 40: 79-93 (in Japanese with English summary).
- Hildebrand, D. F. and T. Hymowitz (1981) Two soybean genotypes lacking lipoxygenase-1. J. Am. Oil Chem. Soc. 58: 583-586.
- Ibrahim, R. (2010) Malaysian Nuclear Agency Gamma Greenhouse. Plant Mutation Reports 2 (2): 52-54.
- Iida, S., E. Amano and T. Nishio (1993) A rice (*Oryza sativa* L.) mutant having a low content of glutelin and a high content of prolamine. Theor. Appl. Genet. 87: 374-378.
- Iida, S., M. Kusaba and T. Nishio (1997) Mutants lacking glutelin subunits in rice: mapping and combination of mutated glutelin genes. Theor. Appl. Genet. 94: 177-183.
- Imbe, T., N. Horisue, Y. Akama, A. Nakane, I. Ando, M. Suto, T. Hata, K. Ise, K. Numaguchi, H. Furutachi and M. Takadate (2004) Odorokimochi – a multipurpose high-yield glutinous rice variety. Bull. Natl. Inst. Crop Sci. 6: 19-35 (in Japanese with English summary).
- International Rice Genome Sequence Project (2005) The map-based sequence of the rice genome. Nature 436: 793-800.
- Ise, K., Y. Akama, N. Horisue, A. Nakane, M. Yokoo, I. Ando, T. Hata, M. Suto, K. Numaguchi, H. Nemoto, H. Furutachi and T. Imbe (2001) "Milky Queen", a new high quality rice cultivar with low amylose content in endosperm. Bull. Natl. Inst. Crop Sci. 2: 39-61 (in Japanese with English summary).
- Ishikawa, G., Y. Takada and T. Nakamura (2006) A PCR-based method to test for the presence or absence of  $\beta$ -conglycinin  $\alpha'$ - and  $\alpha$ -subunits in soybean. Mol. Breed. 17: 365-374.
- Iwata, M. and T. Omura (1971) Linkage analysis of reciprocal translocation method in rice plant (*Oryza sativa* L.). II. Linkage groups corresponding to the chromosome 5, 6, 8, 9, 10 and 11. Sci. Bull. Fac. Agric. Kyushu Univ. 30: 137-153 (in Japanese with English summary).
- Jenks, M. A., R. J. Joly, P. J. Peters, P. J. Rich, J. D. Axtell and E. N. Ashworth (1994) Chemically-induced cuticle mutation affecting epidermal conductance to water vapor and disease susceptibility in *Sorghum bicolor* (L.) Moench. Plant Physiol. 105: 1239-1245.
- Ji, H., S. Kim, Y. Kim, H. Kim, M. Eun, I. Jin, Y. Cha, D. Yun, B. Ahn, M. Lee, G. Lee, U. Yoon, J. Lee, Y. Lee, S. Suh, W. Jiang, J. Yang, P. Jin, S. R. McCouch, G. An and H. Koh (2010) Inactivation of the CTD phosphatase-like gene *OsCPL1* enhances the development of the abscission layer and seed shattering in rice. Plant J. 61: 98-106.
- Jordan, W. R., R. L. Monk, F. R. Miller, D. T. Rosenow, L. E. Clark and P. J. Shouse (1983) Environmental physiology of sorghum: I. Environmental and genetic control of epicuticular wax load. Crop Sci. 23: 552-558.
- Kang, S.-Y., J.-B. Kim, G.-J. Lee and D. S. Kim (2010) Gamma Phytotron: A new chronic gamma irradiation facility. Plant Mutation Reports 2 (2): 50-51.
- Kasuga, S., N. Inoue, H. Kaidai and H. Watanabe (2001) Effects of brown midrib and bloomless genes on the resistance to sheath blight (*Rhizoctonia solani* Kuehn) in sorghum. Grassl. Sci. 47: 256-261.
- Kato, H., H. Yoshioka, R. Iba, S. Ueda, S. Tsutsumi, H. Mitsueda, K. Wakasugi, T. Nakahara and H. Takeda (2006) A newly bred rice variety "Minamiyutaka" for whole-crop silage. Bull. Miyazaki Agricultural Experiment Station 41: 51-60 (in Japanese with English summary).
- Kinoshita, T., S. M. Rahman, T. Anai and Y. Takagi (1998) Genetic analysis of restriction fragment length polymorphism on the fatty acid synthesis in soybean mutants and their progenies: II. High oleic acid mutants with two microsomal  $\omega$ -6 fatty acid desaturase cDNAs as probes. Bull. Fac. Agr. Saga Univ. 83: 37-42.



- Kitagawa, K., M. Nagae, M. Uchida, K. Inoue, K. Murata, T. Masuda, T. Yoshioka and K. Kotobuki (1999) A new Japanese pear cultivar, 'Kotobuki Shinsui'. *Bull. Tottori Hort. Exp. Stn.* 3: 1-13 (in Japanese).
- Kitamura, K., C. S. Davies, N. Kaizuma and N. C. Nielsen (1983) Genetic analysis of a null-allele for lipoxygenase-3 in soybean seeds. *Crop Sci.* 23: 924-927.
- Kitamura, K. and N. Kaizuma (1981) Mutant strains with low level of subunits of 7S globulin in soybean seed. *Jpn. J. Breed.* 31: 353-359.
- Kitamura, K., T. Kumagai and A. Kikuchi (1985) Inheritance of lipogenase-2 and genetic relationships among genes for lipogenase-1, 2 and -3 isozymes in soybean seeds. *Jpn. J. Breed.* 35: 413-420.
- Koga, Y., H. Uchiyama, S. Samoto, S. Ishizaka, Y. Fujita, K. Okuno, Y. Uehara, M. Nakagahra, H. Horiuchi, K. Miura, K. Maruyama, T. Yamada, T. Yagi and K. Mori (1989) Breeding a new rice cultivar "Kinuhikari". *Bull. Hokuriku Natl. Agric. Exp. Stn.* 30: 1-24 (in Japanese with English summary).
- Kokubun, M. and S. Akao (1994) Inheritance of supernodulation in soybean mutant En65000. *Soil Sci. Plant Nutr.* 40: 715-718.
- Konishi, S., T. Izawa, S. Y. Lin, K. Ebata, Y. Fukuta, T. Sasaki and M. Yano (2006) An SNP caused loss of seed shattering during rice domestication. *Science* 312: 1392-1396.
- Kusaba, M., K. Miyahara, S. Iida, H. Fukuoka, T. Takano, H. Sassa, M. Nishimura and T. Nishio (2003) *Low glutelin content1*: A dominant mutation that suppresses the glutelin multigene family via RNA silencing in rice. *Plant Cell* 15: 1455-1467.
- Li, C., A. Zhou and T. Sang (2006) Genetic analysis of rice domestication syndrome with the wild annual species, *Oryza nivara*. *New Phytol.* 170: 185-194.
- Li, X., Y. Song, K. Century, S. Straight, P. Ronald, X. Dong, M. Lassner and Y. Zhang (2001) A fast neutron deletion mutagenesis-based reverse genetic system for plants. *Plant J.* 27: 235-242.
- Ma, J. F., S. Nagao, C. F. Huang and M. Nishimura (2005) Isolation and characterization of a rice mutant hypersensitive to Al. *Plant Cell Physiol.* 46: 1054-1061.
- Masuda, T., T. Yoshioka, K. Inoue, K. Murata, K. Kitagawa, H. Tabira, A. Yoshida, K. Kotobuki and T. Sanada (1997) Selection of mutants resistant to black spot disease by chronic irradiation of gamma-rays in Japanese pear 'Osanijisseiki'. *J. Japan. Hort. Sci.* 66 (1): 85-92.
- Masuda, T., T. Yoshioka, T. Sanada, K. Kotobuki, M. Nagae, M. Uchida, K. Inoue, K. Murata, K. Kitagawa and R. Yoshida (1998) A new Japanese pear cultivar 'Osa Gold', resistant mutant to the black spot disease of Japanese pear (*Pyrus pyrifolia* Nakai) induced by chronic irradiation of gamma-rays. *Bull. Natl. Inst. Agrobiol. Resour. (Japan)* 12: 1-11 (in Japanese with English summary).
- Mizuno, H., H. Kawahigashi, J. Ogata, H. Minami, H. Kanamori, H. Nakagawa and T. Matsumoto (2013) Genome inversion caused by gamma irradiation contributes to downregulation of a WBC11 homolog in bloomless sorghum. *Theor. Appl. Genet.* 126 (6): 1513-1520.
- Mochizuki, T. and S. Hara (2000) Usefulness of low protein rice in diet therapy in patients with chronic renal failure. *Jpn. J. Nephrol.* 42: 24-29 (in Japanese with English summary).
- Morita, R., M. Kusaba, S. Iida, T. Nishio and M. Nishimura (2007) Knockout of glutelin genes which form a tandem array with a high level of homology in rice by gamma irradiation. *Genes Genet. Syst.* 82: 321-327.
- Morita, R., M. Kusaba, S. Iida, H. Yamaguchi, T. Nishio and M. Nishimura (2009) Molecular characterization of mutations induced by gamma irradiation in rice. *Genes Genet. Syst.* 84: 361-370.
- Nagano, A. J., M. Fukazawa, M. Hayashi, M. Ikeuchi, H. Tsukaya, M. Nishimura and I. Hara-Nishimura (2008) AtMap1: a DNA microarray for genome deletion mapping in *Arabidopsis thaliana*. *Plant J.* 56: 1058-1065.

- Nagasawa, N., K. Hibara, E. P. Heppard, K. A. Vander Velden, S. Luck, M. Beatty, Y. Nagato and H. Sakai (2013) *GIANT EMBRYO* encodes CYP78A13, required for proper size balance between embryo and endosperm in rice. *Plant J.* 75: 592-605.
- Naito, K., M. Kusaba, N. Shikazono, T. Takano, A. Tanaka, T. Tanisaka and M. Nishimura (2005) Transmissible and nontransmissible mutations induced by irradiating *Arabidopsis thaliana* pollen with  $\gamma$ -rays and carbon ions. *Genetics* 169: 881-889.
- Nakagawa, H. (2008) Mutation breeding, status quo and future. *Techno Innovation* 68: 6-12 (in Japanese).
- Nakagawa, H. (2009) Induced mutations in plant breeding and biological researches in Japan, In "Induced plant mutations in the genomic era" ed. Q. Y. Shu, Food and Agriculture Organization of the United Nations, Rome, 51-58.
- Nakagawa, H. (2010) Gamma radiation facilities at the Institute of Radiation Breeding. *Plant Mutation Reports*, 2 (2): 55-56.
- Nakagawa, H., T. Anai, A. Okabe, K. Takahashi, M. Hajika and Y. Takagi (2011a) Mutation Breeding of Soybean in Japan, In "Breeding of Pulse Crops" eds. Khan S and M. I. Kozgar, Kalyani Publishers, New Delhi, India: 55-84.
- Nakagawa, H., M. Gau and H. Okuizumi (2013) Introduction of genetic resources and breeding of cellulosic crops and forage grasses. *Agriculture and Horticulture* 88 (7): 754-768 (in Japanese).
- Nakagawa, H., M. Nishimura and H. Okuizumi (2011b) Agronomic characteristics and aphid resistance of *bm* mutant lines of sweet sorghum developed through gamma-ray irradiation. *Japanese Journal of Grassland Science* 57 (Suppl.): 103 (in Japanese).
- Nakagawa, H., H. Okuizumi and M. Nishimura (2011c) Characteristics and inheritance of *bm* mutant induced by gamma-ray irradiation. *Breeding Research* 13 (Suppl. 1): 111 (in Japanese).
- Nakamura, D., H. Yokoh, Y. Hirota, K. Nonaka, T. Shigyo, H. Oh-ishi, O. Shigetomi, Y. Takagi and H. Kishikawa (1991) A new soybean cultivar "Murayutaka". *Bull. Saga Agric. Exp. Stn.* 27: 21-42 (in Japanese with English summary).
- Nakashima, T., T. Ueno and H. Fukami (1982) Structure elucidation of AK-toxins, host specific phyto-toxic metabolites produced by *Alternaria kikuchiana* Tanaka. *Tetrahedron Lett.* 23: 4469-4472. (1982).
- Nakashima, T., T. Ueno, H. Fukami, T. Taga, H. Masuda, K. Osaki, H. Otani, K. Kohmoto and S. Nishimura (1985) Isolation and structure of AK-toxin I and II. Host-specific phytotoxic metabolites produced by *Alternaria alternata* Japanese phytotype. *Agri. Biol. Chem.* 49: 807-815.
- Nelson, O. and D. Pan (1995) Starch synthesis in maize endosperms. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 46: 475-496.
- Nemoto, H., S. Iida, H. Maeda, T. Ishii, N. Nakagawa, T. Hoshino, M. Sakai, M. Okamoto, H. Shinoda and T. Yoshida (2001) A new rice cultivar with giant embryo "Haiminori". *Bull. Chugoku Natl. Agric. Exp. Stn.* 22: 25-40 (in Japanese with English summary).
- Nishimura, M., M. Kusaba, K. Miyahara, T. Nishio, S. Iida, T. Imbe and H. Sato (2005) New rice varieties with low levels of easy-to-digest protein, 'LGC-Katsu' and 'LGC-Jun'. *Breed. Sci.* 55: 103-105.
- Nishimura, S., M. Sugihara, K. Kohmoto and H. Otani (1978) Two different phases in pathogenicity of the *Alternaria* pathogen causing black spot disease of Japanese pear. *J. Fac. Agr. Tottori Univ.* 13: 1-10.
- Nybom, N. (1954) Mutation type in barley. *Acta Agric. Scand.* 4: 430-456.
- Ogawa, T., N. Bando, H. Tsuji, H. Okajima, K. Nishikawa and K. Sasaoka (1991) Investigation of the IgE-binding proteins in soybean by immunoblotting with the sera of the patients with atopic dermatitis. *J. Nutr.*

Sci. Vitaminol. 37: 555-565.

- Ogawa, T., E. Tayama, K. Kitamura and N. Kaizuma (1989) Genetic improvement of seed storage proteins using three variant alleles of 7S globulin subunits in soybean. *Jpn. J. Breed.* 39: 137-147.
- Okada, T., T. Sugishita, T. Murakami, H. Murai, T. Saikusa, T. Horino, A. Onoda, O. Kajimoto, R. Takahashi and T. Takahashi (2000) Effect of defatted rice germ enriched with GABA for sleeplessness depression, autonomic disorder by oral administration. *Nippon Shokuhin Kagaku Kogaku Kaishi* 47: 596-603 (in Japanese).
- Okuno, K., H. Fuwa and M. Yano (1983) A new mutant gene lowering amylose content in endosperm starch of rice, *Oryza sativa* L. *Jpn. J. Breed.* 33 (4): 387-394.
- Panikashvili, D., S. Savaldi-Goldstein, T. Mandel, T. Yifar, R. B. Franke, R. Hofer, L. Schreiber, J. Chory and A. Aharoni (2007) The *Arabidopsis* DESPERADO/AtWBC11 transporter is required for cutin and wax secretion. *Plant Physiol.* 145: 1345-1360.
- Peiretti, R. A., I. Amini, D. E. Weibel, K. J. Starks and R. W. Mcnew (1980) Relationship of bloomless (*bm bm*) sorghum to greenbug resistance. *Crop Sci.* 20: 173-176.
- Rahman, S. M., Y. Takagi, K. Kubota, K. Miyamoto and T. Kawakita (1994) High oleic acid mutant in soybean induced by X-ray irradiation. *Biosci. Biotech. Biochem.* 58 (6): 1070-1072.
- Rahman, S. M., Y. Takagi, K. Miyamoto and T. Kawakita (1995) High stearic acid soybean mutant induced by X-ray irradiation. *Biosci. Biotech. Biochem.* 59 (5): 922-923.
- Rahman, S. M., T. Kinoshita, T. Anai and Y. Takagi (1998) Genetic analysis of restriction fragment length polymorphism on the fatty acid synthesis in soybean mutants and their progenies: III. High stearic acid mutants with a stearyl-acyl carrier protein desaturase cDNA as a probe. *Bull. Fac. Agr. Saga Univ.* 83: 43-48.
- Saikusa, T., T. Horino and Y. Mori (1994a) Accumulation of  $\gamma$ -aminobutyric acid (GABA) in the rice germ during water soaking. *Biosci. Biotechnol. Biochem.* 58: 2291-2292.
- Saikusa, T., T. Horino and Y. Mori (1994b) Distribution of free amino acids in the rice kernel fractions and the effect of water soaking on the distribution. *J. Agric. Food Chem.* 42:1122-1125.
- Sakai, M., H. Nakano, M. Okamoto, K. Tamura, R. Kaji, Y. Tamura, T. Kataoka and R. Mizobuchi (2013) "Ruriaoba", a new rice forage variety for double harvesting. *Bull. Natl. Agric. Res. Cent. Kyushu Okinawa Reg.* 60: 1-12 (in Japanese with English summary).
- Samoto, S. and D. Kanai (1975) Studies on the mutation breeding in rice. I. Short stiff mutations induced by gamma-ray irradiation to the rice variety Koshihikari. *Jpn. J. Breed.* 25: 1-7 (in Japanese with English summary).
- Samoto, M., K. Takahashi, Y. Fukuda, S. Nakamura and Y. Kawamura (1996) Substantially complete removal of the 34 kD allergenic soybean protein, Gly m Bd 30K, from soy milk of a mutant lacking the  $\alpha$ - and  $\alpha'$ -Subunit of conglycinin. *Biosci. Biotechnol. Biochem.* 60: 1911-1913.
- Sanada, S. (1988) Selection of resistant mutants to black spot disease of Japanese pear by using host specific toxin. *Jpn. J. Breed.* 38: 198-204.
- Sanada, T., K. Kotobuki, M. Nishida, H. Fujita and T. Ikeda (1993) A new Japanese pear cultivar 'Gold Nijisseiki', resistant to black spot disease of Japanese pear. *Jpn. J. Breed.* 43: 455-461 (in Japanese with English summary).
- Sasaki, T. and B. Burr (1998) International Rice Genome Sequencing Project: the effort to completely sequence the rice genome. *Curr. Opin. Plant Biol.* 3: 138-141.

- Sato, H., Y. Suzuki, M. Sakai and T. Imbe (2002) Molecular characterization of *Wx-mq*, a novel mutant gene for low amylose content in endosperm of rice (*Oryza sativa* L.). *Breed. Sci.* 52: 131-135.
- Satoh, H. (1981) New endosperm mutations induced by chemical mutagens in rice, *Oryza sativa* L. *Jpn. J. Breed.* 31: 316-326.
- Satoh, H. and N. Iwata (1990) Linkage analysis in rice. On three mutant loci for endosperm properties, *ge* (giant embryo), *du-4* (dull endosperm-4) and *flo-1* (floury endosperm-1). *Jpn. J. Breed.* 40 (Suppl. 2): 268 (in Japanese).
- Sun, T., H. M. Goodman and F. M. Ausubel (1992) Cloning the Arabidopsis *GAI* locus by genomic substitution. *Plant Cell* 4: 119-128.
- Stadler, L. J. (1930) Some genetic effects of X-rays in plants. *J. Hered.* 21: 3-19.
- Takagi, Y. (1969) The second type of gamma-ray sensitive gene *RS<sub>2</sub>* in soybean *Glycine max* (L.) Merrill. *Gamma Field Symposia* 8: 83-94.
- Takagi Y. and T. Anai (2006) Development of novel fatty acid composition in soybean oil by induced mutation. *Oleoscience* 6 (4): 195-203 (in Japanese with English abstract).
- Takagi, Y., A. B. M. M. Hossain, T. Yanagita, T. Matsueda and A. Murayama (1990) Linolenic acid content in soybean improved by X-ray irradiation. *Agric. Biol. Chem.* 54(7): 1735-1738.
- Takagi, Y., A. B. M. M. Hossain, T. Yanagita and S. Kusaba (1989) High linolenic acid mutant induced by X-ray irradiation. *Jpn. J. Breed.* 39: 403-409.
- Takagi, Y., T. Matsuo and H. Kishikawa (1986) Inheritance of oleic and linolenic acid in soybean seeds. *Jpn. J. Breed.* 36: 163-176 (in Japanese with English summary).
- Takagi, Y. and S. M. Rahman (1995) Variation of different fatty acids in mutants in comparison with natural soybean varieties. *Bull. Fac. Agr. Saga Univ.* 79: 23-67
- Takagi, Y., S. M. Rahman and T. Anai (1998) Construction of novel fatty acid composition in soybean oil by induced mutation. *Gamma Field Symposia* 37: 17-28.
- Takagi, Y., S. M. Rahman, H. Joo and T. Kawakita (1995) Reduced and elevated palmitic acid mutants in soybean developed by X-ray irradiation. *Biosci. Biotechnol. Biochem.* 59 (9): 1778-1779.
- Takahashi, K., S. Shimada, H. Shimada, Y. Takada, T. Sakai, Y. Kono, T. Adachi, K. Tabuchi, A. Kikuchi, S. Yumoto, S. Nakamura, M. Ito, H. Banba and A. Okabe (2004) A new soybean cultivar "Yumeminori" with low allergenicity and high content of 11S globulin. *Bull. Natl. Agric. Res. Cent. Tohoku Reg.* 102: 23-39 (in Japanese with English summary).
- Takahashi, K., S. Yumoto, S. Nakamura, Y. Mizuno and K. Kitamura (1996) Inheritance of the  $\alpha$ -subunit deficiency of  $\beta$ -conglycinin in soybean (*Glycine max* L. MERRILL) line induced by  $\gamma$ -ray irradiation. *Breed. Sci.* 46: 251-255.
- Takahashi, M., J. Arihara, N. Nakayama, M. Kokubun, S. Shimada, K. Takahashi and M. Hajika (2003) Breeding of supernodulating soybean cultivar "Sakukei 4". *Bull. Natl. Inst. Crop Sci.* 4: 17-28.
- Takahashi, M., M. Kokubun and S. Akao (1995) Characterization of nitrogen assimilation in a supernodulating soybean mutant En6500. *Soil Sci. Plant Nutr.* 41 (3): 567-575.
- Takano, M., N. Inagaki, X. Xie, N. Yuzurihara, F. Hihara, T. Ishizuka, M. Yano, M. Nishimura, A. Miyao, H. Hirochika and T. Shinomura (2005) Distinct and cooperative function of phytochromes A, B, and C in the control of deetiolation and flowering in rice. *Plant Cell* 17: 3311-3325.
- Toda, M. (1982) The breeding of four new mutant varieties by gamma-rays in rice. *Gamma Field Symposia* 21: 7-18.

- Ueno, T., T. Nakashima, M. Uemoto, H. Fukami, S. N. Lee and N. Izumiya (1977) Mass spectrometry of *Alternaria mali* toxins and related cyclodepsipeptides. *Biomed. Mass Spectrom.* 4: 134-142.
- Ukai, Y. (1990) Application of a new method for selection of mutants in a cross-fertilizing species to recurrently mutagen-treated populations of Italian ryegrass. *Gamma Field Symposia* 29: 55-69.
- Ukitsu, H., T. Kuromori, K. Toyooka, Y. Goto, K. Matsuoka, E. Sakuradani, S. Shimizu, A. Kamiya, Y. Imura, M. Yuguchi, T. Wada, T. Hirayama and K. Shinozaki (2007) Cytological and biochemical analysis of COF1, an *Arabidopsis* mutant of an ABC transporter gene. *Plant Cell Physiol.* 48(11): 1524-1533.
- Van Harten, A. M. (1998) Mutation breeding - theory and practical applications, Cambridge University Press, Cambridge, UK: 353pp.
- Weibel, D. E. and K. I. Starks (1986) Greenbug nonpreference for bloomless sorghum. *Crop Sci.* 26: 1151-1153.
- White, H. B. J., F. W. Quackenbush and A. H. Probst (1961) Occurrence and inheritance of linolenic and linoleic acid in soybean seeds. *J. Am. Oil Chem. Soc.* 38: 113-117.
- Wu, P., B. Zhao, J. Yan, A. Luo, Y. Wu and D. Senadihra (1997) Genetic control of seedling tolerance to aluminum toxicity in rice. *Euphytica* 97: 289-293.
- Yamaguchi, I. (2001) Forty years of mutation breeding in Japan – Research and fruits. *Gamma Field Symposia* 40: 1-14.
- Yamashita, Y., T. Kinoshita, S. M. Rahman, K. Okada, T. Anai and Y. Takagi (1998) Genetic analysis of restriction fragment length polymorphism on the fatty acid synthesis in soybean mutants and their progenies: I. Low linolenic acid mutants with a microsomal  $\omega$ -3 fatty acid desaturase cDNA as a probe. *Bull. Fac. Agr. Saga Univ.* 83: 31-36.
- Yoshida, Y. (1962) Theoretical studies on the methodological procedures of radiation breeding: I. New methods in autogamous plants following seed irradiation. *Euphytica* 11: 95-111.
- Yoshioka, T., Y. Ito and T. Masuda (2001) Induction of mutant resistant to alternaria blotch of apple by gamma-ray irradiation. *Gamma Field Symposia* 39: 69-79.
- Zhou, Y., D. Lu, C. Li., J. Luo, B. Zhu, J. Zhu, Y. Shanguan, Z. Wang, T. Sang, B. Zhou and B. Han (2012) Genetic control of seed shattering in rice by the APETALA2 transcription factor *SHATTERING ABORTION1*. *Plant Cell* 24: 1034-1048.

## Appendix 1. Direct-use mutant cultivars

No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
<b>Rice</b>											
<i>Oryza sativa</i> L.											
1	Reimei	Fuji-minori	1959	Gamma ray 20kR	National Institute of Agricultural Sciences (Hiratsuka)	1966	-	Aomori PAES	Shorter stem; lodging resistance; high-yielding	1138	Futsuhara (1968) <sup>5)</sup>
2	Miyama-nishiki	Takane-nishiki	1972	Gamma ray 30kR	IRB	1978	-	Nagano PAES	High ratio of white core; very suitable for sake brewery	1129	
3	Miyuki-mochi	Toyo-nishiki	1973	Gamma ray 20kR	IRB	1979	-	Nagano PAES	Became glutinous	1131	
4	Shimano-sakigake	Toyo-nishiki	1973	Gamma ray 20kR	IRB	1982	1983	335 Nagano PAES	Big grain; very high-yielding; rice for forage	1141	
5	Shirakabab-nishiki	Reimei	1973	Gamma ray 20kR	IRB	1982	1983	336 Nagano PAES	Big grain; white core; brewery use	1142	
6	Iwate 21	Sasa-nishiki	1979	Gamma ray 20kR	IRB	1986	-	Iwate PAES	Shorter stem; earlier maturity; good taste	1156	Ishikawa <i>et al.</i> (1988) <sup>18)</sup>
7	Iwate 26	Koshihikari	1979	Gamma ray	IRB	1989	-	Iwate PAES	Extremely early maturing; shorter stem; good taste	2445	
8	Hatsu-akane	Sasa-nishiki	1984	Protoplast culture	Mitsui Chemical Ltd.	1988	1990	2508 Mitsui Chemical, Inc.	Shorter stem	2446	
9	Hatsu-yume	Koshi-hikari	1985	Protoplast culture	Plantech Research Institute	1988	1990	2509 Plantech Research Institute	Shorter stem	2447	Ito (1991) <sup>19)</sup>
10	Suzu-takara	Akenohoshi	1984	EMS treatment	Sumitomo Chemical Company	1988	1990	2510 Sumitomo Chemical Co.	Shorter stem; earlier maturity	1165	
11	Sumi-takara	Kogane-bare	1986	Anther culture	Sumitomo Chemical Company	1987	1991	2627 Sumitomo Chemical Co.	No hairs on the leaf blade; smaller seed	2448	
12	Yume-kaori	Tsukino-hikari	1986	Protoplast culture	Plantech Research Institute	1990	1993	3573 Plantech Research Institute	Short stem; lower 1000-kernel weight	2449	Ito (1991) <sup>19)</sup>
13	Fujimi-mochi	Aki-chikara	1988	Gamma ray	IRB	1991	1994	4052 ZEN-NOH	Became glutinous; white seed color	2450	
14	Rinx-Kobayashi	Rinx89	1989	NMU	Plant Laboratory, Kirin Brewery Co., Ltd.	1991	1994	4170 Plant Laboratory, Kirin Brewery Co., Ltd.	Seed-shattering resistance (no seed shattering)	2451	
15	Hareyaka	Sasa-nishiki	1989	Protoplast culture	Plantech Research Institute	1993	1995	4413 Plantech Research Institute	High lodging tolerance	2452	
16	Yume-gokochi	Koshihikari	1986	Protoplast culture	Plantech Research Institute	1992	1995	4709 Plantech Research Institute	Lower amylose	2454	
17	Syonan 3 Gou	Aki-chikara	1988	Radiation		1994	1996	5125 ZEN-NOH	Long and thin grain; high ratio of white core	2453	
18	Meguriai Akane-fuji	Norin No. 22	1980	EMS	Hiroshima PARC	1994	1996	5187 Hiroshima PARC	Short stem; for brewery	2355	Maeshige <i>et al.</i> (1995) <sup>27)</sup>
19		Norin No. 8	1974	EMS	IRB	1994	1997	5307 IRB and Institute of Processing Rice Breeding	Low amylose	2456	

No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
20	Oita 3 Gou	Norin No. 22	1988	Anther culture	Oita PARC	1993	1997	5308	Oita PARC	10% shorter stem; slightly small grain	2457	
21	Milky Queen	Koshihikari	1985	NMU	NARC	1995	1998	6385	NARC	Lower amylose content	2458	Ise <i>et al.</i> (2001) <sup>(6)</sup>
22	Odoroki-mochi	Kanto No. 146	1988	Gamma ray 30kR	IRB	1996	1998	6386	NARC	Became glutinous	2459	Imbe <i>et al.</i> (2004) <sup>(4)</sup>
23	Yume-ekubo	Sasa-nishiki	1988	Protoplast culture	Plantech Research Institute	1996	2000	8546	Plantech Research Institute	Short stem; late maturing	2460	
24	Sakata-mezuru	Mezuru	1988	Gamma ray	IRB	1996	2001	8628	Sakata City	Short stem	2461	
25	Iwata 15 Gou	Kinuhikari	1989	Seed callus	Japan Tobacco Inc.	1996	2001	8632	Japan Tobacco Inc.	Low amylose	2462	
26	Hana-kahori	Yamada-nishiki	1988	Radiation	IRB	1986	2002	9784	NARO	Short stem	2463	
27	LGC-1	Nihonmasari	1988	EI	IRB	2001	2002	10469	IRB	Low glutelin contents in grain	2464	Iida <i>et al.</i> (1993) <sup>(11)</sup>
	Moh-retsu	Rinx89	1989	EMS	Plant Laboratory, Kirin Brewery Co., Ltd.	1988	2003	10961	Plant Laboratory, Kirin Brewery Co., Ltd.	Long stem; tough stem; shattering resistance	2465	
28	Flower Hope	Koshihikari	1985	Gamma ray	IRB	1995	2003	11356	IRB	Low content of globulin in the grain	2466	
30	Kazoku Danran	Nihon-masari	1988	EI	IRB	2000	2004	12176	IRB and Institute of Allergen Free Technology	Low allergen	2467	
31	Minami-yutaka	Moh-retsu	2001	Gamma ray 200Gy	IRB	2004	2007	15001	Miyazaki PAES	Seed-shattering resistance	2886	Kato <i>et al.</i> (2006) <sup>(22)</sup>
32	Homare-fuji	Yamada-nishiki		Gamma ray	IRB	2010	18111	Shizuoka PAES	Semi-dwarf			Shizuoka Prefecture (2010) <sup>(49)</sup>
33	Shizuku-hime	Matsuyama-mitsui		Callus from mature grain	Ehime PAES	2010	19046	Ehime PAES	White center of grain appearing 11-20%			Kanetou <i>et al.</i> (2010) <sup>(21)</sup>
34	Kumika 1 gou	Taichu 65 gou		Anther culture		2010	19048	Kumiai Kagaku and Tohoku Univ.	Herbicide tolerance			
	Yuki-hime-habutae-mochi	Shiga-habutae-mochi		Anther culture of regenerated plant from callus	Shiga PARC	2010	19052	Shiga PARC	Long stem			<a href="http://www.pref.shiga.lg.jp/g/nogyo/saibai/files/yukihime.pdf">http://www.pref.shiga.lg.jp/g/nogyo/saibai/files/yukihime.pdf</a> (in Japanese)
36	Yuki-no-megumi	Yukihikari		Gamma ray	IRB	2010	19410	NARO (Institute of Crop Science)	Large embryo			<a href="https://www.naro.affrc.go.jp/patent/breed/0100/0107/001690.html">https://www.naro.affrc.go.jp/patent/breed/0100/0107/001690.html</a> (in Japanese)
37	Hatsu-yamabuki	Kinuhikari		Gamma ray	IRB	2012	22049	NARO and JIRCAS	Yellow endosperm color; semi-transparent			Kaji <i>et al.</i> (2010) <sup>(20)</sup>
38	Ruri-aoba	Tapoturi		Gamma ray	IRB	2012	21771	NARO	Non or low seed-shattering			Sakai <i>et al.</i> (2013) <sup>(43)</sup>
39	Hoshinoko	Hoshi-no-yume		Gamma ray	IRB	2012	21384	NARO	White colored brown-rice			<a href="https://www.naro.affrc.go.jp/project/resu/Its/laboratory/harc/2008/cryo08-03.html">https://www.naro.affrc.go.jp/project/resu/Its/laboratory/harc/2008/cryo08-03.html</a>
40	Kikusui-HDI gou	Kikusui		Gamma ray	IRB	2012	22079	Keiichi OKAZAKI (Kikusui Shuzou)	Later maturing			
41	EM10	Kinnmaze		NMU	Kyushu University	2014	22986	Kyushu University	Smaller grain; white colored brown-rice			Nishi <i>et al.</i> (2001) <sup>(37)</sup>
42	Amiro-mochi	Kinnmaze		NMU	Kyushu University	2014	22987	Kyushu University	Smaller grain; white colored brown-rice			

No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution Agency	Dev. year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
	Koshihikari Kan 43 1 gou	Koshihikari		Carbon ion beam	Japan Atomic Energy Agency	2015	24338	NIAES and NARO (Institute of Crop Science)	Accumulation of cadmium (Cd) in the grain is less than that of Koshihikari.		
<b>Wheat</b>											
<i>Triticum aestivum</i> L.											
1	Zenkouzi 1 komugi	Iga-Chikugo-Oregon	1958	Gamma ray 30kR	National Institute of Agricultural Sciences (Hiratsuka)	1969	-	- Nagano PAES	Shorter stem (15-20cm lower than the origin); lodging tolerance	1144	
2	Shirowase-komugi	Shirowase-komugi	1966	Gamma ray 25kR	Kyushu University	1977	-	- Kyushu NAES	Erect plant-type; large inflorescence; slightly heavier head type	1143	Nonaka <i>et al.</i> (1979) <sup>38)</sup>
3	Wheat Noh PL.7	Kanto No. 107	1988	EMS	IRB	1994	1999	7169 NARC	Lower amylose content; lower 1,000 kernel weight	2469	Yamaguchi <i>et al.</i> (1998) <sup>56)</sup>
4	Wheat Noh PL.8	Tani-kei A6099	1991	Sodium azide (Na <sub>3</sub> N)	IRB	2000	2003	11365 NARC	Low amylose content	2470	Kiribuchi-Otobe <i>et al.</i> (2001) <sup>24)</sup>
<b>Barley</b>											
<i>Hordeum vulgare</i> L.											
1	Haya-shinriki	Aka-shinriki	1957	Gamma ray 40kR	National Institute of Genetics	1961	-	- National Institute of Genetics	Extremely early maturing; good quality	1118	
2	Gamma 4	Kirin eyoku 1 gou	1958	Gamma ray 15kR	Kanagawa Prefectural Industrial Experiment Station	1962	-	- Barley Research Center; Kirin Brewery Co., Ltd.	Short stem (1.5cm shorter); slightly early maturing (1-2 day earlier)	1116	
3	Amagi Nijo 1 gou	Fuji Nijo	1965	X-ray 20kR	Tokyo University (Faculty of Agriculture)	1971	-	- Barley Research Center; Kirin Brewery Co., Ltd.	Early maturing (7 days earlier); short stem (14cm shorter)	1114	
4	Fuji Nijo II	Fuji Nijo	1967	BUDR(1mM)+ gamma ray 1kR	Tokyo University (Faculty of Agriculture)	1974	-	Barley Research Center; Kirin Brewery Co., Ltd.		1115	
<b>Barnyard millet</b>											
<i>Echinochloa esculenta</i> (A. Braun) H. Scholz											
1	Chojurou-mochi	Nogebie		Gamma ray	IRB	2012	21495	Iwate University	Became glutinous		Hoshino <i>et al.</i> (2010) <sup>10)</sup>
2	Nebarikko 1 gou	Mojappe		Gamma ray	IRB	2012	21577	Iwate Prefecture	Medium stem length; fewer awns; earlier maturity		Nakajo <i>et al.</i> (2013) <sup>35)</sup>
3	Nebarikko 2 gou	Mojappe		Heavy ion beam	Riken	2012	21578	Iwate Prefecture	Shorter stem length; very few awns		Nakajo <i>et al.</i> (2013) <sup>35)</sup>
4	Nebarikko 3 gou	Mojappe		Gamma ray	IRB	2012	21579	Iwate Prefecture	Shorter stem length		Nakajo <i>et al.</i> (2013) <sup>35)</sup>
5	Yume-sakiyo	Nogebie		Gamma ray	IRB	2013	22559	Iwate University	Short stem; earlier maturity		
<b>Job's tear</b>											
<i>Coix ma-yuen</i> Roman.											
1	Hato-musume	Okayama Zairai	1980	Gamma ray 20kR	IRB	1992	1993	3634 Tohoku NAES	Early maturing; short stem	159	Okuyama <i>et al.</i> (1995) <sup>40)</sup>



No.	Cultivar Name	Original Variety	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
2	Hato-hikari	Okayama Zairai	1980	Gamma ray 200Gy	IRB	1995	1996	5069	Tohoku NAES	Early maturing; short stem	2471	Ishida <i>et al.</i> (1995) <sup>17)</sup>
<b>Buckwheat</b>												
<i>Fagopyrum esculentum</i> Moench												
1	Dewa-kaori	Mogami Wase	1988	Colchicine	Yamagata PAES	1995	1996	5069	Yamagata PAES	Larger grain	2472	
2	Hokkai 3 Gou	Botan-soba	1979	Colchicine	NARC for Hokkaido Region	2001	2005	12967	NARO	Tetraploid; large seed	2473	
3	Gamma-no-irodori	Botan-soba		Gamma ray	IRB	2005	22872		IRB, NIAS	Higher rutin content in the seed, smaller seed		
4	Cobalt-no-chikara	Botan-soba		Gamma ray	IRB	2013	22873		IRB, NIAS	Smaller seed		
5	Ruchiking	Botan-soba		Gamma ray (repeated irradiation for the generations)	IRB	2013	22874		IRB, NIAS	Very high rutin content in the seed; smaller seed		
<b>Tartary buckwheat</b> <i>Fagopyrum tataricum</i> (L.) Gaertn.												
1	Daizen	No name		Colchicine	Tokita Seed Company	2010	19044					
2	Hokkai T9 gou	Dattan		Colchicine	NARO	2010	19466		NARO			Suzuki <i>et al.</i> (2008) <sup>11)</sup>
3	Hokkai T10 gou	Hokkai T8 gou		Ethyl methane sulfonate (EMS)	NARO	2010	19526		NARO	Pale red flower color; less maturing seed		Suzuki <i>et al.</i> (2008) <sup>11)</sup>
4	Shinano Kurotsubu	Bhate Parpar		Colchicine	Shinshu University	2010	19527		Shinshu University			
5	Daruma Dattan	Hokkai T8		Gamma ray	IRB	2013	22633		NIAS (IRB)	Semi-dwarf		
6	Ion-no-kousai	Routundatum		Helium ion beam	JAEA, Takasaki	2013	22634		NIAS (IRB) and JAEA	Shorter stem; light leaf		
<b>Amaranthus</b> <i>Amaranthus</i> L.												
1	New Aztec	Mexico-kei	1989	Gamma ray 50kR	IRB	2001	2006	14399	NARO (National Institute of Crop Science)	Early maturing; short stem	2474	Katsuta <i>et al.</i> (2001) <sup>23)</sup>
<b>Rapeseed</b> <i>Brassica napus</i> L.												
1	Haya-natane	Michinoku-natane	1955	Colchicine 0.03× 24h	Fukuoka PAES	1961			Fukuoka PAES	Maturing 3 days earlier than the original variety (earliest of all varieties)	1720	
2	Hanakkori	Wase-kei Saishin×Jyo-	1990	Colchicine treatment	Yamaguchi PAES	1995	1999	7324	Yamaguchi PAES	Morphology	2475	
<b>Soybean</b> <i>Glycine max</i> (L.) Merr.												
1	Raiden	Nemashirazu	1960	Gamma ray 10kR	National Institute of Agricultural Sciences (Hiratsuka)	1966			Tohoku NAES	Earlier maturity (25 days); shorter stem lodging resistance	1594	Laboratory of Soybean Breeding (1970) <sup>26)</sup>

No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
2	Raiko	Nemashirazu	1960	Gamma ray 10kR	National Institute of Agricultural Sciences (Hiratsuka)	1969			Tohoku NAES	Earlier maturity (15 days); shorter stem lodging resistance	1595	Laboratory of Soybean Breeding (1970) <sup>26)</sup>
3	Wase-suzunari	Oku-shirome	1975	Gamma ray 10kR	IRB	1983	1984	610	Tohoku NAES	Earlier maturity (extremely early)	1597	Hashimoto <i>et al.</i> (1985) <sup>8)</sup>
4	Mura-yutaka	Fuku-yutaka	1984	X-ray 24kR	Saga Univ. Facul. Agric.	1988	1990	2156	Saga PAES	From brown hilum to white hilum	2476	Nakamura <i>et al.</i> (1991) <sup>36)</sup>
5	Kosuzu	Natto-Kotsubu	1979	Gamma ray 10kR	IRB	1987	1990	2397	Tohoku NAES	Earlier maturity (medium); lodging resistance	1596	Hashimoto <i>et al.</i> (1988) <sup>7)</sup>
6	Ryokusui	Fukura	1984	Gamma ray 20kR	IRB	1988	1980	2516	Iwate PAES	Later maturity; brown seed color	1598	
7	Saotome	Aji-ichiban	1986	Gamma ray 15kR	Tokyo Metropolitan Isotope General Research Institute	1991	1994	4055	Mitsui Chemical Ltd.	Yellow seed color; higher pod density	2477	
8	Kokoro-zukushi	Aji-ichiban	1986	Gamma ray 15kR	Tokyo Metropolitan Isotope General Research Institute	1991	1994	4115	Mitsui Chemical Ltd.	Earlier maturity; higher pod density	2478	
9	Murasaki-zukin	Kyoto No. 1 (Shin-Tanba)	1975	Gamma ray 10Gy	Kyoto Univ. Facul. Agric.	1992	1995	4715	Kyoto PARI	Later maturity; for green soybean-use	2479	
10	Ichi-hime	(Kanto No. 2/Kanto No. 1)F2	1989	Gamma ray 15kR	IRB	1996	1997	5369	Kyushu NAES	All lipoxygenase knock-out	2480	Hajika <i>et al.</i> (2002) <sup>6)</sup>
11	Taki-hime	Komaki Dadacha	1988	Gamma ray 8kR	IRB	1996	2000	8365	Yanagawa Saisyu Kenkyukai	Shorter stem; high pod density; earlier maturity	2481	
12	Aomori-toyomaru	Kemame (Aomori Local var.)	1989	Gamma ray	IRB	1995	2001	8642	Aomori Prefectural Agriculture and Forestry Research Center	Earlier maturity; shorter stem; ecological mutation (summer-type)	2482	
13	Aomori-fukumaru	Kemame (Aomori Local var.)	1989	Gamma ray	IRB	1995	2001	8643	Aomori Prefectural Agriculture and Forestry Research Center	Earlier maturity; shorter stem; ecological mutation (summer-type)	2483	
14	Akita-midori	Aome-daizu	1990	Gamma ray 10kR	IRB	1998	2002	9650	Akita PAES	Earlier maturity	2484	Sasaki <i>et al.</i> (2000) <sup>46)</sup>
15	Yume-minori	Kari-kei No.434	1991	Gamma ray	IRB	2001	2004	12280	Tohoku NAES	Low-allergic	2485	Takahashi <i>et al.</i> (2004) <sup>52)</sup>
16	Sayane	Saya-musume	1995	Gamma ray	IRB	2002	2006	13749	Snow Brand Ltd.	Earlier maturity; shorter main stem length	2487	
17	Akita-honoka	Hidden		Somaclonal variation from somatic embryo	Akita PAES	2015	24350		Akita PAES	Earlier maturity; larger bean size		Sato <i>et al.</i> (2015) <sup>47)</sup>

No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
<b>Adzuki bean</b>												
1	Beni-nambu	Mombetsu	1968	Gamma ray 10kR	IRB	1978			Iwate PAES	Earlier maturity (10 days)	406	Sato <i>et al.</i> (1982) <sup>48)</sup>
<b>Broad bean</b>												
		<i>Vicia faba</i> L.										
1	Rin-rei	Niigata Zairai one-bean type	1982-8	Gamma ray irradiation (recurrent irradiation)	Tokyo Metropolitan Isotope General Research Institute	1988	1991	2695	Hokusei Nouji Co.	Dwarf	2490	
<b>Sweet potato</b>												
		<i>Ipomoea batatas</i> (L.) Lam.										
1	Sweet Garden	CNI367-2	1990	Gamma-ray irradiation to tissue culture	Kagoshima Biotechnology Institute	1993	1998	6390	Miwa Green Co.	Leaf-shape	2491	
<b>Potato</b>												
		<i>Solanum tuberosum</i> L.										
1	White Baron	Danshaku	1989	Protoplast culture	Hokuren	1985	1997	5965	Hokuren	Less browning after cutting	2492	
2	Jagakids Purple	Neo Delicious	1986	Protoplast culture	Plant Laboratory, Kirin Brewery Co., Ltd.	1990	1994	4054	Plant Laboratory, Kirin Brewery Co., Ltd.		2493	
<b>Sugarcane</b>												
		<i>Saccharum officinarum</i> L.										
1	Nan-ci	N11	1976	Gamma ray to the plant	IRB	1985	1986	1024	Kagoshima University; Faculty of Agriculture	Larger plant size; High yielding	375	
<b>Chinese matgrass</b>												
		<i>Cyperus malaccensis</i> Lam. ssp. <i>monophyllus</i> (Vahl) T. Koyama										
1	Toyo-midori	Ohoi 2 Gou	1969	Gamma ray to plant 115kR	IRB	1979			Ooita Prefectural Igyou Shidousyo	Thicker stem; lodging tolerance; resistance to fungal disease	373	
<b>Rush</b>												
		<i>Juncus effusus</i> L. var. <i>decipiens</i> Buchenau										
1	Seto-nami	Asa-nagi	1963	Gamma ray to plant 68kR	IRB	1982	1983	397	Hiroshima PAES, Rush Breeding Station	More longer stems; High yielding; good quality; beautiful straws	345	Sadahira <i>et al.</i> (1982) <sup>41)</sup>
2	Fuku-nami	Asa-nagi	1965	Gamma ray to plant 96kR	IRB	1984	1986	941	Hiroshima PAES, Rush Breeding Station	Early-harvest and high yielding; less browning of leaf tops; stronger straw	346	Sadahira <i>et al.</i> (1984) <sup>42)</sup>
3	Chikugo-midori	Iso-nami	1989	Meristem culture	Fukuoka ARC	1997	2001	9033	Fukuoka ARC	More long stem number; higher weight of total long stems	2494	Nakahara <i>et al.</i> (1998) <sup>34)</sup>
<b>Lettuce</b>												
		<i>Lactuca sativa</i> L.										
1	Evergreen	Way-a-head	1959	<sup>32</sup> P100μCi and gamma ray 42kR	Hyogo PAES	1966			Hyogo PAES	Later bolting under high summer temperature	2219	

No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
2	Giant Green	Way-a-head	1959	<sup>32</sup> p	Hyogo PAES	1966	1995	4778	Hyogo PAES	More heat tolerance	2220	
3	Satilo	Empire	1990	EMS	Sumitomo Chemical company	1992	1995	4778	Sumitomo Chemical Co.	Shrunken leaf; wave in leaf edge	2495	
<b>Japanese parsley</b> <i>Oenanthe javanica</i> (Blume) DC.												
1	Miyagi VWD 1 Gou	Shimane-midori	1994	Protoplast culture	Miyagi Prefectural Agriculture and Horticulture Research Center	1996	2002	10071	Miyagi Prefectural Agriculture and Horticulture Research Center	Resistant to leaf disease	3163	
<b>Butterbur</b> <i>Petasites japonicus</i> (Siebold & Zucc.) Maxim.												
1	Okayama Nohshi B1 Gou	Aichi-wase	1985	Callus (meristem)	Okayama PAES	1992	1996	5017	Okayama PAES	Less anthocyanin in leaf stem	3164	Hikawa <i>et al.</i> (1998) <sup>9)</sup>
<b>Tomato</b> <i>Solanum lycopersicum</i> L.												
1	Kyoryoku-reikou 1	Syogyoku	1967	Gamma ray irradiation to pollen 2340R	IRB	1974			Musashi Breeding Station Co.	Resistant to TMV and wilt disease	2221	
<b>Strawberry</b> <i>Fragaria</i> L.												
1	Shin-Nyohou	Nyohou	1963	Callus formation	Tochigi PAES, Tochigi Branch	1989	1989	2048	Tochigi PAES, Tochigi Branch	Earlier harvest type; high yielding	3213	
2	Akita Berry	Morioka 16 Gou	1967	Meristem culture	Akita Prefectural College of Agriculture	1990	1992	3224	Akita Prefectural College of Agriculture	Resistant to black leaf spot disease ( <i>Alternaria alternata</i> )	3214	
3	Anther	Nyohou	1986	Anther culture	Ibaraki PAES	1991	1994	4176	Ibaraki PAES	Early maturing	3215	Ezura <i>et al.</i> (1998) <sup>2)</sup>
4	Himatsuri	Toyonoka	1988	Meristem culture	Kyushu Tokai University	1992	1995	4593	Kyushu Tokai University	Bright red fruit skin; light red flesh color	3216	Fukuoka <i>et al.</i> (1996) <sup>3)</sup>
5	Smile Heart	Ai-berry	1993	Tissue culture	Shikoku Research Institute Inc.	1994	1998	6563	Shikoku Research Institute Inc.	Resistant to phytophthora rot ( <i>Phytophthora nicotinae</i> )	3217	
<b>Burdock</b> <i>Atractium lappa</i> L.												
1	Cobalt Wase	Yanagawa-risou	1969	Gamma ray 10kR	IRB	1978	1981	73	Yanaken Co.	Earlier root growth; earlier maturing (growing days: 120 days)	2218	
2	Cobalt Goku-wase	Yanagawa-nakate	1969	Gamma ray 10kR	IRB	1978	1981	74	Yanaken Co.	Shorter root; extremely early maturing (growing days 100 days)	2216	
3	Cobalt Okute	Yanagawa-nakate	1969	Gamma ray 10kR	IRB	1978	1981	75	Yanaken Co.	Less pithy tissue; higher storability; late maturing	2217	
4	Tsune-yutaka	Yanagawa-risou	1969	Gamma ray 10kR	IRB	1984	1986	1233	Yanaken Co.	Thicker root; thick at upper root	2225	

No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
5	Tegaru	Cobalt Goku-wase	1982	Gamma ray 10kR	Tokyo Metropolitan Isotope General Research Institute	1991	1995	4422	Yanaken Co.	Shorter root; thick at upper root	2496	
<b>Taro</b>												
<i>Colocasia esculenta</i> Schott												
1	Fukugashira	Yatsugashira	1984	MNU	Saga University, Faculty of Agriculture	1989	1992	3172	Saga PAES	Round mother corm; higher cooking and processing suitability	2497	
2	Chiba-manu	Dotare	1995	Soft X-ray	Chiba PAES	2004	2007	15137	Chiba PAES	Tuber shape	2901	Suzuki <i>et al.</i> (2006) <sup>50)</sup>
3	Ehime-Noushi V2 gou	Onna-wase	1994	Callus from meristem part	Ehime PAES	2000	2008	16024	Ehime PAES	Number and size of grand-son tuber is larger		Nakagawa <i>et al.</i> (2015) <sup>33)</sup>
<b>Wasabi (Japanese horseradish)</b>												
<i>Wasabia japonica</i> (Miq.) Matsum.												
1	Green Magic	Matsuma	1991	Meristem culture	Miyoshi Co.	1993	1997	5312	Miyoshi Co.	Longer stem; less anthocyanin at leaf stem; light color at root stem; stronger bitterness		
2	Amagi-nishiki	Matsuma	1988	Tissue culture	Miyoshi Co.	1991	1998	6030	Miyoshi Co.	Less pithy tissue	3219	
<b>Scarlet runner bean</b>												
<i>Phaseolus coccineus</i> L.												
1	Shiro-hanakko	Oo-shirohana	1991	Gamma ray		2002	2002	14409	Hokkaido Prefectural Central Agriculture Experiment Station	Higher pod width		Minami, M. and H. Sato (2009) <sup>29)</sup>
<b>Limonium</b>												
<i>Limonium</i> Mill.												
1	Tall Pink Emirre	Tall Emirre	1962	Gamma ray	Tokyo Metropolitan Isotope General Research Institute	1993	1998	6100	Miyoshi Co.	Petal color: pale pink	3181	
2	Neo Misty Blue	Misty Blue	1990	Gamma ray	Tokyo Metropolitan Isotope General Research Institute	1994	1998	6101	Miyoshi Co.	Less number of floral stalks	3182	
3	Dai-fura	Dai-fura Pink	1992	Gamma ray irradiation to tissue culture	Tokyo Metropolitan Isotope General Research Institute	1995	1999	7104	Daiichi Seed Co., Ltd.	Sepal color: bright reddish purple	3183	
4	Dai-fura Super	Dai-fura Pink	1992	Gamma ray irradiation to tissue culture	Tokyo Metropolitan Isotope General Research Institute	1995	2000	7938	Daiichi Seed Co., Ltd.	Sepal color: bright purple; less number of floral stalks	3184	
5	Dai-lady Rose	Super Lady	1991	Gamma ray irradiation to tissue culture	Tokyo Metropolitan Isotope General Research Institute	1996	2000	8490	Daiichi Seed Co., Ltd.	Petal color: vivid yellowish green	3185	
6	Dai-lady White	Super Lady	1991	Gamma ray irradiation to tissue culture	Tokyo Metropolitan Isotope General Research Institute	1996	2001	8971	Daiichi Seed Co., Ltd.	Sepal color: pinkish white	3186	

No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
<b>Showy baby's breath</b> <i>Gypsophylla elegans</i> M. Bieb.											
1	Kishu Kasumi 1 Gou	Perfecta	1989	Meristem culture	Wakayama Research Center of Agriculture, Forestry and Fisheries Tokyo Metropolitan Isotope General Research Institute	1994	2000	7762 Wakayama Research Center of Agriculture, Forestry and Fisheries Miyoshi Co.	More wax on the leaf surface; top of the petal, V-shape Sepal shape	3220	Miyamoto <i>et al.</i> (2003) <sup>30)</sup>
2	Buranka	Diamond	1993	Gamma ray		1997	2002	9667 Miyoshi Co.		3165	
3	Ayami	Bristle Fairy	1993	Colchicine	Sumitomo Chemical company	1993	1999	7094 Sumitomo Chemical Company	Shorter internode length; more petal number	3166	
<b>Gymnaster</b> <i>Aster savatieri</i> Makino											
1	Seto-no-otome	Hama-otome	1991	Gamma ray irradiation to meristem culture	IRB	1994	1998	6896 Minoru Industrial Co., Ltd.	More ligulose flowers; deep flower color	3167	
2	Minoru-otome	Hama-otome	1990	Gamma ray	IRB	1995	2000	7831 Minoru Industrial Co., Ltd.	Flower shape; flower color: bright reddish purple	3168	
<b>Prairie gentian</b> <i>Eustoma grandiflorum</i> (Raf.) Shimmers											
1	Purple Robin	Pastel Murasaki	1992	Gamma ray	IRB	1995	1999	7266 IRB and Naganoken Nourin	Small-flowered; spray type	2004	
2	Red Robin	Morugen Rhoto	1992	Gamma ray	IRB	1995	1999	7267 IRB and Naganoken Nourin	Small-flowered; spray type	2006	
3	Tokyo El gou	Crystal Yellow		Gamma ray	Tokyo Metropolitan Agriculture and Forestry Research Center	2011	2015	24321 Tokyo Metropolitan Agriculture and Forestry Research Center			Miyashita <i>et al.</i> (2014) <sup>31)</sup>
4	Izu-Ohshima E3 gou	Tenryu Otome		Gamma ray	Tokyo Metropolitan Agriculture and Forestry Research Center	2005	2009	18330 Tokyo Metropolitan Agriculture and Forestry Research Center			Miyashita <i>et al.</i> (2014) <sup>31)</sup>
<b>Carnation</b> <i>Dianthus carvophyllus</i> L.											
1	Scarlet Bell	Angel	1977	Gamma ray 31kR	Tokyo Metropolitan Isotope General Research Institute	1980	1983	379 Miyoshi Co.	Flower color: from deep pink to red	53	
2	Mrs. Elegant	SP	1996	Tissue culture	Plant Laboratory, Kirin Brewery Co., Ltd.	1997	2001	9247 Plant Laboratory, Kirin Brewery Co., Ltd.	Small flowered; smaller number of flowers per flower stalk	2498	
3	Boh-red	Nora	1995	Meristem culture	Akita Prefectural Agriculture, Forestry and Fisheries	1998	2002	10550 Akita Prefectural Agriculture, Forestry and Fisheries	Flower color: vivid red	2499	

No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
4	Kirikami Red (Mrs. Red)	Mrs. Elegant	1997	Tissue culture	Plant Laboratory, Kirin Brewery Co., Ltd.	1999	2002	10926	Plant Laboratory, Kirin Brewery Co., Ltd.	Surface of flower: deep orange red	2500	
5	Garden Spice carnation Salmon	Garden Spice carnation Cool Pink	1998	X-ray to tissue culture	Plant Laboratory, Kirin Brewery Co., Ltd.	1999	2004	12103	Plant Laboratory, Kirin Brewery Co., Ltd.	Flower surface: vivid pink	3169	
6	Garden Spice carnation Pink	Garden Spice carnation Cool Pink	1998	X-ray to tissue culture	Plant Laboratory, Kirin Brewery Co., Ltd.	1999	2004	12103	Plant Laboratory, Kirin Brewery Co., Ltd.	Flower color: pale pink	2502	
7	Your Red	Nora	1995	Meristem culture	Akita Prefectural Agriculture, Forestry and Fisheries	2000	2005	13051	Akita Prefectural Agriculture, Forestry and Fisheries	Flower color: vivid red; suspended flower; large flower	2503	Arai (2001) <sup>1)</sup>
8	Garden Spice carnation Marble	Garden Spice carnation pearl White Marble	1995	X-ray to tissue culture	Plant Laboratory, Kirin Brewery Co., Ltd.	2000	2005	13052	Plant Laboratory, Kirin Brewery Co., Ltd.	Flower color: pinkish white with vivid purple red stripe	2504	
9	Misty Pink Ion	Vital	2000	C ion beam on regenerating callus	JAEA, Takasaki	2002	2005	13585	Plant Laboratory, Brewery Co., Ltd.	Flower color; petal shape	2505	
10	Dark Pink Ion	Vital	2000	C ion beam on regenerating callus	JAEA, Takasaki	2002	2005	13586	Plant Laboratory, Brewery Co., Ltd.	Flower color; petal shape	2506	
11	Red Vital Ion	Vital	2000	C ion beam on regenerating callus	JAEA, Takasaki	2002	2005	13587	Plant Laboratory, Brewery Co., Ltd.	Flower color: bright red	2507	
12	Red Vital	Vital	2001	C ion beam on leaf	JAEA, Takasaki	2004	2008	16064	Barbre & Branc S. A	Double flower with bright red color		Okamura <i>et al.</i> (2006) <sup>39)</sup>
13	Beam Cherry	Vital		C ion beam on leaf	JAEA, Takasaki	2009	2009	17297	Barbre & Branc S. A	Flower color		
14	Loro Red	Loro	2007	Gamma ray	IRB	2007	2010	19121	Miyazaki Prefecture	Early flowering; flower color		<a href="http://www.pref.miyazaki.lg.jp/contents/org/nosei/mae-station/topics/21.html">http://www.pref.miyazaki.lg.jp/contents/org/nosei/mae-station/topics/21.html</a>
15	Loro Pink	Loro	2007	Gamma ray	IRB	2007	2010	19122	Miyazaki Prefecture			<a href="http://www.pref.miyazaki.lg.jp/contents/org/nosei/mae-station/topics/21.html">http://www.pref.miyazaki.lg.jp/contents/org/nosei/mae-station/topics/21.html</a>
<b>Clematis</b>												
<i>Clematis</i> L.												
1	Pastorale	Hakuba	1986	Gamma ray 2kR	IRB	1989	1992	3130	Mr. Fukutarou MIYATA	Flower color; flower shape	3170	
2	Nachtmusik	Hakuba	1986	Gamma ray 2kR	IRB	1989	1992	3131	Mr. Fukutarou MIYATA	Flower color; flower shape (double flowered)	3171	

No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
<b>Chrysanthemum</b> <i>Chrysanthemum x morifolium</i> Ramat.												
1	Ki-uzushio	Uzushio	1962	Gamma ray 2.5kR	IRB	1983	1986	1044	Seikoen Ltd.	Flower color: from white to yellow		1987
2	Baigiku Rainbow White	Seikou-no-kurenai	1983	Gamma ray	IRB	1985			Seikoen Ltd.	Flower color: from red to white		1991
3	Baigiku Rainbow Yellow	Seikou-no-kurenai	1983	Gamma ray	IRB	1985			Seikoen Ltd.	Flower color: from red to yellow		1993
4	Baigiku Rainbow Peach	Seikou-no-kurenai	1983	Gamma ray	IRB	1985			Seikoen Ltd.	Flower color: from red to rose color		1988
5	Baigiku Rainbow Pink	Seikou-no-kurenai	1983	Gamma ray	IRB	1985			Seikoen Ltd.	Flower color: from red to pink		1990
6	Baigiku Rainbow Orange	Seikou-no-kurenai	1983	Gamma ray	IRB	1985			Seikoen Ltd.	Flower color: from red to reddish yellow		1989
7	Pink Orizuru	Orizuru	1986	Gamma ray	IRB	1988	1990	2537	Seikoen Ltd.	Flower color: from white to pink		1997
8	Hae-no-hatsuyuki	Taihei	1987	Gamma ray to tissue culture	IRB	1990	1995	4305	IRB and Okinawa PAES	Flower color: yellowish white		2508
9	Hae-no-kirameki	Taihei	1987	Gamma ray to tissue culture	IRB	1990	1995	4306	IRB and Okinawa PAES	Flower color: yellowish orange		2512
10	Hae-no-kurenai	Taihei	1987	Gamma ray to tissue culture	IRB	1990	1995	4307	IRB and Okinawa PAES	Flower color: purple pink		2509
11	Hae-no-miyarabi	Taihei	1987	Gamma ray to tissue culture	IRB	1990	1995	4308	IRB and Okinawa PAES	Flower color: pale yellowish pink		2513
12	Hae-no-yugure	Taihei	1987	Gamma ray to tissue culture	IRB	1990	1995	4309	IRB and Okinawa PAES	Flower color: pink		2514
13	Hae-no-kagayaki	Taihei	1987	Gamma ray to tissue culture	IRB	1990	1995	4310	IRB and Okinawa PAES	Flower color: bright yellow		2515
14	Iero Sei Roza (Reagan Yellow)	Sei Roza	1989	Gamma ray 2kR	IRB	1992	1997	5398	Seikoen Ltd.	Flower color: bright greenish yellow		2516
15	Oreiji Sei Roza (Reagan Orange)	Sei Roza	1989	Gamma ray 2kR	IRB	1992	1997	5399	Seikoen Ltd.	Flower color: orange		2517
16	Dipu Sei Roza (Reagan Royal)	Sei Roza	1989	Gamma ray 2kR	IRB	1992	1997	5400	Seikoen Ltd.	Flower color: deep purplish pink		2718
17	Paru Sei Roza (Reagan Pearl)	Sei Roza	1989	Gamma ray 2kR	IRB	1992	1997	5401	Seikoen Ltd.	Flower color: vivid purplish pink		2519
18	Howaito Sei Roza (Reagan White)	Sei Roza	1989	Gamma ray 2kR	IRB	1992	1997	5402	Seikoen Ltd.	Flower color: yellowish white		2520
19	Yellow Prism	Goldstock Bar	1991	Gamma ray to tissue culture	Plant Laboratory, Kirin Brewery Co., Ltd.	1997	1997	5874	Plant Laboratory, Kirin Brewery Co., Ltd.	Erect flower petal; petal color: bright yellow		2521



No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
20	Pearl Prism	Goldstock	1992	Tissue culture	Plant Laboratory, Kirin Brewery Co., Ltd.	1993	1997	5875	Plant Laboratory, Kirin Brewery Co., Ltd.	Erect flower petal; more transparent petal	2522	
21	Amazon	Bred line	1992	Gamma ray to tissue culture	Plant Laboratory, Kirin Brewery Co., Ltd.	1998	1998	6167	Plant Laboratory, Kirin Brewery Co., Ltd.	Petal color: deep orange red	2523	
22	Joy Light	Bred line	1992	Gamma ray	IRB	1998	1998	6218	ZEN-NOH		2524	
23	Joy Coral	Bred line	1992	Gamma ray	IRB	1998	1998	6219	ZEN-NOH		2525	
24	Joy Royal	Bred line	1993	Gamma ray	IRB	1998	1998	6220	ZEN-NOH		2526	
25	Joy Apricot	Bred line	1993	Gamma ray	IRB	1998	1998	6221	ZEN-NOH		2527	
26	Royal Wedding Rose	Goldstock	1992	Tissue culture	Plant Laboratory, Kirin Brewery Co., Ltd.	1998	1998	6638	Plant Laboratory, Kirin Brewery Co., Ltd.	Flower color: deep purple pink	2528	
27	Joy Light Yellow	Joy Light Salmon	1994	Gamma ray	IRB	1994	1999	7096	ZEN-NOH	Flower color: pale yellow	2529	
28	White Lineker OW-1	Goldstock Lineker	1995	X ray to tissue culture	Plant Laboratory, Kirin Brewery Co., Ltd.	2000	1998	8318	Plant Laboratory, Kirin Brewery Co., Ltd.	Flower color: white	2530	
29	Joy Prelude Apricot	Joy prelude Pink	1995	Gamma ray	IRB	1996	2000	8477	ZEN-NOH		2531	
30	Joy Prelude Coral	Joy prelude Pink	1995	Gamma ray	IRB	1996	2000	8479	ZEN-NOH		2532	
31	Etenraku	Kotobuki	1992	Tissue culture	Yamagata Prefectural Horticulture Experiment Station	1997	2001	8867	Yamagata Prefectural Horticulture Experiment Station	Shorter stem	2533	
32	Pretty Wedding	Royal Wedding	1996	X ray to tissue culture	Plant Laboratory, Kirin Brewery Co., Ltd.	2001	2001	8878	Plant Laboratory, Kirin Brewery Co., Ltd.	Flower color (bi-color; outside: pink; inside: white)	2534	
33	Remukoiero (Remco Yellow)	Remco	1995	Radiation	Kinki University	1998	2001	9570	Seikoen Ltd.		2535	
34	Ray Sunrise	Erias	1991	Soft X ray	Shizuoka PAES	1997	2002	10538	Shizuoka PAES	Petal color (abaxial: orange; adaxial: yellow)	2536	Yamada <i>et al.</i> (1999) <sup>54)</sup>
35	Dreaming	Dream Nurse	1993	Soft X ray 200Gy	Shizuoka PAES	2000	2004	12420	Shizuoka PAES	Flower color: from white to yellow	2537	Yamada <i>et al.</i> (2002) <sup>55)</sup>
36	Princess Kagawa	Pink Seiko	1997	Tissue culture + X ray	Kagawa PAES	2001	2004	12415	Kagawa PAES	Stem color and petal color: reddish purple	2538	Furuichi <i>et al.</i> (2003) <sup>4)</sup>
37	Emi-akari	Sei-un	1997	Soft X ray irradiation to meristem culture	Aomori Prefectural Agriculture and Forestry Research Center	2003	2006	14099	Aomori Prefectural Agriculture and Forestry Research Center	Ligulate petal color: pale greenish yellow		



No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
47	Ion-no-kouki	Taihei		Callus generated from petals was irradiated with a carbon ion beam	JAEA, Takasaki	2002	2006	14570	IRB and JAEA, Takasaki	Petal color	3399	
48	Ion-no-mahou	Taihei		Callus generated from petals was irradiated with a neon ion beam	JAEA, Takasaki	2002	2006	14571	IRB and JAEA, Takasaki	Petal color	3400	
49	Sei-faust-cream	Sei-faust		Gamma ray	IRB	2007	2010	19333	Seikoen Ltd.	Petal color: 155A		
50	Sei-piaje-white	Piaje		Gamma ray	IRB	2007	2010	19185	Seikoen Ltd.	Petal shape		
51	Sei-piaje-pink	Piaje		Gamma ray	IRB	2007	2010	19186	Seikoen Ltd.	Petal color: 187C		
52	Sei-piaje-yellow	Piaje		Gamma ray	IRB	2007	2010	19187	Seikoen Ltd.	Petal color: 9A		
53	Sei-rocket-yellow	A developed line		Gamma ray	IRB	2007	2010	19335	Seikoen Ltd.	Flower disc color: 144B		
54	Sei-patrick-yellow	Sei-patrick		Gamma ray	IRB	2007	2010	19336	Seikoen Ltd.	Flower color (182A); shape (flower disc is smaller)		
55	Sei-patrick-sermon	Sei-patrick		Gamma ray	IRB	2007	2010	19337	Seikoen Ltd.	Flower color: 49D		
56	Sei-roys-yellow	Sei-bingo		Gamma ray	IRB	2007	2010	19003	Seikoen Ltd.	Flower color: 4A		
57	Arajin 2	(Aladdin Arajin)		Ion beam	JAEA, Takasaki	2007	2010	19096	Kagoshima PAES	Fewer flower buds		Ueno <i>et al.</i> (2013) <sup>53)</sup>
58	Lemon smile	Egao		Soft X-ray	Aomori Prefectural Agriculture and Forestry Research Center	2011	2013	22281	Aomori Prefectural Agriculture and Forestry Research Center and Seikoen Ltd.	Flower color (cream)		
59	Akebono-no-mai	Taihou-senryu		Soft X-ray irradiation to shoot-apex culture	Aomori Prefectural Agriculture and Forestry Research Center	2014	2016	24889	Aomori Prefectural Agriculture and Forestry Research Center and Seikoen Ltd.			
60	Akane-no-mai	Taihou-senryu		Soft X-ray irradiation to shoot-apex culture	Aomori Prefectural Agriculture and Forestry Research Center	2014	2016	24890	Aomori Prefectural Agriculture and Forestry Research Center and Seikoen Ltd.			
1	Rosetone	Rohito-horun	1987	Gamma ray	Tokyo Metropolitan Isotope General Research Institute	1992	1998	6608	Hanano-Yamato Co.	Leaf shape: ellipse	3172	

Lily

Lilium L.

No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
2	Light Memory	Rohto-horun	1987	Gamma ray	Tokyo Metropolitan Isotope General Research Institute	1992	1998	6609	Hanano-Yamato Co.	Petal color: pale pink; deep purple stripe in the center rib of petal	3173	
3	Ivory Memory	Rohto-horun	1988	Gamma ray to tissue culture	Tokyo Metropolitan Isotope General Research Institute	1993	1998	6610	Hanano-Yamato Co.	Petal color: pale orange yellow	3174	
4	Coral Bouquet	Rohto-horun	1987	Gamma ray	Tokyo Metropolitan Isotope General Research Institute	1993	1999	7544	Hanano-Yamato Co.	No pollen production; smaller stem, umbel	3175	
<b>Cymbidium</b>												
<i>Cymbidium Sw.</i>												
1	Cocktail Dress	Inasa	1987	Tissue culture (mericlone)	Purchase	1992	1997	5707	Mr. Makoto MATSUO	Color and shape of petal	3221	
<b>Kalanchoe</b>												
<i>Kalanchoe Adans.</i>												
1	Harvest Moon	<i>K. miniata/K. polyphylloukar icus</i>	1985	Gamma ray	IRB	1992	1996	5148	Mr. Isao KOBAYASHI	Flower shape	2542	
<b>Caladium</b>												
<i>Caladium Vent.</i>												
1	Soyogi	Candydam	1991	X ray	Kagoshima PAES	1993	2000	8250	Kagoshima PAES		3176	
<b>Petunia</b>												
<i>Petunia Juss.</i>												
1	Kirimaji Cherry Red (Purple Wave)	Kirimaji	1998	X-ray to protoplast culture	Plant Laboratory, Kirin Brewery Co., Ltd.	2000	2004	12082	Plant Laboratory, Kirin Brewery Co., Ltd.	Monotone flower color: vivid red	3177	
2	Sun-lobein	Revolution		Heavy ion beam irradiation to callus developed from axillary buds	Riken	2001	2006	13913	Suntory Flowers Ltd and Keisei Rose Co.	Flower color: JHS color chart 9507		
3	Seto-fuku-white- A bariegata	A developed line		Ion beam	The Wakasawan Energy Research Center	2007	2009	17653	Hakusan International Co. and The Wakasawan Energy Research Center	Flower color and plant type		
4	Hamapetu 3 gou	Calen Beach		X-ray	Shizuoka PAES	2007	2010	18926	The Yokohama nursery Co.	Flower color: 76C		
5	Sunsurfsitrou	An original line		Radiation		2007	2014	23217	Suntory Flowers Ltd.			
<b>Begonia</b>												
<i>Begonia L.</i>												
1	Ryoku-ha	Winter Queen	1966	Gamma ray 10kR	Tokyo Metropolitan Isotope General Research Institute	1976	-	-	Tokyo Metropolitan Industrial Technology Research Institute	Leaf color; variation in leaf spot; higher ozone/oxidant sensitivity	1985	

No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
2	Ginsei	Winter Queen	1966	Gamma ray 10kR	Tokyo Metropolitan Isotope General Research Institute	1976	-	-	Tokyo Metropolitan Industrial Technology Research Institute	Leaf color; variation in leaf spot	1981	
3	Orange Iron	Iron Cross	1971	Gamma ray 0.5kR	Tokyo Metropolitan Isotope General Research Institute	1976	-	-	Tokyo Metropolitan Industrial Technology Research Institute	Leaf color; leaf shape; brown purple iron-cruciate leaf spot	1984	
4	Mini-mini Iron	Iron Cross	1971	Gamma ray 1kR	Tokyo Metropolitan Isotope General Research Institute	1976	-	-	Tokyo Metropolitan Industrial Technology Research Institute	Dwarf; leaf spot; leaf shape	1983	
5	Big Cross	Iron Cross	1971	Gamma ray 3kR	Tokyo Metropolitan Isotope General Research Institute	1976	-	-	Tokyo Metropolitan Industrial Technology Research Institute	Leaf color; variation in leaf spot; larger iron-cruciate spot	1980	
6	Kaede Ion	Iron Cross	1972	Gamma ray 10kR	Tokyo Metropolitan Isotope General Research Institute	1976	-	-	Tokyo Metropolitan Industrial Technology Research Institute	Leaf shape; variation in leaf spot; reddish brown maple leaf shaped spot	1982	
<b>Resal pelargonium</b> <i>Pelargonium</i> L'Her. ex Ait.												
1	Capri Ice	Strawberry Thunder	1976	Petal culture	Ehime University	1985	1988	1096	Daiichi Seed Co., Ltd.		3222	
2	Capri Lullaby	Strawberry Thunder	1976	Anther culture	Ehime University	1985	1986	1098	Daiichi Seed Co., Ltd.		3223	
<b>Sonerila picta</b> <i>Sonerila</i> Roxb.												
1	Splash	<i>Sonerila picta</i> korth	1995	Chemical	Kansai TEC Co.	1997	2003	11451	Kansai TEC Co.	More spot on leaves	3178	
<b>Glossy abelia</b> <i>Abelia</i> R. Br.												
1	Meihan Hane-1	Hane-1 tsukubane-utsugi	1972	Gamma ray 3kR	Tokyo Metropolitan Isotope General Research Institute	1976	-	-	Tokyo Metropolitan Industrial Technology Research Institute	Leaf spot; slightly dwarf	1978	
<b>Japanese azalea</b> <i>Rhododendron indicum</i> (L.) Sweet												
1	Cobalt	Takasago	1964	Gamma ray 5kR	National Institute of Agricultural Sciences (Hiratsuka)	1972	-	-	Tochigi PAES	More branching; dwarf (useful for bonsai)	1986	
<b>Rose</b> <i>Rosa</i> L.												
1	Bridal Sonia	Sonia	1979	Gamma ray to plant 15kR	IRB	1982	1985	801	Kanagawa Prefectural Horticulture Experiment Station	Flower color: from salmon to pale pink	1994	
2	San Jol	Santa Monica	1985	Gamma ray	IRB	1987	1990	2264	Mr. Yuitsu MIZUTANI	Flower color: vivid purple	2543	pink

No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
3	Ohmiya-bito	Queen Elizabeth	1964	Gamma ray	IRB	1976	1990	2384	Keisei Rose Nurseries, Inc.	Flower color: from pink to pinkish white	1996	
4	Miss Ohmiya	Queen Elizabeth	1964	Gamma ray	IRB	1976	1990	2385	Keisei Rose Nurseries, Inc.	Flower color: from pink to vivid pink	1995	
5	Banbina	Green Eyes	1987	Chemical	KYOWA	1989	1995	4274	KYOWA	Flower color: pale pink	2544	
6	Ichi-anda	Asami Red	1983	Chemical	Mishima-shi	1996	2000	8349	Mr. Keichi ICHIKAWA	Small leaf; thread type	2545	
7	Ichi-runa	Aarus Mail Goal	1994	Chemical	Mishima-shi	1996	2000	8350	Mr. Keichi ICHIKAWA	Flower color and shape	2546	
8	Ichi-jiru	Ichity	1994	Gamma ray	Tokyo Metropolitan Isotope General Research Institute	1997	2001	8704	Mr. Keichi ICHIKAWA	Flower color and shape	2547	
9	Hitachi Smile	Samantha	1990	Gamma ray	IRB	2002	2007	15714	IRB and Okinawa Prefecture	Flower color and shape		
10	Hitachi Poeny	Samantha	1990	Gamma ray	IRB	2002	2007	15715	IRB and Okinawa Prefecture	Flower color and shape		
<b>Oncidium</b>												
<i>Oncidium</i> Group												
1	Gold One	Gower Ramsey	1986	Mericlone culture	Mr. Masayoshi SHIMOKAWA	1992	1995	4463	Mr. Masayoshi SHIMOKAWA	Flower color		
2	Sherley Baby White Flush Rika	Sherley Baby Sweet Fragrance	1988	Mericlone culture	Mr. Saneharu MATSUMOTO	1990	1997	5601	Mr. Saneharu MATSUMOTO	Flower color		
<b>Salvia</b>												
<i>Salvia splendens</i> Ker Gawl.												
1	Magunasu Cherry	Cherry Sage	2001	Colchicine	Toyota Motor Cooperation	2003	2006	14059	Toyota Motor Cooperation	Larger petal		
2	TL585	Lady In red		Ion beam irradiation to seed	JAEA, Takasaki	2012	2015	24473	Takii & Co, Ltd.	Shorter stem		
<b>Delphinium (larkspur)</b>												
<i>Delphinium</i> L.												
1	Star Dust	Bluecloud	1996	Colchicine	Toyama City Agriculture Center	1996	2001	8746	Toyama City Agriculture Center	Larger petal; leaf shape		
2	Sunny Sky Blue	Plage Sky Blue	1988	Tissue culture	Kaneko Seeds	2000	2005	12735	Kaneko Seeds	Shorter plant; shorter inflorescence		
<b>Torenia (wishbone flower)</b>												
<i>Torenia</i> L.												
1	San-reniramu	San-renimu	1998	Colchicine chromosome doubling	Suntory Flowers Ltd.	1999	2004	12092	Suntory Flowers Ltd.	Plant type: stoloniferous; flower color	3187	
2	San-renirahopasu	San-renihopasu	1998	Colchicine chromosome doubling	Suntory Flowers Ltd.	1999	2004	12093	Suntory Flowers Ltd.	Flower color of labium: uniformly colored	3188	

No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
3	San-reniraba	San-renireba	1999	Colchicine chromosome doubling	Suntory Flowers Ltd.	2000	2005	12993	Suntory Flowers Ltd.	Flower color of labium: uniformly colored	3189	
4	Sunrenripin	A developed line		Ion beam	Riken (Wako)	2007	2010	19301	Suntory Flowers Ltd.	Petal color: 72B		
5	Sunrenripihokai	Sunrenipihō		Colchicine	Suntory Flowers Ltd.	2003	2013	22609	Suntory Flowers Ltd.	Earlier flowering		
6	Sunrekodou	A developed line		Heavy ion beam	Riken (Wako)	2011	2013	22611	Suntory Flowers Ltd.	Petal color: 8B		
<b>Margaret</b> <i>Argyranthemum frutescens</i> (L.) Sch. Bip.												
1	Star Light Ripple	Zairai-shiro	1998	Soft X-ray to immature ovary culture	Shizuoka PAES	2002	2005	12779	Shizuoka PAES	Shorter plant type; larger stock		
2	Angel Maisu	not written	2002	Soft X ray to cuttings	Shizuoka PAES	2003	2006	14146	Shizuoka PAES	Flower color		
3	Southern Venus White	An original line	2002	Soft X ray to plants	Shizuoka PAES	2005	2007	14971	Shizuoka PAES			
4	Canary Queen	Peach Queen	2003	Soft X ray to plants	Shizuoka PAES	2005	2007	14992	Shizuoka PAES	Ligulate flower color change		<a href="http://www.geocities.jp/yunakisaragi/x-ma-garetto-kamariakuuin.html">http://www.geocities.jp/yunakisaragi/x-ma-garetto-kamariakuuin.html</a> (in Japanese)
<b>Lavender</b> <i>Lavandula angustifolia</i> Mill.												
1	Cosmic Blue	Yuulong	1998	Tissue culture of leaf	Kyoto Prefectural University	2001	2006	14044	SAKATA SEED CORPORATION	Plant type: spreading, color of calyx: deep purple; more flowers in the inflorescence		
<b>Cytisus (Scotch broom)</b> <i>Cytisus scoparius</i> (L.) Link												
1	Mei Shower	Crimson King	1990	Gamma ray to tissue culture	IRB	1993	1998	7027	IRB and Meiji Seika Kaisha, Ltd.	Extremely short plant; flower color: vivid yellow	3190	
2	Mei Wako	Crimson King	1990	Gamma ray to embryo culture	IRB	1993	1998	7028	IRB and Meiji Seika Kaisha, Ltd.	Extremely short plant; flower color: ground color: pale yellow; brown, deep purple pink	3191	
3	Mei Rose	Crimson King	1990	Gamma ray to embryo culture	IRB	1993	1998	7029	IRB and Meiji Seika Kaisha, Ltd.	Medium plant height; flower color: ground color: pale greenish yellow, brown purple pink	3192	
4	Mei Eve	Crimson King	1990	Gamma ray to embryo culture	IRB	1994	1998	7030	IRB and Meiji Seika Kaisha, Ltd.	Extremely short plant; flower color: ground color: pinkish white, brown reddish purple	3193	

No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
5	Mei Hiro	Crimson King	1990	Gamma ray to embryo culture	IRB	1993	2002	10948	IRB and Meiji Seika Kaisha, Ltd.	Outer standard ground color: rainbow; outer wing color: deep reddish brown	3194	
6	Mei Samson	Park Woody	1990	Gamma ray	IRB	1993	2002	10949	IRB and Meiji Seika Kaisha, Ltd.	Outer wing color: several colors; small wing	3195	
7	Mei Fanny	Crimson King	1990	Gamma ray to embryo culture	IRB	1997	2002	10950	IRB and Meiji Seika Kaisha, Ltd.	Outer standard and wing ground color: deep red	3196	
8	Mei Lord	Crimson King	1990	Gamma ray to embryo culture	IRB	1997	2002	10951	IRB and Meiji Seika Kaisha, Ltd.	Outer keel color: brown; wide keel	3197	
<b>Abelmoschus</b> <i>Abelmoschus manihot</i> Medik												
1	Hiroshima Local No. 1	Hiroshima Zairai	1956	Gamma ray 0.5kR	Kyoto University	1967	—	—	Kagawa University	Higher resistance to phytophthora rot	341	
2	Hiroshima Local No. 3	Hiroshima Zairai	1956	Gamma ray 0.5kR	Kyoto University	1967	—	—	Kagawa University	Higher resistance to phytophthora rot	342	
3	Hiroshima Local No. 5	Hiroshima Zairai	1956	Gamma ray 0.5kR	Kyoto University	1967	—	—	Kagawa University	Higher resistance to phytophthora rot	343	
4	Hiroshima Local No. 8	Hiroshima Zairai	1956	Gamma ray 0.5kR	Kyoto University	1967	—	—	Kagawa University	Higher resistance to phytophthora rot	344	
<b>Hibiscus (common rose mallow)</b> <i>Hibiscus moscheutos</i> L.												
1	Shirasagi-no-yume	Sakai-no-hana	1981	Gamma ray 30kR	Osaka Prefectural Radiation Research Center (now Osaka Prefectural University)	1987	1987	1311	Osaka Prefectural Radiation Research Center (now Osaka Prefectural University)	Flower color: pinkish white-vivid purple pink with purple red stripe	1979	
<b>Gentiana</b> <i>Gentiana</i> L.												
1	Miyama Love	Sasa-rindou		Colchicine treatment	Sky Blue Seto Co.	2001	2005	13126	Sky Blue Seto Co.	Late maturing		
2	Shimano Love	Ezo-rindou		Colchicine treatment and hybridization between chromosome-doubled plants	Sky Blue Seto Co.	2001	2005	13128	Sky Blue Seto Co.	Chromosome doubling (tetraploid)		
<b>Verbena</b> <i>Verbena</i> L.												
1	San-mariko-rabi	San-mariribi	1998	Ion beam	Riken (Wako)	2001	2005	12999	Suntory Flowers Ltd.	Flower color: vivid purple; eye-shaped spot on the petal		
2	San-mari-sakura	San-marrisa		Radiation	No description	2003	2011	20462	Suntory Flowers Ltd.	Plant type		



No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
3	San-mari-peach	A developed line		Heavy nitrogen ion beam irradiation to flower buds	(Wako)	2006	2008	16854	Suntory Flowers Ltd.	Flower color and shape		
<b>Osteospermum</b>												
<i>Osteospermum</i> L.												
1	Viento Flamingo	Mother Symphony		Ion beam	JAEA, Takasaki	2007	2009	18549	JAEA and Hanasekiguchi Co.	Petal color: 29D		Iizuka <i>et al.</i> (2011) <sup>(2)</sup>
2	Viento Labios Flamingo			Radiation	JAEA, Takasaki	2011	2016	24758	JAEA and Hanasekiguchi Co.			
<b>Bidens</b>												
<i>Bidens</i> L.												
1	San-bideki	Bidens triplinervia	1997	Colchicine	Suntory Flowers Ltd.	2001	2004	12262	Suntory Flowers Ltd.	Ever-flowering		
<b>Gardenia</b>												
<i>Gardenia jasminoides</i> J. Ellis												
1	Vald	A domestic line		Colchicine treatment to seeds	Toyota Motor Corporation		2005	13621	Toyota Motor Corporation	Leaf shape		
<b>Asian melastoma</b>												
<i>Melastoma</i> L.												
1	Pink Pearl	Belgra	1996	Gamma ray irradiation to callus	GIFUSEED Co. Ltd.		2005	13519	GIFUSEED Co. Ltd.	Pattern of leaf surface		
<b>Morning glory</b>												
<i>Ipomoea indica</i> (Burm.) Merr.												
1	IRBii Light Blue	Capetown Blue		Chronic gamma ray irradiation in the Gamma Field	IRB, NIAS	2012	2015	24599	IRB, NIAS, Fukukaen Nursery & Bulb Co., Ltd.	Flower color: 98B		
<b>Italian ryegrass</b>												
<i>Lolium multiflorum</i> Lam. ssp. <i>italicum</i> (A. Br.) Volkart												
1	Miyuki-aoba	Taka-kei No. 4		Maternal selection after colchicine treatment	Hokuriku NAES	1983	1984	649	Hokuriku NAES	Larger leaf	2548	
<b>Japanese lawngress</b>												
<i>Zoysia</i> Willd.												
1	Winter Carpet	Tsukuba-kei	1990	Gamma ray	IRB	1992	1995	4299	IRB and Sumitomo Metal Industries, Ltd.	Earlier green color of leaf	3201	
2	Winter Field	Tsukuba-kei (Z. <i>Matrella</i> ; <i>manilagrass</i> )	1990	Chronic gamma ray irradiation (Gamma field)	IRB	1994	1996	5254	IRB and Sumitomo Metal Industries, Ltd.	Better stolon vigor; narrower and shorter leaf	3202	

No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
3	Kopurosu	Tottori-kourai	1995	X ray to meristem culture	Yamaguchi PAES	1998	2002	10638	Kaisui Chemical Industry Co.	Better spring vigor; earlier green color of leaf; no browning in winter	3203	
4	Tee Emu 9	<i>Z. murella</i> ecotype	2000	Soft X-ray irradiation to the seeds	Toyota Motor Corporation	2003	2006	14320	Toyota Motor Corporation	More flower heads; earlier browning in autumn		
5	Shiba Noh Pl 1	Ecotype		Tissue culture of seeds and regeneration from callus	NARO, National Institute of Livestock and Grassland Science	2007	2009	18204	NARO	Thicker turf development		
<b>Creeping bentgrass <i>Agrostis stolonifera</i> L.</b>												
1	Spring	Pennecross	1970	Gamma ray	Kyushu University	1982	1983	469	Nihon Ryokuei Co.	Resistance to summer heat	293	
2	Chiba Green B-2	Pennecross	1991	Soft X ray to seedling	Chiba PAES	1996	1997	5439	Chiba PAES	Resistant to brown patch disease	2549	
3	My Comfort	Pennecross	1992	Tissue culture	Chiba PAES	1996	2001	9043	Maekawa MFG Co.	Taller plant	2550	
<b>Guineagrass <i>Panicum maximum</i> Jacq.</b>												
1	PL 1 (Nekken No. 1)	guineagrass Noh	GR297	Colchicine	Coastal Plain Experiment Station, Tifton, USDA, GA, USA	1990	1993	3708	Tropical Agriculture Research Center (Now JIRCAS)	Chromosome doubling; shorter plant		Nakagawa <i>et al.</i> (1992) <sup>32)</sup>
<b>Red clover <i>Trifolium pratense</i> L.</b>												
1	Taisetsu	Sapporo		Colchicine	Hokkaido NAES	1991	1991	2934	Hokkaido NAES	Chromosome doubling; larger leaf	3205	
<b>Apple <i>Malus pumila</i> Mills.</b>												
1	Mori-hou-fu 3A	Fuji	1963	Gamma ray 3kR to scion	IRB	1963			IRB	Resistance to disease	2551	
2	Houiku Indo	Indo	1992	Gamma ray irradiation to meristem culture	IRB	2003	2007	15022	IRB	Resistant to Alternaria disease		
<b>Japanese pear <i>Pyrus pyrifolia</i> (Burm. f.) Nakai var. <i>culta</i> (Mak.) Nakai</b>												
1	Gold Nijisseiki	Nijisseiki	1962~8	Chronic irradiation (53m from center of source in Gamma Field)	IRB	1990	1991	2932	IRB	Resistance to black spot disease	278	Sanada <i>et al.</i> (1993) <sup>45)</sup>
2	Kotobuki Sinsui	Sinsui	1989	Gamma ray 80Gy to scion	IRB	1996	1997	5436	IRB and Tottori Prefectural Horticulture Experiment Station	Resistance to black spot disease	279	Kitagawa, K. <i>et al.</i> (1999) <sup>25)</sup>

No.	Cultivar Name	Original Cv. or Line	Treat. Year	Treatment	Treated Institution	Dev. year	Reg. Year	Reg. No. <sup>a</sup>	Institution of Development	Improved Characteristics	IAEA ID <sup>b</sup>	Reference
3	Osa Gold	Osa-nijisseiki 9	1986~9	Chronic irradiation (40m from center of source in Gamma Field)	IRB	1995	1997	5620	IRB and Tottori Prefectural Horticulture Experiment Station	Resistance to black spot disease	2552	Masuda <i>et al.</i> (1998) <sup>28)</sup>
<b>Peach</b>												
1	Fuku-ekubo	Akatsuki	1987	Gamma ray 3kR	IRB	1994	1996	5076	Fukushima Prefectural Fruit Tree Experiment Station	Early maturing	2553	Sakuma <i>et al.</i> (1999) <sup>44)</sup>
2	RS	Shimizu Hakutou Hakutou	1996	Gamma ray irradiation to nursery stock	IRB	2001	2004	12302	Okayama PAES	Resistant to black spot disease	2554	Inoue <i>et al.</i> (2006) <sup>15)</sup>
<b>Loquat</b>												
1	Shiro-Mogi	Mogi	1981	Gamma ray	IRB	1981	1982	300	Nagasaki Prefectural Fruit Tree Experiment Station		240	
<b>Cherry</b>												
1	Super 6	Koruto	1985	Colchicine treatment to meristem culture	Tenkoen Co.	1988	1997	5374	Tenkoen Co.	Thicker branch; flower petal: round; larger leaf		
2	Roman Nishiki	Masamitsu-nishiki	1987	Colchicine treatment to buds	Mr. Masamitsu SATO	1998	2005	9929	Mr. Masamitsu SATO	Larger leaf; stickiness between seed and skin: medium sticky		
<b>Cherry flower</b>												
1	Nishina-Zaou	Groiko		Carbon ion beam	Riken (Wako)	2007	2009	17785	Riken and Shigehisa ISHII	Leaf shape and flower structure	3355	
2	Nishina-otome	Keiou Zakura		Heavy ion beam	Riken (Wako)	2009	2011	21281	Riken	Perpetual flowering		
<b>Mulberry</b>												
1	Lala Berry	Kataneo	1994	Colchicine treatment to axillary buds	NIAS	2000	2003	11242	NIAS	Thicker and larger leaf		
2	Pop Berry	Daitou-gawa	1994	Colchicine treatment to axillary buds	NIAS	2001	2004	12194	NIAS	Pericinal chimera; heavier fruit		
<b>Tea</b>												
1	Tea Noh Pl 2	Yabukita	1970-1971	Chronic irradiation in gamma field	IRB	1988	1998	6449	IRB	From self-incompatible to self-compatible; deeper leaf color		



**Note:**

\*Reg. No. is the registration number of The Plant Variety Protection System in Japan  
 †IAEA ID is the variety ID number of the Joint FAO/IAEA Mutant Variety Database (<https://mvd.iaea.org/#!Home>)

**Abbreviations:**

ARC: Agricultural Research center  
 ARI: Agricultural Research Institute  
 IRB: Institute of Radiation Breeding  
 JAEA: Japan Atomic Energy Agency  
 JIRCAS: Japan International Research Center for Agricultural Sciences  
 NAES: National Agricultural Experiment Station  
 NARC: National Agriculture Research Center  
 NARO: National Agriculture and Food Research Organization  
 NIAES: National Institute of Agro-environmental Sciences  
 NIAS: National Institute of Agrobiological Sciences  
 PAES: Prefectural Agriculture Experiment Station  
 PARC: Prefectural Agriculture Research Center

**Literature cited**

- 1) Arai, M. (2001) The characteristics of a new carnation variety "Your Red" bred by selection of somaclonal variation. *Tohoku Agric. Research* 54: 235-236 (in Japanese).
- 2) Ezura, H., H. Amagai, M. Kasumi, and Y. Ishizuka (1998) A new strawberry variety "Anteher" derived from somaclonal variants of callus culture. *Bull. Ibaraki Plant Biotech. Inst.* 2: 75-82 (in Japanese with English summary).
- 3) Fukuoka, H., T. Murata, A. Komura, S. Nakao, and H. Kinouchi (1996) A strawberry cultivar "Himatsuri". *Bull. Kyushu Tokai University Faculty of Agriculture* 15: 19-25 (in Japanese with English summary).
- 4) Furuichi, T., K. Murakami, and Y. Matsumoto (2003) Breeding of "Princess Kagawa" - a new cultivar of *Dendranthema X grandiflorum*. *Bull. Kagawa Pref. Agric. Exp. Stn.* 56: 57-60 (in Japanese with English summary).
- 5) Futuhara, Y. (1968) Breeding of a new rice variety Reimei by gamma-ray irradiation. *Gamma Field Symposia* 7: 87-109.
- 6) Hajjika, M., M. Takahashi, K. Igita, S. Sakai, and Y. Nakazawa (2002) A new soybean variety "Ichihime". *Bull. Kyushu Okinawa Agric. Res. Cent.* 40: 79-94 (in Japanese with English summary).
- 7) Hashimoto, K., T. Nagasawa, S. Murakami, I. Watanabe, K. Kokubun, S. Sakai, K. Igita, and A. Okabe (1988) A new soybean variety "Kosuzu". *Bull. Tohoku Natl. Agric. Exp. Stn.* 77: 45-61 (in Japanese with English summary).
- 8) Hashimoto, K., T. Nagasawa, S. Murakami, K. Kokubun, S. Nakamura, T. Koyama, S. Matsumoto, K. Sasaki, and A. Okabe (1985) A new soybean variety "Wasesuzunari". *Bull. Tohoku Natl. Agric. Exp. Stn.* 71: 23-42 (in Japanese with English summary).
- 9) Hikawa, M., T. Nobumori, Y. Naito, S. Okatake, T. Kawai, K. Ishida, H. Kimoto, and S. Fujii (1998) A new butterbur variety "Okayamanoshi B1 gou". *Bull. Okayama Pref. Agric. Exp. Stn.* 16: 21-23 (in Japanese).
- 10) Hoshino, T., T. Nakamura, Y. Seimiya, T. Kamada, G. Ishikawa, A. Ogasawara, S. Sagawa, M. Saito, H. Shimizu, M. Nishi, M. Watanabe, J. Takeda, and Y. Takahata (2010) Production of a fully waxy line and analysis of waxy genes in the allohexaploid crop, Japanese barnyard millet. *Plant Breed.* 129: 349-355.
- 11) Iida, S., E. Amano, and T. Nishio (1993) A rice (*Oryza sativa* L.) mutant having a low content of glutenin and a high content of prolamine. *Theor. Appl. Genet.* 87: 374-378.
- 12) Iizuka, M., Y. Kimura, T. Okada, Y. Hase, A. Tanaka, and M. Sekiguchi (2011) A new cultivar "Viento Flamingo" of *Osteospermum*. *Bull. Gumma Agric. Res. Cent.* 8: 89-92 (in Japanese with English summary).
- 13) Imakire, S., S. Nagayoshi, K. Koriyama, and K. Ueno (2006) Breeding process and characteristics of standard type chrysanthemum cultivar "Araddin". *Bull. Kagoshima Pref. Agric. Exp. Stn.* 34: 15-19 (in Japanese).  
 Japanese with English summary.
- 14) Imbe, T., N. Horisue, Y. Akama, A. Nakane, I. Ando, M. Suto, T. Hata, K. Ise, K. Numaguchi, H. Furutachi, and M. Takadate (2004) "Oodorokimochi" - A multipurpose high-yield glutinous rice variety. *Bull. Natl. Inst. Crop Sci.* 6: 19-35 (in Japanese with English summary).
- 15) Inoue, K., Y. Fujii, Y. Sasabe, M. Otsuka, H. Kagami, H. Nasu, T. Kimura, S. Yoda, and S. Kasuyama (2006) New peach cultivar "Shimizu hakuto RS" which has moderately resistant to black spot. *Bull. Okayama Pref. Agric. Exp. Stn.* 24: 11-15 (in Japanese with English summary).
- 16) Ise, K., Y. Akama, N. Horisue, A. Nakane, M. Yokoo, I. Ando, T. Hata, M. Suto, K. Numaguchi, H. Nemoto, H. Furutachi, and T. Imbe (2001) "Milky Queen", a new high-quality rice cultivar with low amylose content in endosperm. *Bull. Natl. Inst. Crop Sci.* 2: 39-61 (in Japanese with English summary).

- 17) Ishida, M., M. Kato, and I. Chiba (1995) Breeding process and characteristics of a new Job's tear cultivar "Hatohikari". Tohoku Agric. Res. 48: 141-142 (in Japanese).
- 18) Ishikawa, H., T. Satoh, T. Uwano, M. Nitta, Y. Kiuchi, and T. Sasaki (1988) The new rice variety "Iwate-21" - developed by radiation breeding - Bull. Iwate Pref. Agric. Exp. Stn. 27: 25-39.
- 19) Ito, R. (1991) On the new rice variety "Hatsuyume" and "Yumekaori" induced by protoplast culture and their breeding back ground. Journal of agricultural science 46: 350-354 (in Japanese).
- 20) Kaji, R., M. Okamoto, M. Sakai, R. Mizobuchi, K. Tamura, H. Hirabayashi, Y. Tamura, S. Fukaura, T. Tomimatsu, and H. Nakano (2010) "Hatsuyamabuki", a new rice variety with yellow endosperm derived from gamma ray mutation. Breeding Research 12: 54-58 (in Japanese).
- 21) Kanetou, A., S. Torii, T. Akiyama, D. Miyoshi, and H. Asaumi (2010) Breeding of new rice cultivar for sake brewery "SIZUKUHIME". Bull. Ehime Pref. Agric. Exp. Stn. 2: 1-6 (in Japanese with English summary).
- 22) Kato, H., H. Yoshioka, R. Iba, H. Mitsuueda, T. Nakahara, H. Takeda, S. Tsutsumi, and K. Wakasugi (2006) A newly bred rice variety "Minamiyutaka" for whole-crop silage. Bull. Miyazaki Agric. Exp. Stn. 41: 51-60 (in Japanese with English summary).
- 23) Katsuta, M., M. Komeichi, Y. Okuyama, Y. Honda, S. Shirato, and M. Nakatani (2001) A new grain amaranthus variety "New Aztec". Bull. Nat. Inst. Crop Sci. 1: 57-70 (in Japanese with English summary).
- 24) Kiribuchi-Otobe, T. Yanagisawa, H. Yoshida, I. Yamaguchi, H. Seko, T. Ushiyama, and Y. Amano (2001) Breeding of "Wheat Norin-PL 8 (Tanikei A6599-4)", a waxy wheat line with starch showing a stable hot paste viscosity. Bull. Natl. Inst. Crop Sci. 2: 63-74 (in Japanese with English summary).
- 25) Kitagawa, K. *et al.* (1999) A new Japanese pear cultivar, "Kotobuki Sinsui". Bull. Tohori Hort. Exp. Stn. 3: 1-13 (in Japanese).
- 26) Laboratory of Soybean Breeding (1970) The new soybean varieties "RAIDEN" and "RAIKO" induced by gamma-ray irradiation. Bull. Tohoku Natl. Agric. Exp. Stn. 40: 65-105 (in Japanese with English summary).
- 27) Maeshige, M., T. Tsuchiya, Y. Doi, S. Otake, S. Umemoto, Z. Katsuba, and Y. Sakai (1995) A new paddy rice variety "Meguriai". Bull. Hiroshima Pref. Agric. Res. Ctr. 62: 31-38 (in Japanese with English summary).
- 28) Masuda, T., T. Yoshioka, T. Sanada, K. Kotobuki, M. Nagaraj, M. Uchida, K. Inoue, K. Murata, K. Kitagawa, and A. Yoshida (1998) A new Japanese pear cultivar "Osa Gold", resistant mutant to the black spot disease of Japanese pear (*Pyrus pyrifolia* Nakai) induced by chronic irradiation of gamma-rays. Bull. Natl. Inst. Agrobiol. Resour. 12: 1-11 (in Japanese with English summary).
- 29) Minami, M. and H. Sato (2009) A new scarlet runner bean variety "Shirohanakko". Bull. Hokkaido Central Agric. Exp. Stn. 93: 25-34 (in Japanese with English summary).
- 30) Miyamoto, Y., T. Kobata, and K. Kato (2003) Studies on breeding of *Gypsophila paniculata* L. (2) Breeding process and characteristics of a new cultivar "Kisyu-kasumi 1". Bull. Wakayama Res. Cent. Agric. Forest. Fisheries 4: 33-41 (in Japanese with English summary).
- 31) Miyashita, C., H. Sakaemori, H. Minami, and R. Uzawa (2014) A new registered weak-roseette-type parental variety of eustoma. Technical Report of Tokyo Metropolitan Agric. Forest. Res. Cent. 9: 15-22 (in Japanese).
- 32) Nakagawa, H. and W. W. Hanna (1992) Induced sexual tetraploids for breeding guineagrass (*Panicum maximum* Jacq.). J. Japan. Grassl. Sci. 38: 152-159.
- 33) Nakagawa, H., N. Shimizu, H. Sato, T. Momonoki, and H. Nakano (1991) Development of a sexually reproducing tetraploid line for breeding apomictic guineagrass. TARC Newsletter Vol. 2, No. 3: 4-5.
- 34) Nakagawa, T., H. Asaumi, and M. Tamaki (2015) "Ehime-Noushi V2 gou", a new taro cultivar in Ehime Prefecture. Bull. Ehime Pref. Agric. Exp. Stn. 7: 16-20 (in Japanese).
- 35) Nakahara, T., T. Sumiyoshi, N. Morifuji, K. Hirashima, M. Koga, H. Matsui, M. Okuma, S. Fujitomi, and Y. Uchimura (1998) A new rush cultivar "CHIKUGOMIDORI" derived from somatic embryo. Bull. Fukuoka Agric. Res. Cent. 17: 57-61 (in Japanese with English abstract).
- 36) Nakajo, S., S. Hasegawa, H. Yoshida, S. Urushibara, A. Abe, T. Abe, N. Fukunishi, H. Ryoutou, and Y. Ohshimizu (2013) Breeding new varieties of Japanese barnyard millet with shorter culms and low amylose content: "Nebarikko 1", "Nebarikko 2" and "Nebarikko 3". Res. Bull. Iwate. Agric. Res. Ctr. 12: 43-60 (in Japanese with English summary).
- 37) Nakamura, D., H. Yokoh, Y. Hirota, K. Nonaka, T. Shigyo, H. Oh-ishi, O. Shigetomi, Y. Takagi, and H. Kishikawa (1991) A new soybean cultivar "Murrayutaka". Bull. Saga Agric. Exp. Stn. 27: 21-42 (in Japanese with English summary).
- 38) Nishi, A., Y. Nakamura, N. Tanaka, and H. Sato (2001) Biochemical and genetic analysis of the effects of amylose-extender mutation in rice endosperm. Plant Physiol. 127: 459-472.
- 39) Nonaka, S., Y. Yoshida, S. Kitahara, M. Tsuru, S. Taya, H. Araki, and H. Gocho (1979) A new wheat cultivar "SHIROWASE-KOMUGI". Bull. Kyushu Natl. Agric. Exp. Stn. 20: 209-220 (in Japanese with English summary).
- 40) Okamura, M., M. Momose, S. Watanabe, A. Tanaka, and Y. Hase (2006) Development of stress tolerant variety series in ornamentals by ion beam breeding. Proc. The 3rd Ion Beam Breeding Workshop (<http://jolissreh-inter.tokai-sc.jaea.go.jp/search/servelet/search?5002578>).
- 41) Okuyama, Y., S. Sugawara, K. Shindo, K. Seki, N. Ishikura, S. Tanosaki, T. Endo, M. Shibata, and M. Ishida (1995) A new job's tear variety "Hatomusume". Bull. Tohoku Natl. Agric. Exp. Stn. 89: 1-10 (in Japanese with English Summary).
- 42) Sadahira, M., S. Hamada, T. Akagi, Y. Shimoyamane, Y. Nakano, T. Yoshizaki, H. Kurata, and T. Ushiro (1982) Raising of new mat rush variety "Setonami" by radiation treatment. Bull. Hiroshima Pref. Agric. Exp. Stn. 45: 105-112 (in Japanese with English summary).
- 43) Sadahira, M., S. Hamada, Y. Shimoyamane, T. Akagi, H. Kurata, T. Yoshizaki, and T. Ushiro (1984) Breeding of "Fukunami", a new variety of mat rush, by gamma-ray irradiation. Bull. Hiroshima Pref. Agric. Exp. Stn. 48: 105-112 (in Japanese with English summary).
- 44) Sakai, M., H. Nakano, M. Okamoto, K. Tamura, R. Kaji, Y. Tamura, T. Kataoka, and R. Mizobuchi (2013) "Ruriaoba", a new rice forage variety for double harvesting. Bull. Kyushu Okinawa Agric. Res. Cent. 60: 1-12 (in Japanese with English Summary).

- 45) Sakuma, N., T. Munakata, H. Yamaga, T. Kunisawa, and H. Matsuno (1999) New peach cultivar "Fukuekubo". Bull. Fukushima Pref. Agric. Exp. Stn. 17: 1-5 (in Japanese with English summary).
- 46) Sanada, T., K. Kotobuki, M. Nishida, H. Fujita, and T. Ikeda (1993) A new Japanese pear cultivar "Gold Nijisseiki", resistant to black spot disease of Japanese pear. Jpn. J. Breed. 23: 455-461 (in Japanese with English summary).
- 47) Sasaki, K., Y. Sato, M. Suzuki, K. Inoue, H. Igarashi, T. Kutsuzawa, A. Kikawa, Y. Mizokoshi, J. Fujimoto, and K. Okada (2000) Breeding of a new green seed-coated soybean cultivar "Akitamidori" and its characteristics. Bull. Akita Agric. Exp. Stn. 41: 1-16 (in Japanese with English summary).
- 48) Sato, T., Y. Himori, S. Tsubaki, and H. Sano (2015) Breeding of new green soybean varieties "Akinoushi-40" and "Akita-Honoka". Bulletin of Akita Pref. Exp. Stn. 55: 40-59 (in Japanese with English Summary).
- 49) Satoh, H., Y. Takahata, N. Kamata, K. Sasaki, F. Furusawa, Y. Kamiyama, Y. Ohno, K. Sasaki, N. Takahashi, and S. Maita (1982) A new azuki bean variety, Beni-nanbu. Gamma Field Symposia 21: 31-36.
- 50) Shizuoka Prefecture (2010) Characteristics of a new rice recommended cultivar "Homare-fuji", Atarashii Nougyou Gijyutsu (New Agricultural Technologies). Shizuoka Pref. Res. Inst. of Agric. and Forest.: 8 pp. (in Japanese).
- 51) Suzuki, K., M. Ohara, M. Ino, T. Takeuchi, M. Ito, H. Sakiyama, and S. Ushio (2006) Breeding of a new eddo (*Colocasia esculenta* L.) cultivar, "Chiba maru". Bull. Chiba Pref. Agric. Res. Cent. 5: 65-74 (in Japanese with English summary).
- 52) Suzuki, T., M. Kimura, M. Kawakatsu, K. Nakatsuka, S. Kim, Y. Mukasa, S. Yokota, H. Yamauchi, S. Takigawa, T. Noda, N. Hashimoto, and C. Matsuura-Endo (2008) "Hokkai T9" and "Hokkai T10": new Tertiary buckwheat varieties for sprouts and dried powders. Bull. Natl. Agric. Res. Cent. Hokkaido Reg. 188: 45-53 (in Japanese with English summary).
- 53) Takahashi, K., S. Shimada, H. Shimada, Y. Takada, T. Sakai, Y. Kono, T. Adachi, K. Tabuchi, A. Kikuchi, S. Yumoto, S. Nakamura, M. Ito, H. Banba, and A. Okabe (2004) A new soybean cultivar "Yumeminori" with low allergenicity and high content of 11S globulin. Bull. Natl. Agric. Res. Cent. Tohoku Reg. 102: 23-39 (in Japanese with English abstract).
- 54) Ueno, K., S. Nagayoshi, S. Imakiire, K. Koriyama, T. Minami, A. Tanaka, Y. Hase, and T. Matsumoto (2013) Breeding of new chrysanthemum cultivar "Aladdin 2" through stepwise improvements of cv. "Jimba" using ion beam re-irradiation. Hort. Res. (Japan) 12 : 245-254 (in Japanese with English abstract).
- 55) Yamada, H., F. Baba, Y. Hagiwara, and H. Ohtsuka (1999) Breeding of flower color mutant cultivar "Ray sunrise" of spray-type chrysanthemum (*Dendranthema grandiflora*) by soft X-ray irradiation. Bull. Shizuoka Agric. Exp. Stn. 44: 37-41 (in Japanese with English summary).
- 56) Yamada H., H. Inagaki, Y. Hagiwara, and H. Otsuka (2002) Breeding of flower color mutant cultivar "Dreaming" of spray-type chrysanthemum (*Dendranthema grandiflora*) by soft X-ray irradiation. Bull. Shizuoka Agric. Exp. Stn. 47: 43-48 (in Japanese with English summary).
- 57) Yamaguchi, I., C. Kiribuchi-Otobe, T. Yanagisawa, T. Ushiyama, H. Seko, Y. Amano, S. Miyagawa, A. Kuroda, and S. Oda (1998) Breeding of "Wheat Norin-PL 7 (Tanikei A6099)", a new germplasm accession with low amylose content. Bull. Natl. Agric. Res. Cent. 29: 1-11 (in Japanese with English summary).

## Appendix 2. Indirect-use mutant cultivars

No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
<i>Oryza sativa</i> L.										
Yamada-nishiki										
1	Hyo-kei-sake 18		I-M106*	*Gamma ray irradiation to cv. Norin No. 8	Hyogo PAES	1972		Short and tough culm; good quality; very large seed	1125	
2	Mutsuhonami	Etsu-nan No. 39	Fu-kei No. 70 (Reimei)*	*cf. cv. Reimei (Appendix 1)	Aomori PAES, Fujisaka Branch	1973		Short and tough culm; high-yielding	1133	
3	Kagahikari	R6-1*	Fu-kei No. 72	*Gamma ray to cv. Koshihikari	Ishikawa PAES	1973		Early maturity	1126	
4	Hanahikari	stripe36/Bi-kei No. 53/Bi-kei No. 53	Fu-kei No. 70 (Reimei)*	*cf. cv. Reimei (Appendix 1)	Yamagata PAES	1975		Short and tough culm; early maturity	1122	Kamei <i>et al.</i> (1980) <sup>(58)</sup>
5	Houhai	Ao-kei No. 62 (Kojyou-nishiki)	Reimei*	*cf. cv. Reimei (Appendix 1)	Aomori PAES	1975		Short and tough culm; cold tolerance	1124	
6	Akihikari	Ouu No. 269 (Toyonishiki)	Reimei*	*cf. cv. Reimei (Appendix 1)	Aomori PAES, Fujisaka Branch	1976		Short and tough culm; very high-yielding	1120	Kushibuchi <i>et al.</i> (1977) <sup>(65)</sup>
7	Hayahikari	Fu-kei No. 70 (Reimei)*	Ouu No. 269 (Toyonishiki)	*cf. cv. Reimei (Appendix 1)	Tohoku NAES	1976		Semi-dwarf; lodging resistance	1123	Hirano <i>et al.</i> (1977) <sup>(26)</sup>
8	Nadahikari	5810-19/Hyo-kei Sake No. 18*	Tou-Kin-kei No. 1011	*cf. cv. "Hyo-kei-sake No. 18"	Hyogo PAES	1977		Short culm; big seed	1136	
9	Fujihikari	R151*	Fu-kei No. 71**/No. 67/Koshihikari B1F1	*Gamma ray to cv. Koshihikari; **Gamma ray to cv. Fujiminori	Chugoku NAES			Extremely early-maturing; short culm	1121	Fujii <i>et al.</i> (1981) <sup>(1)</sup>
10	Sachiminori	Manryou/R4B*	Yamase-nishiki	*Gamma ray to "Pi No. 4"	Hokuriku NAES	1978		Slightly tough stem; resistant to rice blast (Pi-ta')	1140	Samoto <i>et al.</i> (1979) <sup>(41)</sup>
11	Katsura-wase	Fu-kei No. 67*/Fu-kei No. 71*	Koshihikari	*Gamma ray to the seed of cv. Fujiminori	Kagoshima PAES	1978		Early maturing; short and tough culm	1127	
12	Miyamishiki	Kanto No. 79*	Todoroki-wase	*Gamma ray (20kR) to the seed of cv. Koshihikari	Miyazaki PAES	1978		Good quality; good taste	1130	Uchiyamada <i>et al.</i> (1979) <sup>(85)</sup>
13	Niigata-wase	Fu-kei No. 91*	Chou 60	*A descendant (the 1st generation) of cv. Reimei	Niigata PAES	1979		Short and tough culm	1137	Ichikawa <i>et al.</i> (1981) <sup>(88)</sup>
14	Mine-ssahi	Kanto No. 79*	Kihou	*Gamma ray (20kR) to the seed of cv. Koshihikari	Aichi PAES	1980	100	Good taste; short culm	1128	Morimoto <i>et al.</i> (1980) <sup>(86)</sup>
15	Musashi-kogane	Tama-kei No. 56*	Aichi No. 21	*cv. Reimei/cv. Nipponbare	Saitama PAES	1981	239	Short culm; lodging resistance	1132	Shiobara <i>et al.</i> (1982) <sup>(54)</sup>
16	Mutsu-komachi	Mutsu-nishiki	Fu-kei No. 104 (Akihikari)*	*cf. cv. Akihikari (the 1st generation of cv. Reimei)	Aomori PAES	1981	241	Short culm; lodging resistance	1135	Ono <i>et al.</i> (1982) <sup>(33)</sup>
17	Mutsu-kaori	Mutsu-nishiki	Fu-kei No. 104 (Akihikari)*	*cf. cv. Akihikari (the 1st generation of cv. Reimei)	Aomori PAES	1981	242	Short culm; lodging resistance	1134	Ono <i>et al.</i> (1982) <sup>(34)</sup>
18	Rokko-nishiki	5810-19/Hyo-kei Sake No. 18*	Tou-Kin-kei No. 1011	*cf. cv. "Hyo-kei sake No. 18"	Hyogo PAES	1981	326	Large seed	1139	
19	Miyakaori	Iwaga	Ouu No. 282*	*Named as cv. Hayahikari (the 1st generation of cv. Reimei)	Miyagi PAES Furukawa Branch	1983	588	Semi-dwarf	2986	Oikawa <i>et al.</i> (1991) <sup>(30)</sup>
20	Natsuhikari	Ka-kei No. 639*	Sei-nan No. 45	* Named as "Katsura-wase" (the 3rd generation of a gamma ray induced strain "Fu-kei 71")	Kagoshima PAES	1983	598	Tough stems; lodging resistance	2987	Iwashita <i>et al.</i> (1984) <sup>(49)</sup>
21	Megumi-mochi	Aichi-mochi No. 27	Fu-kei No. 102*	*cv. Reimei/cv. Fujiminori/cv. Somewake	Aichi ARC	1983	623	Short and tough culm; high-yielding	1145	Koumura <i>et al.</i> (1983) <sup>(83)</sup>



No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. Year	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
22	Hatsukogane	Fu-kei No. 102*	Matsumae	*cv. Reimei/cv. Fujiminori/cv. Somewake	Aomori PAES, Fujisaka Branch	1984	1985	889	Short and tough stem	1146	Shimura <i>et al.</i> (1986) <sup>(53)</sup>
23	Okuhomare	Hyo-kei-sake No. 18* Reimei**	Reimei**	*cf. cv. "Hyo-kei-sake No. 18"; **cf. cv. Reimei (Appendix I)	Hyogo PAES	1984			Semi-dwarf	2988	
24	Ibuki-wase	Mine-asahi*	Hourei	*cf. cv. "Mine-asahi" (the 1st generation of gamma ray induced mutant line "Kanto No. 79" from cv. Koshihikari)	Aichi ARC	1985	1986	1204	Cold tolerance	1148	Akama <i>et al.</i> (1985) <sup>(2)</sup>
25	Mutsu-homare	Todoroki-wase/Akibikari*	Fuji 329	*cf. cv. Akibikari (the 1st generation of cv. Reimei)	Aomori PAES	1985	1986	1206	Short culm; good taste	1149	Yamazaki <i>et al.</i> (1987) <sup>(210)</sup>
26	Hanafubuki	Ao-kei No. 79 (Oku-homare)*	Fu-kei No. 103**	*cf. cv. Okuhomare; **A descendant (the 1st generation) of cv. Reimei	Aomori PAES	1985	1988	1494	Semi-dwarf	2989	Tanabu <i>et al.</i> (1987) <sup>(72)</sup>
27	Tamaminori	Tama-kei No. 56*	Aichi No. 21	*cv. Reimei/cv. Nipponbare	Saitama PAES	1986	1986	1196	Tough culm; semi-dwarf	2990	Niwayama <i>et al.</i> (1987) <sup>(25)</sup>
28	Hyogo-Kitanishiki	Nadahikari*	Gohyaku-mangoku	*cf. cv. Nadahikari	Hyogo PAES	1986	1987	1374		2991	
29	Akchikara	Hokuriku No. 101*	Akibikari**	*A descendant of cv. Reimei; **A descendant (the 1st generation) of cv. Reimei	Hokuriku NAES	1986	1987	179	Short and tough culm; high-yielding	1147	Koga <i>et al.</i> (1987) <sup>(78)</sup>
30	Manyo-mochi	77-5133*	To-kou No. 76	*A glutinous mutant of cv. Reimei	Toyama PAES	1987	1988	1738	Glutinous	2992	Yamamoto <i>et al.</i> (1989) <sup>(208)</sup>
31	Seihou No. 1	Akibikari*	Nishihomare	*cf. cv. Akibikari (the 1st generation of cv. Reimei)	Mr. Seiichi HIROKI	1987	1988	1799		2993	
32	Aichi-no-kaori	Hatsushimo	Mine-asahi*	*A descendant of Gamma ray (20KR) induced mutant line "Kanto 79" from cv. Koshihikari	Aichi ARC	1987	1988	1802	Good taste; big seed	1151	Koomura <i>et al.</i> (1989) <sup>(82)</sup>
33	Michinoku-wase	Bi-kei No. 113*	Bi-kei No. 110**	*, **A descendant of cv. Reimei	Yamagata PAES	1988			Semi-dwarf	2994	Sato <i>et al.</i> (1988) <sup>(49)</sup>
34	K. muhikari	Shu2800/Hokuriku No. 100*	Nago-yutaka	*Gamma ray to cv. Koshihikari	Hokuriku NAES	1988	1989	2037	Short culm; good taste; lodging resistance	1160	Koga <i>et al.</i> (1989) <sup>(77)</sup>
35	Tsugaru-otome	Ouu 305	Mutsu-kaori*	*cf. cv. Mutsu-kaori; a descendant (the 2nd generation) of cv. Reimei through cv. Akibikari	Aomori PAES	1988	1990	2389	Good taste; cold tolerance; short culm	1153	Takadate <i>et al.</i> (1990) <sup>(61)</sup>
36	Heisei-mochi	Musashi-kogane*	Musashi-mochi	*cf. cv. "Musashi-kogane"; a descendant (the 2nd generation) of cv. Reimei	Saitama PAES	1988	1990	2394	Lodging resistance; resistant to stripe virus	1157	Niwayama <i>et al.</i> (1991) <sup>(27)</sup>
37	Nijihikari	Tori-kei No. 4*	Satominori	*Gamma ray to cv. "Norin No. 8"	Tottori PAES	1988	1991	2867	Short culm; lodging resistance	1163	
38	Oochikara	BG1*	Shu3116**	*A descendant (the 1st generation) of mutant cv. Taihou; **A descendant (the 3rd generation) of "R4-B" through cv. Sachiminori	Hokuriku NAES	1989	1990	2271	Large seed; tough stem and lodging resistance	2295	Kobayashi <i>et al.</i> (1990) <sup>(71)</sup>
39	Koihime	Koshikari/Reimei*	Akibikari*	*cf. cv. Reimei (Appendix I); **cf. cv. Akibikari (the 1st generation of cv. Reimei)	Aomori PAES, Fujisaka Branch	1989	1990	2329	High-yielding; cold tolerance	1161	Shimura <i>et al.</i> (1990) <sup>(52)</sup>
40	Hirohikari	Fu-kei No. 130*	Akibikari**	*cv. Koshikari/cv. Reimei/"Fu-kei No. 104"; **cf. cv. Akibikari	Hiroshima PAES	1989	1990	2392	Tough culm; high-yielding; good taste	1152	Maeda <i>et al.</i> (1990) <sup>(89)</sup>
41	Hirohonami	Fu-kei No. 130*	Yoneshiro	*cv. Koshikari/cv. Reimei/"Fu-kei No. 104"	Hiroshima PAES	1989	1990	2393	Tough culm	2996	Maeda <i>et al.</i> (1990) <sup>(89)</sup>

No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
42	Yumeminori	Eitaman No. 119	Musashi-kogane*	*cf. cv. "Musashi-kogane": a descendant (the 2nd generation) of cv. Reimei	Saitama PAES	1989	3222	Lodging resistance	1166	Niwayama <i>et al.</i> (1994) <sup>(26)</sup>
43	Koshi-no-hana	Shounai No. 32	To-kei 196*	*Gamma ray to cv. Koshihikari	Toyama PAES	1980	2869	Low amylose; panicle weight type	2997	
44	Kitaou	Fu-kei No. 108	Fu-kei No. 113*	*A descendant (the 2nd generation) of cv. Reimei	Aomori PAES, Fujisaka Branch	1990	2871	Short culm	2998	Kobayashi <i>et al.</i> (1993) <sup>(72)</sup>
45	Kaguya-mochi	Chubo37*	Fu-kei No. 126 (Kochiminori)	*A descendant (the 2nd generation) of cv. Reimei	Aomori PAES, Fujisaka Branch	1990	2872	Semi-dwarf; middle to short culm	2999	Horisue <i>et al.</i> (1992) <sup>(31)</sup>
46	Hagi-no-kaori	Tohoku No. 125	Furu2986 (cv. Miyakaori)/Tohoku No. 125	*A descendant (the 2nd generation) of cv. Reimei	Miyagi PAES, Furukawa Branch	1991	3044	Semi-dwarf	3000	Sasaki <i>et al.</i> (1994) <sup>(43)</sup>
47	Aya	Douhoku No. 43 (Naga-kei 84271)*	Kita-ake	*NM391 (a gamma ray induced mutant from cv. "Nihon-masari")/cv. Ishikari	Hokkaido AES, Kamikawa Station	1991	3283	Good taste	1169	Tanno <i>et al.</i> (1997) <sup>(73)</sup>
48	Yama-uta	Bi-kei No. 108*/Akihikari**	Fu-kei No. 127***	*, **, **A descendant of cv. Reimei	Aomori PAES, Fujisaka Branch	1991	3284	Semi-dwarf	3001	Kobayashi <i>et al.</i> (1993) <sup>(70)</sup>
49	Yukimi-mochi	Fuji-mochi712*	Fu-kei-mochi No. 119**	*, **A descendant of cv. Reimei	Aomori PAES, Fujisaka Branch	1991	3285	Semi-dwarf	3003	Horisue <i>et al.</i> (1993) <sup>(30)</sup>
50	Aneko-mochi	Fu-kei-mochi No. 133*	Fu-kei-mochi No. 119**	*, **A descendant (the 2nd generation) of cv. Reimei	Aomori PAES, Fujisaka Branch	1991	3286	Semi-dwarf	3004	Horisue <i>et al.</i> (1993) <sup>(32)</sup>
51	Yuki-gesyou	Oou No. 301	Bi-kei No. 94*	*Bi-kei 67"/cv. Reimei/cv. Sasanishiki	Yamagata PAES	1991	3344	Semi-dwarf	3005	
52	Doman-naka	Chubu No. 42*	Shoumai No. 29**	*Named as cv. "Ibuki-wase" (the 1st generation of cv. Mine-asahi), **A descendant of cv. Reimei	Yamagata PAES	1991	3345	Short stem	1168	
53	Hae-nuki	Shoumai No. 29*	Akita No. 31	*A descendant (the 3rd generation) of cv. Reimei	Yamagata PAES	1991	3346	Semi-dwarf	3006	Kyoya <i>et al.</i> (2002) <sup>(86)</sup>
54	Hatsu-nanori	Natsuhikari	Niigata-wase*	*A descendant (the 2nd generation) of cv. Reimei; cf. cv. "Niigata-wase"	Kochi PARC	1991	4051	Semi-dwarf	3007	Nakamura <i>et al.</i> (1993) <sup>(118)</sup>
55	Hashiri-aji	Niigata-wase*	Shoumai No. 32	*A descendant (the 2nd generation) of cv. Reimei; cf. cv. "Niigata-wase"	Niigata PAES	1991	5784	Semi-dwarf	3008	Hoshi <i>et al.</i> (1998) <sup>(36)</sup>
56	Ougi-wase	Inaba-wase	Yo128 (Kagahikari)*	*cf. cv. Kagahikari; a descendant (the 1st generation) of a gamma ray induced mutant of cv. Koshihikari	Ishikawa PAES	1991		Large grain; a good grain quality	3009	Matsumoto <i>et al.</i> (1993) <sup>(94)</sup>
57	Gin-no-sei	Aikawa No. 1*	Aki-kei 53	*A mutant of cv. Akiyutaka	Akita PAES	1993	3343		1167	Takahashi <i>et al.</i> (1999) <sup>(63)</sup>
58	Kinu-no-hada	Chubu-mochi No. 37	Akihikari*	*cf. cv. Akihikari; a descendant (the 1st generation) of cv. Reimei	Akita PAES	1993	3574	Tough stem; high-yielding	3010	Kato <i>et al.</i> (1995) <sup>(88)</sup>
59	Tatsuko-mochi	Chubu-mochi No. 37	Akihikari*	*cf. cv. Akihikari; a descendant (the 1st generation) of cv. Reimei	Akita PAES	1993	3575	Tough stem; high-yielding	3011	Kato <i>et al.</i> (1995) <sup>(88)</sup>
60	Ippon-jime	Gohyaku-mangoku	Houhai*	*cf. cv. Houhai; a descendant (the 1st generation) of cv. Reimei	Niigata PAES	1993	4172	Semi-dwarf	3012	Sasaki <i>et al.</i> (1994) <sup>(144)</sup>
61	Natsu-no-tayori	Fu-kei No. 125*	Ouu No. 309	*cv. Sasaminori/cv. Reimei	Kagoshima PAES	1993	4349	Semi-dwarf	3013	Yatou <i>et al.</i> (1994) <sup>(212)</sup>
62	Ume-tsukushi	Kimuhikari*	Koshihikari	*cf. cv. Kimuhikari	Fukuoka PAES	1993	4414	Good taste; lodging resistance	3014	Imabayashi <i>et al.</i> (1995) <sup>(42)</sup>

No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. Year	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
63	Inahikari	Chubu No. 35	Mine-asahi*	*A descendant (the 1st generation) of Gamma ray (20kR) induced mutant line "Kanto 79" from cv. Koshihikari	Aichi ARC	1993	1995	4415	Short culm; good taste	3015	Inoue <i>et al.</i> (1993) <sup>(63)</sup>
64	Matsuri-bare	Tsuki-no-hikari	Mine-asahi*	*A descendant of Gamma ray (20kR) induced mutant line "Kanto 79" from cv. Koshihikari	Aichi ARC	1993	1995	4416	Tough culm	3016	Syumura <i>et al.</i> (1994) <sup>(59)</sup>
65	Maihime	Ao-kei No. 69	Fu-kei No. 130*	*cv. Koshihikari/cv. Reimei//Fu-kei No. 104*	Aomori PAES, Fujisaka Branch	1993	1995	4471	Tough culm	3017	Horisue <i>et al.</i> (1994) <sup>(34)</sup>
66	Fukuhibiki	Kochihibiki	Ouu 316 gou*	*A descendant (the 3rd generation) of cv. Reimei	Tohoku NAES	1993	1995	4710	Semi-dwarf	3018	Higashi <i>et al.</i> (1994) <sup>(21)</sup>
67	Mitsu-tarou	Shimano-sakigake*	Akita-komachi	*cf. cv. "Shinano-sakigake" (a gamma ray induced mutant cultivar from cv. Toyonishiki)	Mitsui Chemicals, Inc.	1993	1996	5007		3019	
68	Owara-bijin	Shounai No. 32 (Hana-no-mai)	To-kei 196*	*A mutant strain induced through gamma ray irradiation to cv. Koshihikari	Toyama PAES	1994	1995	4409	Early-maturing, short culm	3020	Kaneda <i>et al.</i> (1996) <sup>(60)</sup>
69	Dewahikari	Shounai No. 32 (Hana-no-mai)	Ouu No. 302*	*A descendant (the 2nd generation) of cv. Reimei	Akita PAES	1994	1996	5065	Semi-dwarf	3021	
70	Aki-roman	Mine-asahi*	Nakate-shinsenbon	*A descendant (the 1st generation) of Gamma ray (20kR) induced mutant line "Kanto 79" from cv. Koshihikari	Hiroshima PAES	1994	1996	5185	Good taste	3022	Tsuchiya <i>et al.</i> (1995) <sup>(63)</sup>
71	Hoso-omote	Shin-hou No. 38 (Shimano-sakigake)*	Miyano No. 23	*cf. cv. "Shinano-sakigake" (a gamma ray induced mutant cultivar from cv. Toyonishiki)	Nagano PAES	1994	1996	5247		3023	
72	Nebari-gachi 94	Akimshiki	Tan-kei 1915*	*Named as "Noh PL No. 13" (a hybrid between two MINU induced mutants)	Plant Laboratory, Kirin Brewery Co., Ltd.	1994	1997	5844	Low amylose	3024	
73	Sai-no-yume	Kanto PL3	Tama-kei No. 74*	*cf. cv. "Yume-minori"; a descendant (the 3rd generation) of cv. Reimei	Saitama PAES	1994	1998	6445	Semi-dwarf	3025	Tokura <i>et al.</i> (1999) <sup>(77)</sup>
74	Dewa-sansan	Miyama-nishiki*	Ao-kei Sake No. 97 (Hanafubuki)**	*cf. cv. "Miyama-nishiki" (a gamma ray induced mutant cultivar from cv. "Takane-nishiki"; **cf. cv. Hanafubuki; a descendant (the 2nd generation) of cv. Reimei)	Yamagata PAES	1995	1997	5545	High % of white core rice; suitable for sake-brewery; semi-dwarf	3026	
75	Otome-gokoro	Akita-komachi	Akihikari*	*cf. cv. Akihikari; a descendant (the 1st generation) of cv. Reimei	Okayama PAES	1995	1997	5546	Semi-dwarf	3027	Nihara <i>et al.</i> (1996) <sup>(22)</sup>
76	Yume-kogane	Yuki-hikari	Hatsu-kogane*	*cf. cv. Hatsukogane; a descendant (the 2nd generation) of cv. Reimei	Aomori PAES, Fujisaka Branch	1995	1997	5785	Semi-dwarf	3028	Uehara <i>et al.</i> (1997) <sup>(86)</sup>
77	Hama-yutaka	Fu-kei No. 115×Ouu No. 321*	Fu-kei No. 140 (Kita-ou)**	*A descendant (the 3rd generation) of cv. Reimei; **cf. cv. Kitaou; a descendant (the 3rd generation) of cv. Reimei	Aomori PAES, Fujisaka Branch	1985	1997	5786	Semi-dwarf	3029	Uehara <i>et al.</i> (1997) <sup>(87)</sup>
78	Dontokoi	Kimuhikari*	Hokuriku No. 120	*cf. cv. Kimuhikari	Hokuriku NAES	1995	1997	5845	Good taste; lodging resistance	3030	Uehara <i>et al.</i> (1995) <sup>(93)</sup>
79	Soft 158	Hokuriku No. 127 (Kimuhikari)*	Ken-kei 2078**	*cf. cv. Kimuhikari; **A mutant from EMS treatment to cv. Sasanishiki	Hokuriku NAES	1995	1997	5846	Good taste; lodging resistance	3031	Uehara <i>et al.</i> (1995) <sup>(88)</sup>

No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. Year	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
80	Rinx-wase	KRN3501 (Rinx-Kobayashi)*	KRN3502	*cf. cv. "Rinx-Kobayashi" (Appendix 1)	Plant Laboratory, Kirin Brewery Co., Ltd.	1995	1998	6265	Seed-shattering resistance	3032	
81	Hoho-emi	Aichi No. 52	Mine-asahi*	*A descendant (the 1st generation) of gamma ray (20KR) induced mutant line "Kanto 79" from cv. Koshihikari	Miyazaki PAES	1995	1998	6388	Good taste; tough culm	3033	Takita <i>et al.</i> (1997) <sup>(68)</sup>
82	Shimasayaka	Tsuki-no-hikari	Kinuhikari*	*cf. cv. Kinuhikari	Plant Laboratory, Kirin Brewery Co., Ltd.	1995	1998	6444	Good taste; lodging resistance	3034	
83	Tsukushi-wase	Norin No. 22	Kinuhikari*	*cf. cv. Kinuhikari	Fukuoka PAES	1995	1999	7321	Good taste; lodging resistance	3035	Hamachi <i>et al.</i> (1998) <sup>(8)</sup>
84	Tsugaru-roman	Fu-kei No. 141*	Akita-komachi	*A descendant (the 2nd generation) of cv. Reimei	Aomori PAES	1995	2000	8280	Semi-dwarf	3036	Takadate <i>et al.</i> (1997) <sup>(60)</sup>
85	Ishikawa Sake No. 30	Gohyaku-mangoku	Hanafubuki*	*cf. cv. Hanafubuki; a descendant (the 2nd generation) of cv. Reimei	Ishikawa PAES	1996	1996	5188	Semi-dwarf	3037	
86	Asamurasaki	Tou-mochi 396	Ouu 331 gou (later named as "cv. Yumeminori**	*A descendant (the 4th generation) of cv. Reimei	Tohoku NAES	1996	1998	6504	Semi-dwarf	3038	Higashi <i>et al.</i> (1997) <sup>(24)</sup>
87	Sakitamahime	Kinuhikari*	Yumeminori**	*cf. cv. Kinuhikari; ** A descendant (the 3rd generation) of cv. Reimei through cv. "Musashi-kogane"	Saitama PAES	1996	2000	7644	Good taste; lodging resistance	3039	Tokura <i>et al.</i> (2000) <sup>(76)</sup>
88	Aki-no-sei	Toyo-nishiki	Miyama-nishiki*	*cf. cv. "Miyama-nishiki" (a gamma ray induced mutant cultivar from cv. "Takane-nishiki")	Akita PAES	1996	2000	7755	High % of white core rice; suitable for sake-brewery	3040	Masaki <i>et al.</i> (1999) <sup>(93)</sup>
89	Okini-iri	Chubu No. 47	Ouu No. 313*	*A descendant (the 3rd generation) of cv. Reimei	Tohoku NAES	1996	2000	7812	Semi-dwarf	3041	Higashi <i>et al.</i> (1997) <sup>(23)</sup>
90	Ideyut-mochi	Fuji-mochi773*	Wasetora-mochi	*A descendant (the 2nd generation) of cv. Reimei	Aomori PAES, Fujisaka Branch	1996	2000	7813	Semi-dwarf	3042	Horisue <i>et al.</i> (1997) <sup>(33)</sup>
91	Yume-musubi	Tohoku No. 137	Hokuriku No. 122 (Kinuhikari)*	*cf. cv. Kinuhikari	Miyagi PAES Furuoka Branch	1996	2000	7814	Good taste; tough culm	3043	Matsumaga <i>et al.</i> (2002) <sup>(97)</sup>
92	Asahi-no-yume	Aichi No. 70 (Aichi-no-kaori)*	Aichi No. 56/Aichi No. 65**	* ** - A descendant (the 2nd generation) of a gamma ray induced mutant line "Kanto 79 gou" through "Mine-asahi"	Aichi ARC	1996	2000	7888	Good taste; tough culm	3044	Izawa <i>et al.</i> (2001) <sup>(50)</sup>
93	Yume-izumi	Yumehikari	Kinuhikari*	*cf. cv. Kinuhikari	Kumamoto PARC	1996	2000	8124	Good taste; lodging resistance	3045	Izumi <i>et al.</i> (1998) <sup>(54)</sup>
94	Yume-hitachi	Chiyo-nishiki	Hokuriku No. 122 (Kinuhikari)*	*cf. cv. Kinuhikari	Ibaraki PAES	1996	2000	8213	Good taste; lodging resistance	3046	Suga <i>et al.</i> (2000) <sup>(55)</sup>
95	Rinx-nakate	Rinx Kobayashi*	KRN3505	*cf. cv. "Rinx-Kobayashi" (Appendix 1)	Plant Laboratory, Kirin Brewery Co., Ltd.	1996	2000	8359	Non seed shattering	3047	
96	Iwata No. 12	Asominori	Aichi-no-kaori*	*cf. cv. "Aichi-no-kaori"	Japan Tobacco, Inc.	1996	2001	8631	Semi-dwarf	3048	
97	Hoshi-akari	Hatsuboshi	Miyama-nishiki*	*cf. cv. "Miyama-nishiki" (a gamma ray induced mutant cultivar from cv. "Takane-nishiki") (Appendix 1)	Kako-mai Breeding Institute, Inc. and Tohoku Electric Power Co., Inc.	1996	2001	9116	High % of white core rice; suitable for sake-brewery	3049	

No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. No. <sup>a</sup>	Reg. Year	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
98	Hayatsukushi	Hatsuboshi	<b>Seinan No. 89*</b>	*Named as cv. "Natsu-no-tayori": a descendant (the 2nd generation) of cv. Reimei	Fukuoka ARC	1997	8360	2000	Semi dwarf	3050	Hamachi <i>et al.</i> (1998) <sup>(7)</sup>
99	Aki-geshiki	<b>Kyu-kei 919*</b>	Himohikari	*A descendant of cv. "Mine-asahi"	Miyazaki PAES	1997	8544	2000	Good taste; tough culm	3051	The Miyazaki Breeding Group of Rice (2005) <sup>(74)</sup>
100	Oyama-nishiki	Hida-homare	<b>Akita Sake No. 33*</b>	*A descendant of cv. "Miyama-nishiki" (Appendix 1)	Toyama PAES	1997	9029	2001	High % of white core rice; suitable for sake-brewery	3052	Kaneda <i>et al.</i> (1999) <sup>(59)</sup>
101	Hatajirushi	Tohoku No. 141	<b>Tohoku No. 142*</b> (Ibukiwase/Tohoku No. 125)	*cv. Ibukiwase: a descendant (the 2nd generation) of "Kanto 79 gou" through cv. "Mine-asahi"	2nd Miyagi PAES Furukawa Branch	1997	8832	2001	High % of white core rice; suitable for sake-brewery	3053	Matsumaga <i>et al.</i> (2002) <sup>(98)</sup>
102	Banbanzai	Chiyo-nishiki	<b>Tama-kei No. 74 (cv. Yume-minori)*</b>	*cf. cv. "Yume-minori": a descendant (the 3rd generation) of cv. Reimei through cv. "Musashi-kogane"	Saitama PAES	1997	9304	2001	Semi-dwarf	3054	Tokura <i>et al.</i> (2002) <sup>(75)</sup>
103	Snow Pearl	<b>74wx2N1*</b>	<b>Reimei**</b>	*A mutant of "Norin No. 8"; **cf. cv. Reimei (Appendix 1)	Tohoku NAES	1998	7062	1999	Low amylose; semi-dwarf	3055	Higashi <i>et al.</i> (1998) <sup>(22)</sup>
104	Koimomiji (Hiroshima No. 21)	Sachi-izumi	<b>Fu-kei No. 141*</b>	*A descendant (the 2nd generation) of Hiroshima PARC cv. Reimei	Hiroshima PARC	1998	9301	2001	Semi-dwarf	3056	Maeda <i>et al.</i> (2000) <sup>(90)</sup>
	Datchi-no-kaze	<b>A-kei 558</b> (Matsuribare)*	Aichi No. 78/4/Tsuki-no-hikari/3/Aichi No. 77//Iku D759/A-kei	*cf. cv. Matsuribare: a descendant (the 2nd generation) of a gamma ray induced mutant line "Kanto 79 gou" through cv. "Mine-asahi"	Aichi ARC	1998	8640	2002	Good taste; tough culm	3057	Izawa <i>et al.</i> (2001) <sup>(52)</sup>
105											
106	Koshihikari Toyama BL No. 1	Koshihikari	<b>Koshihikari/BC-ta2*</b>	*Kinuhikari2/Tohoku IL No. 7"	Toyama ARC	1998	9641	2002	Good taste; lodging resistance	3058	Kojima <i>et al.</i> (2003) <sup>(79)</sup>
107	Hanabusa	<b>Douhoku No. 53*</b>	Kita-ake	*A sister line of cv. Aya; a descendant (the 2nd generation) of a gamma ray induced mutant line "NM391"	Hokkaido NAES	1998	9785	2002	Low amylose	3059	Araki <i>et al.</i> (2002) <sup>(8)</sup>
108	Sawa-pikari	<b>Tan-kei 2018*</b>	Asa-no-hikari	*A mutant line through MNU treatment to cv. Kochihibiki	Gunma PAES	1998	9786	2002	Low amylose	3060	Narizuka <i>et al.</i> (1998) <sup>(20)</sup>
109	Awaminori	Hokkai No. 132	<b>Hokuriku No. 122</b> (Kinuhikari)*	*cf. cv. Kinuhikari	Hokuriku NAES	1998	9788	2002	Good taste; lodging resistance	3061	Uehara <i>et al.</i> (1998) <sup>(92)</sup>
110	Haiminori	<b>EM40*</b>	Ake-no-hoshi	*A mutant from MNU treatment to cv. Kinmaze	Chugoku NAES	1999	8008	2000	Giant embryo	3062	Nemoto <i>et al.</i> (2001) <sup>(21)</sup>
111	Yume-ohmi	<b>Etsunan No. 135*</b>	Shi-kei No. 51	*A descendant (the 3rd generation) of cv. Reimei through cv. Akihikari.	Shiga PAES	1999	8944	2001	Semi-dwarf	3063	Noda <i>et al.</i> (1999) <sup>(28)</sup>
112	Mie-no-emi	<b>Yamagata No. 41*</b>	Hitomebore	*"Shounai No. 32"/cv. "Mine-asahi" (a descendant (the 1st generation) of Gamma ray (20kR) induced mutant line "Kanto 79" from cv. Koshihikari)	Mie PARC	1999	9911	2002	Good taste; tough culm	3064	Yamakawa <i>et al.</i> (2000) <sup>(207)</sup>
113	Yume-akari	Akita-komachi	<b>Ao-kei No. 110*</b>	*A descendant (the 4th generation) of cv. Reimei through cv. Mutsukaori	Aomori PAES	1999	9912	2002	Semi-dwarf	3065	Mikami <i>et al.</i> (2000) <sup>(100)</sup>

No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. No. <sup>a</sup>	Reg. Year	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
114	Gin-ginga	Yamagata Sake No. 49*	Akita Sake No. 49**	*Named as cv. "Dewa-sansan" (a descendant (the 3rd generation) of cv. Reimei); **a descendant (the 3rd generation) of cv. Reimei through cv. Akihikari	Iwate PAES, Kerman Station	1999	2002	1047	Semi-dwarf	3066	Odanaka <i>et al.</i> (2000) <sup>(29)</sup>
115	Yume-no-kaori	Hattan-nishiki No. 1	Yamagata Sake No. 49*	*Named as cv. "Dewa-sansan" (a descendant (the 3rd generation) of cv. Reimei)	Fukushima PAES	1999	2003	1096	Semi-dwarf	3067	Sato <i>et al.</i> (2003) <sup>(48)</sup>
116	Aichi-no-kaori SBL	Aichi-no-kaori*	F3 (Aichi-no-kaori*2//Aichi No. 78/Aichi-no-kaori*)	*cf. cv. "Aichi-no-kaori"	Aichi ARC	1999	2003	10968	Good taste; big seed	3068	Izawa <i>et al.</i> (2001) <sup>(51)</sup>
117	Oku-no-murasaki	Tohoku Mochi 149 gou	Ouu 331 gou (later named as cv. Fukuhibiki)*	*A descendant (the 4th generation) of cv. Reimei	NARO NARC Tohoku Region	1999	2003	11088	Semi-dwarf	3069	Yokogami <i>et al.</i> (2000) <sup>(213)</sup>
118	Oshimako 180	Douhoku No. 43 (NM391*/Ishikari)	Fu-kei No. 144**	*Gamma ray to cv. "Nihon-masari"; **A descendant (the 3rd generation) of cv. Reimei	Aomori PAES	1999	2003	11230	Semi-dwarf	3070	
119	Koshi-ibuki	Tohoku No. 143 (Hitomebore)	Yamagata No. 35*	*Named as cv. "Doman-naka" (a descendant (the 3rd generation) of a gamma ray induced mutant line "Kanto 79 gou" through cv. "Mine-asahi" from cv. Koshihikari)	Niigata PAES	1999	2003		Good taste; tough culm		Hoshi <i>et al.</i> (2002) <sup>(37)</sup>
120	Yawarakomaichi	Nishihomare+B140	Tan-kei 2021*	*A mutant line through NMU treatment to cv. Kinmaze	Kyushu NAES	1995	1999	7084	Low amylose content	3072	Okamoto <i>et al.</i> (2001) <sup>(31)</sup>
121	Kan-no-mai	Gohyaku-mangoku	Miyama-nishiki*	*cf. cv. "Miyama-nishiki" (Appendix 1)	Shimane PAES	1995	2000	7890	High % of white core rice; suitable for sake-brewery	3073	Yamamoto <i>et al.</i> (1999) <sup>(39)</sup>
122	Hanakirari	PR3*	Koshihikari	*A mutant from protoplast culture of cv. Nipponbare	Plantec Research Institute	2000	2000	8545	Early-maturing	3074	
123	Chiho-no-kaori	Tohoku No. 144 (Hagi-no-kaori)*	Nankai No. 122	*cf. cv. "Hagi-no-kaori" (a descendant (the 4th generation) of cv. Reimei.)	Miyazaki PAES	2000	2002	1087	Semi-dwarf	3075	Takita <i>et al.</i> (2000) <sup>(69)</sup>
124	Gin-otome	Akita Sake No. 44*	Tohoku No. 141	*Gamma ray to F6 ("Hyo-kei-sake No. 16"/cv. Yoneshiro)	Iwate PAES	2000	2003	1096	Short culm	3076	Sugawara <i>et al.</i> (2012) <sup>(56)</sup>
125	Yume-ippai	Yume-gokochi*	Yume-kahori**	*cf. cv. "Yume-gokochi"; **cf. cv. "Yume-kahori"	Mitsui Chemicals, Inc.	2000	2003	10965	Short culm	3077	
126	Yume-sayaka	Yamagata No. 40*	Oou No. 341	*A descendant (the 3rd generation) of cv. Reimei	Yamagata PAES	2000	2003	10966	Semi-dwarf	3078	Sato <i>et al.</i> (2000) <sup>(45)</sup>
127	Itadaki	Shu485 (Dontokoi)*	Shu4695	*cf. cv. Dontokoi; a descendant (the 2nd generation) of cv. Kinuhikari	NARO (NARC, Hokuriku Center)	2000	2003	11087	Good taste; lodging resistance	3079	Uehara <i>et al.</i> (2000) <sup>(95)</sup>
128	Mine-hibiki	Sachi-izumi	Mine-asahi*	*cf. cv. "Mine-asahi"; A descendant (the 1st generation) of gamma ray induced mutant line "Kanto No. 79" from cv. Koshihikari	Aichi ARC	2000	2003	11089	Good taste; tough culm	3080	Kudo <i>et al.</i> (2000) <sup>(84)</sup>

No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. Year	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
129	Chiyo-no-mochi	Ouu-mochi No. 317* Fu-kei No. 141**	Fu-kei No. 143***	**A descendant (the 3rd generation) of cv. Reimei; ***Named as cv. "Yama-uta" (a descendant (the 4th generation) of cv. Reimei)	Aomori PAES, Fujisaka Branch	2000	2003	11091	Semi-dwarf	3081	Uehara <i>et al.</i> (2001) <sup>(96)</sup>
130	Milky Princess	Kanto No. 163	Kou272 (Milky Queen)*	*cf. cv. "Milky Queen" (Appendix 1)	NARO (National Institute of Crop Science)	2000	2003	11234	Low amylose content	3082	Sato <i>et al.</i> (2008) <sup>(146)</sup>
131	Koshi-no-shizuku	Hyogo-kita-nishiki*	Miyama-nishiki**	*cf. cv. "Hyo-kei-sake No. 18"; **cf. cv. "Miyama-nishiki" (Appendix 1)	JA, Teraru-Echizen	2000	2003	11361		3083	
132	Yume-shizuku	Kimuhikari*	Tohoku No. 143 (Hitomebore)	*cf. cv. Kimuhikari	Saga PARC	2000	2003	11358	Good taste; lodging resistance	3084	Hirota <i>et al.</i> (2001) <sup>(28)</sup>
133	Fusa-no-mai	Sirotate-nishiki	Chubu No. 72*	*A descendant of cv. Reimei (the 3rd generation) and cv. "Mine-asahi" (the 2nd generation)	Chiba PARC	2000	2004	11583	Good taste; tough culm	3085	Wada <i>et al.</i> (2002) <sup>(97)</sup>
134	Fukumirai	Chubu No. 82*	Chiyo-nishiki	**Chubu No. 44"/cv. Mine-asahi; cf. cv. "Mine-asahi"	Fukushima PAES	2000	2004	11842	Good taste; tough culm	3086	Sato <i>et al.</i> (2006) <sup>(147)</sup>
135	Ten-no-midori	PR3*	Koshihikari	*A mutant from protoplast culture of cv. Nipponbare	Mitsubishi Chemical Co.	1996	2001	8833	Early-maturing	3087	
136	Shun-you	LGC-1*	Hokuriku No. 153**	*cf. cv. LGC1 (Appendix 1); **cv. Oochikara (the 2nd generation) of cv. Taihou. x cv. Kochihibiki	NARO (NARC, Hokuriku Center)	2001	2004	12181	Low gluten content in the grain	3088	Uehara <i>et al.</i> (2002) <sup>(94)</sup>
137	Silky Pearl	Tan-kei2019 (Noh PL No. 14)*	Fu-kei No. 143 (cv. Yamauta)**	*A mutant from NMU treatment to cv. Kochihibiki; **cf. cv. Yamauta (the 3rd generation of cv. Reimei)	NARO (NARC for Tohoku Region)	2001	2004	12274	Semi-dwarf	3089	Takita <i>et al.</i> (2002) <sup>(67)</sup>
138	Asa-isuyu	Hokuriku No. 127 (Kimuhikari)*	Douhoku No. 43 (NM391**/Ishikari)	*A descendant (the 2nd generation) of cv. Reimei; **Gamma ray irradiation to cv. Nihonmasari	NARO (NARC, Hokuriku Center)	2001	2004	12180	Semi-dwarf; low amylose content	3090	Uehara <i>et al.</i> (2002) <sup>(91)</sup>
139	Takitate	Ouu No. 343*	Tohoku No. 153	*74wx2N1/Reimei	Miyagi PAES Furukawa Branch	2001	2004	12055	Semi-dwarf	3091	Nagano <i>et al.</i> (2005) <sup>(11)</sup>
140	Yume-no-hana	Yume-gokochi*	Yume-kahori**	*cf. cv. "Yume-gokochi"; **cf. cv. "Yume-kahori"	Plantec Research Institute	2001	2004	12045	Low amylose contents; short stem	3092	
141	Kahoruko	Kanto No. 154 (Sally Queen)	Akihikari*	*cf. cv. Akihikari (a descendant (the 1st generation) of cv. Reimei)	Niigata ARI	2001	2004	11843	Semi-dwarf	3093	
142	Akigumo	Niigata-wase*	Tan-kei 2019**	*cf. cv. "Niigata-wase"; **Named as "Noh PL No. 14"	Niigata ARI	2001	2004	11844	Low amylose content	3094	
143	Natsugumo	Yuki-no-sei	Aya*	*cf. cv. Aya	Niigata ARI	2001	2004	11845		3095	
144	Shithou	Wataboushi	Asamurasaki*	*A descendant of cv. Reimei (the 5th generation)	Niigata ARI	2001	2004	11846	Semi-dwarf	3038	
145	Bemika	Niigata-mochi No. 31/Shimonoi	Niigata-mochi No. 31/Tohoku No. 144 (Hagi-no-kaori)*	*cf. cv. "Hagi-no-kahori" (a descendant (the 4th generation) of cv. Reimei)	Niigata ARI	2001	2004	11847	Semi-dwarf	3097	
146	Bemisarasa	Niigata-mochi No. 31/Shimonoi	Niigata-mochi No. 31/Tohoku No. 144 (Hagi-no-kaori)*	*cf. cv. "Hagi-no-kahori" (a descendant (the 4th generation) of cv. Reimei)	Niigata ARI	2001	2004	11848	Semi-dwarf	3098	

No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. Year	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
147	Natsushizuka	Kanto No. 136/Koshihikari	Natsuhikari*	*cf. cv. Natsuhikari; a descendant (the 4th generation) of a gamma ray induced mutant line "Fu-kei 71 gou"	Shizuoka PAES	2001	2005	12562	Semi-dwarf	3099	Miyata <i>et al.</i> (2001) <sup>(106)</sup>
148	Toji-no-hana (Hyo-kei-sake No. 65)	Hyo-go-kitanishiki*	Hidomare	*cf. cv. "Hyo-go-kita-nishiki"; a descendant (the 4th generation) of cv. Reimei	Hyo-go PAFFRC	2001	2004	12178	Semi-dwarf	3100	
149	Toji-no-yume (Hyo-kei-sake No. 66)	Hyo-go-kitanishiki*	Gin-no-sei**	*cf. cv. "Hyo-go-kita-nishiki"; a descendant (the 4th generation) of cv. Reimei; **cf. cv. "Gin-no-sei"	Hyo-go PARC	2001	2004	12179	Semi-dwarf	3101	
150	Nebari-goshi	21-3-6*	Kinuhikari**	*A mutant of cv. "Todoroki-wase"; **cf. cv. Kinuhikari	Nagano PAES	1997	2002	9644	Good taste; lodging resistance	3102	
151	Koshihikari Niigata BL No. 4 wase*	Koshihikari/Niigata-wase*	Koshihikari	*A descendant (the 2nd generation) of Niigata ARI cv. Reimei	Niigata ARI	1998	2002	1023	Semi-dwarf	3103	Ishizaki, K. (2007) <sup>(46)</sup>
152	Misato-nishiki	Yamada-nishiki	Miyama-nishiki*	*cf. cv. "Miyama-nishiki" (Appendix)	Akita PAES	1998	2002	1023		3104	Masaki <i>et al.</i> (2004) <sup>(92)</sup>
153	Coimustubi	Chubu No. 73*	Tohoku No. 143 (Hitomebore)	*Chubu No. 44/Mine-asahi; cf. cv. "Mine-asahi"	Miyagi PAES Furukawa Branch	1998	2002	1036		3105	Nagano <i>et al.</i> (2005) <sup>(110)</sup>
154	Hoshi-aoba	Tashu-kei 174 (Chugoku No. 113)	Hokuriku No. 130 (Oochikara)*	*cf. cv. Oochikara	NARO (NARC for Chugoku and Shikoku Region)	2000	2003	11360		3106	Maeda <i>et al.</i> (2003) <sup>(88)</sup>
155	LGC Soft	NM391*	LGC-1**	*A mutant line induced through gamma ray irradiation to cv. Nihonmasari; **cf. cv. "LGC-1"; a mutant cultivar induced through EI treatment to cv. Nihonmasari (Appendix I)	NARO (NARC for Chugoku and Shikoku Region)	2002	2005	12564	Low amylose $\beta$ ; low glutenin $\sigma$	3107	Iida <i>et al.</i> (2004) <sup>(41)</sup>
156	Mebae-mochi	EM40*	Chubu-mochi No. 57	*A mutant from NMU treatment to cv. Kinmaze	NARO (NARC, Hokuriku Center)	2002	2005	13188	Giant embryo	3108	Uehara <i>et al.</i> (2003) <sup>(89)</sup>
157	Kusa-yutaka	Chugoku No. 105	Hokuriku No. 130 (Oochikara)*	*cf. cv. Oochikara	NARO (NARC, Hokuriku Center)	2002	2005	13189		3109	Uehara <i>et al.</i> (2003) <sup>(90)</sup>
158	Benigoromo	Fukuhibiki*/A5	Ouu 331 gou (later named as cv. Fukuhibiki)**	* **A descendant of cv. Reimei (the 4th generation)	NARO (NARC, Tohoku Region)	2002	2005	13186	Semi-dwarf	3110	Yamaguchi <i>et al.</i> (2005) <sup>(104)</sup>
159	Akisayaka	Saikai No. 195	Hokuriku No. 148 (Dontokoi)*	*cf. cv. Dontokoi; a descendant (the 4th generation) of a gamma ray induced mutant line "Hokuriku 100 gou" through cv. Kinuhikari	NARO (NARC for Kyushu and Okinawa Region)	2002	2005	13187	Good taste; lodging resistance	3111	Okamoto <i>et al.</i> (2008) <sup>(132)</sup>
160	Mie-no-yume	Aichi No. 92 (Matsuribare)*	Etsunan No. 148	*cf. cv. Matsuribare	Mie PARC	2002	2004	12056		3112	Yamakawa <i>et al.</i> (2002) <sup>(206)</sup>
161	Tsukushi-roman	Chikushi No. 6 (Yume-tsukushi)*	Chubu No. 88 (Mochi-kei 347/Mine-asahi**)	*cf. cv. "Yume-tsukushi"; **cf. cv. "Mine-asahi"	Fukuoka ARC	2002	2005	12959		3113	Hamachi <i>et al.</i> (2003) <sup>(93)</sup>
162	Tabegokochi	Dontokoi*	Koshihikari	*cf. cv. Dontokoi	Monsanto Japan Limited	2002	2005	12827	Good taste; lodging resistance	3114	
163	Tone-no-megumi	Dontokoi*	Koshihikari	*cf. cv. Dontokoi	Monsanto Japan Limited	2002	2005	12828	Good taste; lodging resistance	3115	



No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. Year	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
164	Hanaemaki	Douboku No. 53*/Noh PL No. 11	Kuu-iku No. 139 (Yukimaru)	*A mutant of "Nihon-masari" (NIM391)	NARO (NARC for Hokkaido Region)	2003	2006	14032	Low amylose content	3116	
165	Oborozuki	Kuu-iku No. 150 (Akiho)	Hokkai No. 287*	*in vitro variation of cv. "Kirara 397"	NARO (NARC for Hokkaido Region)	2003	2006	14033		3117	Ando et al. (2006) <sup>4)</sup>
166	Churahikari	Tohoku No. 143 (Hitomebore)	Ouu No. 338*	*A descendant (the 2nd generation) of cv. "Mine-asahi"	NARO (NARC for Tohoku Region)	2003	2006	14034		3118	Yamaguchi et al. (2005) <sup>3)</sup>
167	Sakihikari	Hinohikari	Kinuhikari*	*cf. cv. Kinuhikari	Fukui PAES	2003	2006	13875	Good taste; lodging resistance	3119	Horiuchi et al. (2004) <sup>3)</sup>
168	LGC-Katsu	LGC-1*	89WPKG30-433**	*cf. cv. "LGC-1" (Appendix 1); **A mutant with 26Da globulin deletion induced by gamma ray irradiation to WPK	NARO (National Institute of Crop Science) and NIAS (IRB)	2004	2006	13871	Easy-to-digest protein 50 % decrease	3120	Nishimura et al. (2005) <sup>24)</sup>
169	LGC-Jun	LGC-1*	89WPKG30-433**	*cf. cv. "LGC-1" (Appendix 1); **A mutant with 26Da globulin deletion induced by gamma ray irradiation to WPK	NARO (National Institute of Crop Science) and NIAS (IRB)	2004	2006	13872	Easy-to-digest protein 50 % decrease	3121	Nishimura et al. (2005) <sup>24)</sup>
170	Hana-omoi	Yamada-nishiki	Hanaufubuki*	*A descendant (the 2nd generation) of cv. Reimei	Aomori PAES	2002	2006	13863	Semi-dwarf	3122	Mikami et al. (2003) <sup>102)</sup>
171	Sai-no-kagayaki	Aichi No. 92 (Matsuribare)*	Tama-kei No. 88 (Sai-no-yume)**	*cf. cv. Matsuribare; **cf. cv. "Sai-no-Saitama PAERI yume"	NARO (NARC for Aichi Prefecture)	2002	2005	12699		3123	Arakawa et al. (2003) <sup>7)</sup>
172	Sai-no-krabiyaka	Aichi No. 92 (Matsuribare)*	Tama-kei No. 88 (Sai-no-yume)**	*cf. cv. Matsuribare; **cf. cv. "Sai-no-Saitama PAES yume"	NARO (NARC for Aichi Prefecture)	2002	2005	12700		3124	Arakawa et al. (2003) <sup>7)</sup>
173	Fuku-izumi	Saikai No. 199	Dontokoi*	*cf. cv. Dontokoi	NARO (NARC for Kyushu and Okinawa Region)	2004	2007	14889	Good taste; lodging resistance	3125	Kaji et al. (2006) <sup>56)</sup>
174	Ikuhikari	Etsunan No. 148	Hokuriku No. 148 (Dontokoi)*	*cf. cv. Dontokoi	Fukui PAES	2004	2007	14999	Good taste; lodging resistance	3126	Tomita et al. (2005) <sup>82)</sup>
175	Miya-yutaka	Nankai No. 133	Saikai No. 215 (Yawarakomachi)*	*cf. cv. Yawarakomachi	Miyazaki ARI	2004	2007	15000	Low amylose content	3127	Kato et al. (2006) <sup>64)</sup>
176	Aki-neiro	Koganebare	Milky Queen (Kanto No. 168)*	*A mutant cultivar with low amylose content induced by a chemical mutagen (NMU) + another culture; cf. cv. "Milky Queen" (Appendix 1)	Kumamoto PARC	2001	2005	12566		3128	Mitsukawa et al. (2002) <sup>103)</sup>
177	Seiki-wan	Koshihikari	Haenuki*	*A descendant (the 4th generation) of cv. Reimei	Mr. Masaki OIKAWA	1998	2003	11232	Semi-dwarf		
178	Komurasaki	Tohoku-mochi 149 gou	Tatsuko-mochi*	*A descendant (the 2nd generation) of cv. Reimei through cv. Akihikari	Akita PAES	2000	2004	11841	Semi-dwarf		Matsumoto et al. (2006) <sup>96)</sup>
179	Hikari-shinseiki	Kanto 79 gou*	Jikkoku	*A mutant line induced through gamma ray irradiation to the seed of cv. Koshihikari	Toitiori University	1999	2004	12273	Semi-dwarf		Tomita (2006) <sup>178)</sup>
180	Kaze-naruko	Tsuyuhakaze	Ippon-jime*	*A descendant (the 2nd generation) of cv. Reimei through cv. Houhai	Kochi PARC	2001	2005	12563	Semi-dwarf		Mizobuchi et al. (2003) <sup>107)</sup>
181	Chigo-no-hoho	Wataboushi	Hagi-no-kaori*	*A descendant (the 3rd generation) of cv. Reimei through cv. Houhai	Niigata PAES	2001	2005	12702	Semi-dwarf		Kobayashi (2008) <sup>73)</sup>

No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. Year	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
182	Ekkakou	Todoroki-wase	Hagi-no-kaori*	*A descendant (the 3rd generation) of Niigata PAES cv. Reimei through cv. Houhai	Niigata PAES	2001	2005	12703	Semi-dwarf		Kobayashi (2008) <sup>(73)</sup>
183	Koshiguruma	Hokkai 269 gou*	Niigata wase**	*A descendant of a mutant line "Tankei 2006", whose embryo is very large, that is induced through gamma ray irradiation to cv. Kinmaze; **A descendant (the 2nd generation) of cv. Reimei	Niigata PAES	2001	2005	12704	Large embryo		
184	Akita 63 gou	Oochikara*	Akita 39 gou	*A hybrid variety between 2 mutant lines, cf. cv. Oochikara	Akita PAES	2001	2005	12826			Kodama <i>et al.</i> (2014) <sup>(75)</sup>
185	Fukkura-momoko	Kimuhikari*	Koshihikari	*A mutant cultivar induced through gamma ray irradiation to cv. Koshihikari	Okayama PAES	2002	2005	13297	Good taste; lodging resistance		Hihara <i>et al.</i> (2004) <sup>(25)</sup>
186	Kinpika	An original line	Haemuki*	*A descendant (the 4th generation) of cv. Reimei	Ibigawa Kougyo Co. Ltd.	2002	2006	13636	Semi-dwarf		
187	Hana-hyogo	Hatsugozen	Kimuhikari*	*cf. cv. Kimuhikari	Hyogo PARC	2002	2006	13742	Good taste; lodging resistance		
188	Fuyu-geshiki	Yamagata 40 gou*	Ao-kei 113 gou**	*A descendant (the 3rd generation) of cv. Reimei through cv. Akihikari; **A descendant (the 3rd generation) of cv. Reimei	Aomori PAES	2002	2006	13867	Semi-dwarf		Mikami <i>et al.</i> (2004) <sup>(99)</sup>
189	Koma-no-mai	Yamagata 40 gou*	Fu-kei 164 gou**	*A descendant (the 3rd generation) of cv. Reimei through cv. Akihikari; **A descendant (the 3rd generation) of cv. Reimei through cv. Akihikari	Aomori PAES	2003	2006	13874	Semi-dwarf		Sunohara <i>et al.</i> (2004) <sup>(58)</sup>
190	Yume-mirai	Yume-gokochi*	Yume-kahori**	*A mutant cultivar induced through a protoplast somaclonal variation from cv. Koshihikari; **A mutant cultivar induced through a protoplast somaclonal variation from cv. "Tsuki-no-hikari"	Plantec Research Institute Ltd.	2003	2006	14036	Low amylose content		
191	Tenshi-no-uta	Saikai 201 gou*	Kanto 165 gou**	*A descendant of cv. "Mine-asahi"; **A descendant of cv. Kimuhikari	Saga PARC	2001	2006	14296	Good taste; lodging resistance		
192	Yume-ikkon	Hokuriku 160 gou*	Yume-tsukushi**	*A descendant (the 3rd generation) of cv. Reimei through cv. "Niigata-wase"; **A descendant (the 2nd generation) of cv. Kimuhikari	Fukuoka PAES	2002	2006	14525	Semi-dwarf; good taste; lodging resistance		Hamachi <i>et al.</i> (2004) <sup>(20)</sup>
193	Nasuhikari	Koshihikari	Aichi 87 gou*	*A descendant of cv. "Mine-asahi"	Tochigi PAES	2004	2007	14775			Izawa <i>et al.</i> (2005) <sup>(53)</sup>
194	Hyogo-yume-otome	Kimuhikari*	Aoi-no-kaze	*cf. cv. Kimuhikari	Hyogo PARC	2003	2007	14776	Good taste; lodging resistance		
195	Chiba 28 gou	Chubu 64 gou*	Fusa-otome	*A descendant of cv. "Mine-asahi"	Chiba PARC	2003	2007	14882			Nishikawa <i>et al.</i> (2006) <sup>(23)</sup>
196	Koi-honoka	An original line	Yamauta*	*A descendant (the 3rd generation) of cv. Reimei through cv. Akihikari	Aomori PAES	2004	2007	14884	Semi-dwarf		Mikami <i>et al.</i> (2007) <sup>(10)</sup>
197	Yuki-no-hana	Oshimako180*	Kakehashi	*A hybrid between cv. Reimei and a mutant line of "Nihon-masari"	Aomori PAES	2003	2007	14886	Semi-dwarf		Sunohara <i>et al.</i> (2007) <sup>(57)</sup>

No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. Year	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
198	Yume-aoba	An indica-japonica hybrid line	Fukuhibiki*	*A descendant (the 4th generation) of cv. Reimei	NARO Hokuriku ARC	2004	2007	14887	Semi-dwarf		Miura <i>et al.</i> (2006) <sup>(16)</sup>
199	Nishi-aoba	Oochikara*	Hitohana	*cf. cv. Oochikara	NARO Kyushu Okinawa ARC	2004	2007	14888	Semi-dwarf	3325	Tamura <i>et al.</i> (2007) <sup>(17)</sup>
200	Dewa-no-sato	Gin-fubuki	Dewa-sansan*	*cf. "Dewa-sansan": a descendant of cv. Reimei (the 3rd generation) through cv. Hanafubuki and cv. "Miyama-nishiki" (the 2nd generation)	Yamagata PAES	2004	2007	15119	Semi-dwarf	3361	Yuki <i>et al.</i> (2006) <sup>(215)</sup>
201	Sato-no-yuki	Shou 1658*	Yamagata 63 gou**	*A descendant of a somaclonal variation from the anther culture of cv. "Akitia-komachi"; **A descendant of cv. Kinuhikari	Yamagata PAES	2004	2007	15534	Good taste; lodging resistance		
202	Lake 65	Hinohikari	Kinuhikari*	*cf. cv. Kinuhikari	Shiga PAES	2004	2008	16011	Good taste; lodging resistance		Nakagawa <i>et al.</i> (2005) <sup>(115)</sup>
203	Yukinko-mai	Doman-naka*	Yukinosei	*A descendant (the 2nd generation) of cv. "Mine-asahi"; a descendant (the 3rd generation) of cv. Reimei	Niigata PAES	2004	2008	16012	Semi-dwarf		Ishizaki <i>et al.</i> (2008) <sup>(47)</sup>
204	Fuku-okoshi	Hoso-omote*	Fukuhibiki**	*A descendant of cv. "Shinano-sakigake"; a mutant through gamma ray irradiation to cv. Toyonishiki; **A descendant (the 4th generation) of cv. Reimei	Nagano PAES	2004	2008	16288	Semi-dwarf		
205	Yume-kanae	LGC1*	Hitomebore	*cf. cv. LGC1 (Appendix 1)	Chiba PARC	2004	2008	16289	Low gluten content in the grain		
206	Awayuki-komachi	Ouu 343 gou*	Dewahikari**	*A descendant (the 1st generation) of cv. Reimei; **A descendant (the 3rd generation) of cv. Reimei	Akita PAES	2002	2008	16290	Semi-dwarf		Kodama <i>et al.</i> (2010) <sup>(76)</sup>
207	Himuka-mochi	Mine-no-yuki-mochi*	Miyazaki-mochi	*A descendant of cv. Reimei through cv. Hayahikari	Miyazaki PAES	2005	2008	16466	Semi-dwarf		
208	Dompishyari	Gan-nan 7 gou	Fu-kei 179 gou*	*A descendant (the 4th generation) of cv. Reimei through cv. Hatsukogane	Iwate PAES	2005	2008	16604	Semi-dwarf		Tamura <i>et al.</i> (2007) <sup>(70)</sup>
209	Akimasari	Manpuku-mochi	Akisayaka*	*A descendant (the 2nd generation) of cv. Kinuhikari through cv. Dontokoi	NARO Kyushu Okinawa ARC	2006	2008	16606	Good taste; lodging resistance		Sakai <i>et al.</i> (2006) <sup>(138)</sup>
210	Ayu-no-hikari	EMS*	Fukuhibiki**	*A mutant induced through NMU treatment to cv. Kinmaze; **A descendant (the 4th generation) of cv. Reimei	NARO Hokuriku ARC	2005	2008	16607	Semi-dwarf		Miura <i>et al.</i> (2007) <sup>(64)</sup>
211	Kinumusume	Kinuhikari*	Matsuribare**	*cf. cv. Kinuhikari; **A descendant of cv. "Mine-asahi"	NARO Kyushu Okinawa ARC	2005	2008	16609	Good taste; lodging resistance		Kaji <i>et al.</i> (2009) <sup>(57)</sup>
212	Beko-aoba	Oochikara*	Saikai 203 gou	*cf. cv. Oochikara	NARO Tohoku ARC	2005	2008	16610		3320	Nakagami <i>et al.</i> (2006) <sup>(116)</sup>
213	Nikomaru	Kinumusume*	Hokuriku 174 gou**	*A descendant of cv. Kinuhikari; **A descendant of cv. Kinuhikari through cv. Dontokoi	NARO Kyushu Okinawa ARC	2005	2008	16611			Sakai <i>et al.</i> (2010) <sup>(40)</sup>

No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. Year	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
214	Hikarikko	Koshihikari	Reimei*	*cf. cv. Reimei (Appendix I); A mutant cultivar induced through a gamma ray irradiation to cv. Fujiminori	Mr. Masayuki MURAI	2009	2009	18110	Semi-dwarf		
215	Nangoku-sodachi	Kouiku 30 gou*	Kou-kei 265**	*A descendant of cv. Reimei through cv. Hayahikari; **A descendant of cv. "Katsura-wase", cf. cv. "Katura-wase"	Kochi PARC	2009	2009	18117	Semi-dwarf		Takata <i>et al.</i> (2005) <sup>(66)</sup>
216	Satsuma-yuki-mochi	Mine-no-yuki-mochi*	KG Mochi 102	*A descendant (the 3rd generation) of cv. Reimei through cv. Hayahikari (the Prefectural Institute for Agricultural Development 1st generation)	Kagoshima Prefectural Institute for Agricultural Development	2009	2009	18117	Semi-dwarf		Wakamatsu <i>et al.</i> (2007) <sup>(200)</sup>
217	Mine-no-yuki-mochi	Ouu 302 gou*	Himeno-mochi	*A descendant (the 2nd generation) of cv. Reimei	Hokuriku NAES	1991	1995	4231	Semi-dwarf		Shimizu <i>et al.</i> (1993) <sup>(51)</sup>
218	Massigura	Ouu 341 gou	Yamagata 40 gou*	*A descendant (the 4th generation) of cv. Reimei through cv. Akihikari	Aomori Prefectural Industrial Technology Research Center	2009	2009	18348	Semi-dwarf		
219	New Hikari	Etsunan 148 gou	Milky Queen (Kanto No. 168)*	*cf. cv. "Milky Queen" (Appendix I)	Fuku PAES	2009	2009	18349	Low amylose content		Tomita <i>et al.</i> (2007) <sup>(81)</sup>
220	Hoshimaru	Kami-iku 428 gou	Kuiku 159 gou*	*A descendant (the 4th generation) of cv. Hayamasari through cv. Ishikari	Hokkaido Prefectural Kamikawa Agricultural Experiment Station	2009	2009	18350			
221	Moe-minori	Nankai 128 gou	Haenuki*	*A descendant (the 4th generation) of cv. Reimei through "Shonai 29 gou"	NAKO Tohoku ARC	2009	2009	18351	Semi-dwarf		Kataoka <i>et al.</i> (2007) <sup>(62)</sup>
222	Tachi-aoba	Ha-kei 906*	An original line**	*A descendant (the 3rd generation) of cv. "Mine-asahi" through cv. Maturibare; **A descendant (the 3rd generation) of cv. Kinuhikari	NAKO Kyushu Okinawa ARC	2009	2009	18352			Sakai <i>et al.</i> (2009) <sup>(39)</sup>
223	Yuyake-mochi	Taisuko-mochi*	Bemigoromo**	*A descendant (the 2nd generation) of cv. Reimei through cv. Akihikari; **A descendant of cv. Reimei through cv. Fukuhibiki (the 4th generation)	NAKO Tohoku ARC	2009	2009	18353	Semi-dwarf	3356	Kataoka <i>et al.</i> (2007) <sup>(61)</sup>
224	Kareimai	Miyan 23 gou	Akihikari*	*A descendant (the 1st generation) of cv. Reimei	NAHO Hokuriku ARC	2009	2009	18473	Semi-dwarf		Shigemune <i>et al.</i> (2011) <sup>(50)</sup>
225	Koyuki-mochi	Waiboushi	Yamagata mochi 55 gou*	*A descendant (the 4th generation) of cv. Reimei through cv. "Kaguya-mochi"	Yamagata PAES	2009	2009	18546	Semi-dwarf		Chuba <i>et al.</i> (2007) <sup>(9)</sup>
226	Sue-akari	Aoi-no-kaze	Milky Queen (Kanto No. 168)*	*cf. cv. "Milky Queen" (Appendix I)	Mr. Jiro OZEKI	2008	2008	16008	Low amylose content		
227	Ayanatsuki	Aya*	An original line	*cf. "Aya" (NM391/Ishikari); (the 2nd generation of a gamma ray induced endosperm mutant "NM391"	Kagoshima PAES	2008	2008	16010	Low amylose content		Wakamatsu <i>et al.</i> (2005) <sup>(99)</sup>

No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
228	Mizuhonoka	LGC1*	Hyogo-kitanishiki**	*cf. cv. "LGC 1" (Appendix 1); **A descendant (the 4th generation) of cv. Reimei through cv. Nadahikari	NARO Kinokuniya Chugoku Shikoku ARC	2010	19409	Low gluten content in the grain; semi-dwarf		Iida <i>et al.</i> (2009) <sup>(40)</sup>
229	Manpuku-mochi	Etsunan 144 gou	Fukuhibiki*	*A descendant (the 4th generation) of cv. Reimei	Fukui PAES	2010	19053	Semi-dwarf		Tomita <i>et al.</i> (2008) <sup>(79)</sup>
230	Yukimusubi	Hatajirushi*	Higashi 810**	*A descendant (the 3rd generation) of cv. "Mine-asahi"; **A descendant (the 2nd generation) of cv. Reimei	Miyagi Prefectural Furukawa Agricultural Experiment Station	2010	19502	Tolerance to cold; semi-dwarf		Nagano <i>et al.</i> (2008) <sup>(69)</sup>
231	Koujyu-muryou	Yume-gokochi*	Yume-kahori**	*cf. cv. "Yume-gokochi" (Appendix 1); **cf. cv. "Yume-kahori" (Appendix 1)	Kaiyama Kometen	2010	18775			
232	Seto-no-niji	Hinohikari	Matsuribare*	*A descendant (the 1st generation) of cv. "Mine-asahi"	Yamaguchi Technology Center for Agriculture and Forestry	2010	19691			Hajima <i>et al.</i> (2010) <sup>(6)</sup>
233	Aki-matsuri	Yamahikari	Matsuribare*	*A descendant (the 1st generation) of cv. "Mine-asahi"	Yamaguchi Technology Center for Agriculture and Forestry	2010	19692			Hajima <i>et al.</i> (2012) <sup>(5)</sup>
234	Yume-sorata	Koshihikari	Akihikari*	*A descendant (the 1st generation) of cv. Reimei	Tottori PAES	2010	18780	Semi-dwarf		
235	Tenryu-otome	Inahikari*	Matsuribare**	**A descendant (the 1st generation) of cv. "Mine-asahi"	Nagano PAES	2010	19267			
236	Takane-murasaki	Kaguya-mochi*	Tohoku mochi 149 gou	*A descendant (the 4th generation) of cv. Reimei	Nagano PAES	2010	19045	Semi-dwarf		
237	Kinuhikari-Saitama SBL	Kinuhikari*	Sai-no-kagayaki**	*cf. cv. Kinuhikari; **A descendant of cv. "Mine-asahi"	Saitama PAES	2010	19049	Good taste; lodging resistance		Arakawa <i>et al.</i> (2008) <sup>(6)</sup>
238	Mine-no-murasaki	An original line	Asamurasaki*	*A descendant (the 5th generation) of cv. Reimei through cv. Fukuhibiki	Aichi PAES	3010	19055	Semi-dwarf		Saka <i>et al.</i> (2007) <sup>(37)</sup>
239	Beko-gonomi	An original line*	Fukuhibiki**	*A descendant (the 4th generation) of cv. Reimei; **A descendant (the 4th generation) of cv. Reimei	NARO NARC Tohoku Reg.	2010	19355	Semi-dwarf		Nakagami <i>et al.</i> (2008) <sup>(17)</sup>
240	Yume-matsuri	Asahi-no-yume*	Daichi-no-kaze**	**A descendant of cv. "Mine-asahi"	Aichi PAES	2010	19357			Kato <i>et al.</i> (2008) <sup>(65)</sup>
241	Ishikawa Sake 52 gou	An original line	Ippon-jime*	*A descendant (the 2nd generation) of cv. Reimei through cv. Houhai	Ishikawa PAES	2010	19695	Semi-dwarf		
242	Tomi-no-kaori	Yamadani-shiki	Oyamanishiki*	*A descendant (the 2nd generation) of cv. "Miyama-nishiki", that was induced through gamma ray irradiation to cv. Takanishiki (Appendix 1)	Toyama PAES	2010	19696			Ebitani <i>et al.</i> (2009) <sup>(60)</sup>
243	Mimi-hikari	Hikari-shinseiki*	An original line	*A descendant (the 3rd generation) of "Kanto 79 gou", that was induced through gamma ray irradiation (200 Gy) to the seed of cv. Koshihikari	Mr. Motomori TOMITA	2010	19985			

No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
244	Sai-no-minori	Yumematsuri (Aichi 108 gou)* Sai-no-kagayaki**	*A descendant of (G4 and G5, respectively) a gamma ray induced line "Kanto 79 gou" through cv. "Mine-asahi" and cv. Matsuribare; **A descendant (the 3rd generation) of "Kanto 79 gou" through cv. Matsuribare, a descendant (the 4th generation) of cv. Reimei	Saitama Pref. Agr. Forest. Res. Ctr.	2010	20027	Semi-dwarf		Ooka <i>et al.</i> (2009) <sup>(36)</sup>	
245	Koshi-no-kaori	Kinuhikari*	An indica-japonica hybrid back-crossed to Kinuhikari**	* **cf. cv. Kinuhikari	NARO	2011	20349	Good taste; lodging resistance		Sasahara <i>et al.</i> (2013) <sup>(42)</sup>
246	Shikibu-mochi	Asamurasaki (Ouu mochi 349 gou)*	Fu-kei mochi 170 gou	*A descendant (the 4th generation) of cv. Reimei through cv. Fukuhibiki	Aomori Prefectural Industrial Technology Research Center	2010	20030	Semi-dwarf		
247	Murasaki-no-kimi	Asamurasaki (Ouu mochi 349 gou)*	Fu-kei 176 gou	*A descendant (the 4th generation) of cv. Reimei through cv. Fukuhibiki	Aomori Prefectural Industrial Technology Research Center	2010	20031	Semi-dwarf		
248	Ake-no-murasaki	Asamurasaki (Ouu mochi 349 gou)*	Silky Pearl (Ouu 354 gou)**	*A descendant (the 4th generation) of cv. Reimei through cv. Fukuhibiki; **cf. cv. "Silky Pearl"	Iwate ARC	2011	20710	Semi-dwarf		Takakusagi <i>et al.</i> (2008) <sup>(62)</sup>
249	Gin-sayaka	Syun-yo (Hokuriku 183 gou)*	Gin-ginga (Iwate minami sake 13 gou)**	*A descendant (the 1st generation) of cv. "LGC1" (Appendix 1); **A descendant of cv. Reimei (Appendix 1) and cv. "Miyama-nishiki"	Iwate ARC	2011	20711	Semi-dwarf		
250	Aki-sakari	Awaminori (Hokuriku 159 gou)*	Etsunan 173 gou**	*A descendant (the 1st generation) of cv. Kinuhikari; **A descendant (the 1st generation) of both cv. Kinuhikari and a descendant (the 3rd generation) of cv. Reimei	Fukui PAES	2011	20431	Semi-dwarf		Tomita <i>et al.</i> (2009) <sup>(80)</sup>
251	Kita-aoba	Yume-aoba (Hokuriku 187 gou)*	cv. "Hatsushizuku" and cv. "Nanatsuboshi"	*A descendant (the 4th generation) of cv. Reimei through cv. Fukuhibiki	NARO	2011	20615	Semi-dwarf		
252	Ushi-yutaka	Yume-akari (Ao-kei 125 gou)*	Kanto PL 12 (Tsu-kei 995 gou)**	*A descendant of cv. Reimei; **A descendant of cv. Reimei through cv. "Musashi-kogane"	Aomori Prefectural Industrial Technology Research Center	2011	20530	Semi-dwarf		Maeda <i>et al.</i> (2008) <sup>(87)</sup>
253	Murasaki-koboshi	Kanto 195 gou	Asamurasaki*	*A descendant (the 4th generation) of cv. Reimei through cv. Fukuhibiki	NARO NARC Tohoku Reg.	2011	20714	Semi-dwarf		Kataoka <i>et al.</i> (2010) <sup>(63)</sup>
254	Nebari-yuki	Takitate (Tohoku 172 gou)*	Ao-kei 134 gou**	*A descendant (the 2nd generation) of cv. Reimei through "Ouu 343 gou"; **A descendant (the 3rd generation) of cv. Reimei through cv. Akihikari	Aomori Prefectural Industrial Technology Research Center	2011	20533	Semi-dwarf		Kon <i>et al.</i> (2008) <sup>(80)</sup>

No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
255	Saga-biyori	Tenshi-no-uta*	Aichi-no-kaori SBL**	*A descendant of both cv. "Mine-asahi" (the 3rd generation) and cv. Kinuhikari (the 2nd generation); **A descendant (the 3rd generation) of cv. "Mine-asahi"	Saga PARC	2011	20775			Hirota <i>et al.</i> (2012) <sup>(26)</sup>
256	Genki-tsukushi	Tsukushi-roman (Chikushi 46 gou)*	Tsukushi wase**	*A descendant (the 2nd generation) of Fukuoka PAES cv. Kinuhikari; **A descendant (the 2nd generation) of cv. Kinuhikari	Fukuoka PAES	2011	20744	Good taste; lodging resistance		Wada <i>et al.</i> (2010) <sup>(28)</sup>
257	Tsubu-yutaka	An original line	Fukuhibiki*	*A descendant (the 4th generation) of cv. Reimei	Iwate ARC	2011	20716	Semi-dwarf		Abe <i>et al.</i> (2009) <sup>(1)</sup>
258	Akidawara	Mirenishiki (Kanto 188 gou)	Ikuhikari (Etsunan 176 gou)*	*A descendant (the 2nd generation) of NARO National Institute of Crop Science cv. Kinuhikari through cv. Dontokoi	NARO National Institute of Crop Science	2011	20717	Good taste; lodging resistance		Ando <i>et al.</i> (2011) <sup>(5)</sup>
259	Milky Star	Tohoku 168 gou*	Milky Princess (Tohoku 194 gou)**	*A descendant (the 4th generation) of cv. Reimei through cv. Akihikari; **A descendant (the 1st generation) of cv. "Milky Queen" (Appendix 1)	NARO National Institute of Crop Science	2011	20718	Low amylose content		Ishii <i>et al.</i> (2012) <sup>(44)</sup>
260	Milky Summer	Wa-kei 243	Milky Queen (Kanto No. 168)*	*cf. cv. "Milky Queen" (Appendix 1)	NARO National Institute of Crop Science	2011	20745	Low amylose content		
261	Kinefuri-mochi	Mine-no-yuki-mochi*	Kokonoe-mochi	*A descendant (the 3rd generation) of cv. Reimei through cv. Hayahikari	Gifu PAES	2011	20945	Semi-dwarf		Hirose <i>et al.</i> (2010) <sup>(27)</sup>
262	Mina-yutaka	Fuyu-geshiki*	Fu-kei 186 gou**	*A descendant (the 4th generation) of cv. Reimei through cv. Akihikari; **A descendant (the 3rd generation) of cv. Reimei through cv. Akihikari	Aomori Prefectural Industrial Technology Research Center	2011	20913	Semi-dwarf		Ono <i>et al.</i> (2009) <sup>(35)</sup>
263	Maki-mizuhō	Hoshi-aoba*	Kusa-yutaka**	*A descendant (the 1st generation) of cv. Oochikara; **A descendant of cv. Oochikara	NARO Kyushu Okinawa ARC and JIRCAS	2011	21175	Semi-dwarf		
264	Mogu-mogu-aoba	Hoshi-aoba*	Mizuho-chikara**	*A descendant (the 1st generation) of cv. Oochikara; **A descendant (the 3rd generation) of cv. Reimei through cv. Akihikari	NARO Kyushu Okinawa ARC and JIRCAS	2011	20915	Semi-dwarf		
265	Hime-gonomi	Milky Queen*	Chugoku 169 gou	*cf. cv. "Milky Queen" (an NMU induced mutant cultivar) (Appendix 1)	NARO NARC for Western Region and JIRCAS	2012	21717	Low amylose content		Iida <i>et al.</i> (2011) <sup>(39)</sup>
266	Yamadawara	Izumi 348*	Kanto 192 gou**	*A descendant (the 5th generation) of cv. Reimei through cv. Akihikari; **A descendant (the 3rd generation) of cv. Kinuhikari through cv. Dontokoi	NARO National Institute of Crop Science and JIRCAS	2014	23197	Semi-dwarf		
267	Kita-aoba	Yume-aoba (Hokuriku 187 gou)*	Nanatsuboshi (Kuiku 163 gou)	*A descendant (the 5th generation) of cv. Reimei through cv. Fukuhibiki	NARO Hokuriku ARC	2011	20615	Semi-dwarf		
268	Tsukuba SD2 gou	An original line	Milky Queen*	*cf. cv. "Milky Queen"; an NMU induced mutant cultivar (Appendix 1)	Plant Genome Center Co. Ltd.	2012	21431	Low amylose content		
269	Torihime	Kan-no-mai*	Gyokuei	*A descendant (the 1st generation) of cv. "Miyama-nishiki"	Tottori PAES	2012	21383			

No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. Year	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
270	Yuki-asobi	An original line	Yume-akari*	*A descendant (the 5th generation) of cv. Reimei	Aomori Prefectural Industrial Technology Research Center	2012	2012	21433	Semi-dwarf		Kobayashi <i>et al.</i> (2010) <sup>7a)</sup>
271	Natsu-aoba	Yume-aoba (Hokuriku 187 gou)*	Akichikara**	*A descendant (the 5th generation) of cv. Reimei through cv. Fukuhibiki; **A descendant (the 2nd generation) of cv. Reimei	NARO Hokuriku ARC	2012	2012	21434	Semi-dwarf		Nagaoka <i>et al.</i> (2013) <sup>112)</sup>
272	Jyugemu	Yume-ikkon*	Yamadanishiki	*A descendant of both cv. Reimei and cv. Kinuhikari	Mr. Norimitsu KAWAMURA	2012	2012	22025	Semi-dwarf		
273	Niigata-jiro	Achikari*	Niigata 11 gou	*A descendant (the 1st generation) of cv. Reimei	Niigata PAES	2013	2013	22426	Semi-dwarf		Ishizaki <i>et al.</i> (2014) <sup>48)</sup>
274	Kin-no-megumi	Okimi-iri*	Asominori	*A descendant (the 4th generation) of cv. Reimei	NARO, Toyo Rice Co. Ltd.	2013	2013	22530	Semi-dwarf		Kaji <i>et al.</i> (2013) <sup>35)</sup>
275	Harumi	Koshihikari	Kinuhikari*	*cf. cv. Kinuhikari	Japan Agricultural Co-operatives (JA)	2014	2014	22985	Good taste; lodging resistance		
276	Rakufumai	Dontoki*	Gohyaku-mangoku	*A descendant (the 1st generation) of cv. Kinuhikari	NARO ARC	2014	2014	23198	Good taste; lodging resistance		
277	Tachi-ayaka	Hoshi-aoba*	An original line	*A descendant (the 2nd generation) of "BG1" through cv. Oochikara	NARO Kinki Chugoku Shikoku ARC	2014	2014	23275	Semi-dwarf		
278	Haigokoro	Milky Princess*	An original line	*A descendant (the 1st generation) of cv. "Milky Queen"	NARO Kinki Chugoku Shikoku ARC	2014	2014	23276	Low amylose content		Ishii <i>et al.</i> (2013) <sup>45)</sup>
279	Sato-no-shirayuki gou*	Hokuriku mochi 175 gou*	Aneko-mochi**	**A descendant (the 3rd generation) of cv. Reimei	NARO ARC	2014	2014	23429	Semi-dwarf		
280	Yume-no-mai	Koimusubi (Tohoku 160 gou)*	Syu 6084**	*A descendant (the 3rd generation) of "Kanto 79 gou" through cv. "Mine-asahi"; **A descendant (the 2nd generation) of cv. Kinuhikari	NARO Agricultural Research Center	2014	2014	23320	Good taste; lodging resistance		
281	Tochigi-no-hoshi	Tochigi 11 gou	Nasuhikari (Tochigi 7 gou)*	*A descendant of both "Kanto 79 gou" through cv. "Mine-asahi" (the 5th generation) and cv. Reimei (the 4th generation)	Tochigi PAES	2015	2015	24269	Semi-dwarf		Yamazaki <i>et al.</i> (2012) <sup>211)</sup>
282	Kazusa 1 gou	Koshihikari Kazusa 9 gou	Milky Queen (Kanto No. 168)*	*cf. cv. "Milky Queen" (Appendix 1)	Honda Motor Co., Ltd.	2015	2015	24336	Low amylose content		
283	Koshi-izumi	Hikari-shinseiki*	Yutaka-Koshihikari	*A descendant (the 5th generation) of "Kanto 59 gou"	Tottori University and Shizuoka University	2015	2015	23807			
284	Pikamaru	Kanto 221 gou*	Nikomaru (Saikai 250 gou)**	*A descendant of cv. "Milky Queen" through cv. "Milky Princess"; **A descendant of cv. Kinuhikari	NARO Kyushu Okinawa ARC	2015	2015	24270	Low amylose content; good taste; lodging resistance		



No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. Year	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
285	Konadamon	Hinohikari	Houiku 5 gou*	*A hybrid between an EI induced mutant cv. "LGC1" and a gamma ray induced mutant line "89WPKG30-433)	NARO Kyushu Okinawa ARC	2015	2015	24337	Low digestible protein content; low gluten content; low globulin content) and high non-digestible protein (prolamine) content		
286	Seto-no-kagayaki	Kimumusume (Saikai 232 gou)*	Yumematsuri (Aichi 108 gou)**	*cf. cv. Kinumusume; **cf. cv. Yumematsuri	NARO Kinki Chugoku Shikoku ARC	2015	2015	24361	Good taste with lodging resistance		
287	Emi-no-aki	Mine-haruka (Chubu 111 gou)*	Moeminori (Ouu 382 gou)**	*A descendant of "Kanto 79 gou" through cv. Mineasahi; **A descendant (the 5th generation) of cv. Reimei	NARO Tohoku ARC	2015	2015	24271	Semi-dwarf		
288	Yume-fuwari	Takitate (Tohoku 172 gou)*	LGC-Katsu**	*A descendant (the 2nd generation) of cv. Reimei; **A descendant (the 1st generation) of cv. "LGC1"	NARO Tohoku ARC and JIRCAS	2015	2015	24272	Semi-dwarf		
289	Tachi-hayate	An original line*	Hoshi-aoiba (Chugoku 146 gou)**	*A descendant (the 1st generation) of cv. "Odoroki-mochi" (Appendix 1); **A descendant (the 3rd generation) of Taihou through cv. Oochikara	NARO National Institute of Crop Science	2015	2015	24364			
290	Yamagata mochi 110 gou	Tatsuko-mochi (Akita mochi 45 gou)*	Oochikara**	*A descendant (the 2nd generation) of cv. Reimei through cv. Akihikari; **A descendant (the 2nd generation) of cv. Taihou	Yamagata PAES	2015	2015	24457	Semi-dwarf		
291	Kiraho	Milky Princess*	Iwate 61 gou	*A descendant (the 1st generation) of cv. "Milky Queen" (Appendix 1)	Iwate ARC	2015	2015	24582	Low amylose content		
292	Kin-no-ibuki	Takitate (Tohoku 172 gou)*	Mebae-mochi**	*A descendant (the 3rd generation) of cv. Reimei; **A descendant (the 2nd generation) of a NMU induced mutant line from cv. Kimmaze	Miyagi Prefectural Furukawa Agricultural Experiment Station	2015	2015	24378	Semi-dwarf		
293	Yamagata 95 gou	Yamagata 59 gou*	Chura-hikari (Ouu 366 gou)**	*A descendant of cv. Reimei; **A descendant (the 4th generation) of "Kanto 79 gou" through cv. "Mine-asahi"	Yamagata PAES	2015	2015	24454	Semi-dwarf		
294	Iwai-dawara	Ouu shi 394 gou	Beko-gonomi (Ouu shi 395 gou)*	*A hybrid between 2 descendants (both the 5th generation) of cv. Reimei	NARO National ARC for Tohoku Region	2015	2015	24363	Semi-dwarf		
295	Hana-sayaka	Kuro 1900*	Gin-ginga (Gaman sake 13 gou)**	*A descendant of cv. Reimei; **A descendant of both cv. Reimei and cv. "Miyama-nishiki"	Aomori Prefectural Industrial Technology Research Center	2015	2015	24456	Semi-dwarf		
296	Mine-no-hoshi	Akita-komachi*4/Mine-asahiRF*	Akita-komachi*4/(Chubu 111)	**cf. cv. "Mine-asahi"	Aichi PAES	2015	2015	24365			
297	Ginsan	Iwate 75 gou	Akita 63 gou*	*A descendant (the 3rd generation) of cv. Taihou through cv. Oochikara	Akita PAES	2015	2015	28538			

No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. Year	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
298	Mineharuka	Chubu 100 gou*	cv. "Chiyo-nishiki" X "Kumo 14252"	*A descendant (the 1st generation) of Aichi PAES cv. "Aichi-no-kaori"	Aichi PAES	2010	19054				
<b>Wheat</b>											
<i>Triticum aestivum</i> L.											
1	Akebono-mochi	Kinu-iroha	Tani-kei A6099*	*A mutant from EMS treatment to "Kanto No. 107"	NARC	2000	2000	8363	Extremely low amylose content	3129	Yamaguchi <i>et al.</i> (2003) <sup>215</sup>
2	Ibuki-mochi	Kinu-iroha	Tani-kei A6099*	*A mutant from EMS treatment to "Kanto No. 107"	NARC	2000	2000	8364	Extremely low amylose content	3130	Yamaguchi <i>et al.</i> (2003) <sup>202</sup>
3	Tama-izumi	Kan-kei w364	Kan-kei w361*	*A descendant (the 2nd generation) of gamma ray induced mutant cv. "Shiro-wase-komugi"	NARO (National Institute of Crop Science)	2002	2005	12962		3131	Fujita <i>et al.</i> (2004) <sup>15</sup>
4	Urara-mochi	Bandou-wase	Akebono-mochi *	*A descendant (the 1st generation) of EMS induced mutant line "Tani-kei A6099" from "Kanto 107 gou"	NARO (National Institute of Crop Science)	2009	2009	18434			Fujita <i>et al.</i> (2007) <sup>12</sup>
5	Yume-shiho	Tamaizumi (Kanto w421)*	An original line	*A descendant (the 3rd generation) of gamma ray induced mutant cv. "Shiro-wase-komugi"	NARO (National Institute of Crop Science)	2010	2010	19418			Kiribuchi-Otobe <i>et al.</i> (2009) <sup>69</sup>
6	Yume-akari	Haru-ibuki	Tamaizumi*	*A descendant (the 3rd generation) of gamma ray induced mutant cv. "Shiro-wase-komugi"	Aichi PAES	2014	2014	23409			
<b>Barley</b>											
<i>Hordeum vulgare</i> L.											
1	Nirasaki Nijo 8	Gamma No. 4*	34r127	*Gamma ray irradiation to "Kinn-cyoku No. 1"	Barley Research Center, Kirin Brewery Co., Ltd.	1967		1119			
2	Kawamizuki	U-kei H-83*	U-kei H-79**	*Gamma ray irradiation to cv. "Kanto-nakate-gold", **Gamma ray irradiation to "Asahi No. 5"	Kanto-Kyushu NAES	1979		1118	Early maturing; short and tough culm; constantly high-yielding		Tsuru <i>et al.</i> (1981) <sup>184</sup>
3	Tone-nijo	M4-66*	Nitta-kei 1	*Gamma ray irradiation to "Nitta-nijo No. 1"	Sapporo Breweries, Ltd.	1989	1989	1972		1164	
4	Masakado-mugi	Ea52*	Kanto-kawa No. 53	*Gamma ray irradiation to the plant of "Takebayashi-Ibaraki No. 1"	NARC	1989	1991	2688	Early maturing; resistant to stripe virus disease	1154	Makino <i>et al.</i> (1992) <sup>91</sup>
5	Komaki-nijo	(Nitta-kei 1 × Tochi-kei 85)F2	M4-66*	*Gamma ray irradiation to "Nitta-nijo No. 1"	Sapporo Breweries, Ltd.	1996	1996	5066		3132	
6	Sayakaze	Kanto-kawa No. 70*	Kanto-kawa No. 68 (Suzukaze)	*"Ea52" × "Kanto-kawa No. 60"; "Ea52"; Gamma ray irradiation to the plant of "Takebayashi-Ibaraki No. 1"	NARO (National Institute of Crop Science)	2003	2003	14301	Stripe virus disease resistance	3133	Yoshioka <i>et al.</i> (2005) <sup>214</sup>
7	Sakitama nijo	Tone nijo (Nitta-kei 25)*	Yasu-kei 58	*A descendant (the 2nd generation) of gamma ray induced "M4-66" through cv. "Tone nijo"	Sapporo Breweries, Ltd.	2003	2003	11235			
8	Kashima-goal	Sayakaze (Kanto kawa 78 gou)*	Kanto hadaka 78 gou	*A descendant (the 2nd generation) of Gamma ray induced disease resistant mutant "Ea52"	NARO Kinki Chugoku Shikoku NARC	2012	2012	21880	Stripe virus disease resistance		

No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference	
<b>Soybean</b>											
		<i>Glycine max</i> (L.) Merr.									
1	Nanbu-shirome	10KR-3-7 (Raïden)*	Kirami-nagaha	*cf. cv. Raïden (Appendix 1)	Tohoku NAES	1977		Medium-maturing	1593	Matsumoto <i>et al.</i> (1979) <sup>(95)</sup>	
2	Tomo-yutaka	Tohoku No. 52*	Kari-kei No. 102	*Raïden/"Toiku No. 109"	Tohoku NAES	1990	2878		3134	Watanabe <i>et al.</i> (1990) <sup>(201)</sup>	
3	Ryuhou	Suzu-yutaka	Kari-kou 343*	*A descendant (the 2nd generation) of cv. Raïden	Tohoku NAES	1995	5859		3135	Nakamura <i>et al.</i> (1996) <sup>(113)</sup>	
4	Suzu-no-ne	Kari-kei No. 244*	Kari-kei No. 221**	*A descendant (the 2nd generation) of cv. Raïden; **Named as cv. Kosuzu; cf. cv. Kosuzu (Appendix 1)	Tohoku NAES	1995	6206		3136	Nakamura <i>et al.</i> (1996) <sup>(114)</sup>	
5	Eru-star (L-star)	Fuku-yutakar/Kyu-kou 506*	Mura-yutaka**	*Named as cv. Ichihime; cf. cv. Ichihime (Appendix 1); **cf. cv. "Mura-yutaka" (Appendix 1)	Kyushu NAES	2000	8646		3137	Takahashi <i>et al.</i> (2003) <sup>(65)</sup>	
6	Suzu-sayaka	Suzu-yutaka	Kyu-kou 365F2(γ)-M4*	*A mutant line induced through Gamma ray irradiation to a hybrid between "Kanto 102 gou" and "Kan-kei 1 gou (later named as cv. Yumeiyutaka)"	NARO (NARC for Tohoku Region)	2003	14042	Complete absence of lipoxygenases		Yumoto <i>et al.</i> (2006) <sup>(217)</sup>	
7	Suzu-kaori	Kari-kou 778F5*	Kosuzu**	*A progeny of cv. Kosuzu and "Kitsurin No. 13"; **cf. cv. Kosuzu	NARO (NARC for Tohoku Region)	1995	15131		3139	Kono <i>et al.</i> (2006) <sup>(81)</sup>	
8	Kanto 100 gou	Enrei	En6500*	*A mutant line induced through EMS treatment to cv. Enrei	NARO (NARC)	2002	13531	Supermodulating character	3412	Akao & Kouchi (1992) <sup>(3)</sup> ; Takahashi <i>et al.</i> (2003) <sup>(64)</sup>	
9	Olerich50	M23*	LOLL**	*A high oleic acid mutant line induced through X-ray irradiation to cv. Bay;** A hybrid of M5 x M24, both of which are low linoleic acid content mutant through X-ray irradiation to cv. Bay	Saga University	1996	2008	16459			
10	Kinu-sayaka	Karikei 508 gou*	Developed line	*A lipoxygenase free mutant induced through a gamma ray irradiation	NARO (NARC for Tohoku Region)	1995	2008	16460	All lipoxygenase free	Kato <i>et al.</i> (2007) <sup>(67)</sup>	
11	Kyo-shirotanba	Murasaki-zukin*	Tama-daikoku	*Gamma ray irradiated mutant cultivar; cf. cv. "Murasaki-zukin"	Kyoto Prefecture	2013	22491				
12	Satohonoka	Kanto 100 Gou*	Nanbu-shirome**	*EMS treated mutant; cf. "Kanto 100 gou"; **derived from cv. Raïden (Appendix 1)	Iwate University and Akita Prefectural University	2014	23737				
13	Takamaru	Fukubuki	Kanto 100 gou*	*EMS treated super-nodulating mutant; cf. "Kanto 100 gou"	Iwate University and Akita Prefectural University	2014	23738	Supermodulating character			
14	Murasaki-zukin 3 gou	Murasaki-zukin 2 gou	Murasaki-zukin*	*Gamma ray induced mutant cultivar	Kyoto Prefecture	2015	23850				
15	Kogane-sayaka	Eru-star*	Sachi-yutaka	*cf. cv. Eru-star	NARO (National Institute of Crop Science)	2016	24955				
<b>Rush</b>											
		<i>Juncus effusus</i> L. var. <i>decipiens</i>	Buchtenau								
1	Hinomidori	Shimomasuda-zairai	Seto-nami*	*cf. cv. "Seto-nami" (Appendix 1)	Kumamoto PARC	1998	2001	9034	Long and many stems	3141	Nakazawa <i>et al.</i> (1999) <sup>(119)</sup>

No.	Cultivar Name	Maternal Line	Paternal Line	Origin of Parental Line	Institution of Development	Dev. Year	Reg. No. <sup>a</sup>	Reg. No. <sup>a</sup>	Mutant Characteristics Transferred	IAEA ID <sup>b</sup>	Reference
2	Suzukaze	<b>Hinomidori*</b>	An original line	*cf. cv. Hinomidori	Kumamoto PARC	2015	23967				Fushimizu <i>et al.</i> (2015) <sup>(4)</sup>
<b>Tomato</b>											
<i>Solanum lycopersicum</i> L.											
1	Kyoryoku-oogata-reikou	IRB301-31* <b>Sekai-ichi</b>	(Anahu <sup>×</sup> Ichihara) <sup>×</sup> Sekai-ichi	*Gamma ray irradiation to a wild relative species	Musashi Breeding Station Co.	1984			Resistance to both TMV and crown and root rot	2222	
2	Kagyoku	<b>Noh PL No. 4*</b>	Noh PL No. 3	*Gamma ray irradiation to a wild relative species	National Institute of Vegetables	1984	1986	964	Resistance to crown and root rot	2223	Yamakawa <i>et al.</i> (1987) <sup>(205)</sup>
3	Ryugyoku	<b>Noh PL No. 5*</b>	<b>Noh PL No. 4**</b>	* **Gamma ray irradiation to a wild relative species	National Institute of Vegetables	1984	1986	965	Resistance to fusarium crown and root rot	2224	Yamakawa <i>et al.</i> (1987) <sup>(205)</sup>
<b>Eggplant</b>											
<i>Solanum melongena</i> L.											
1	Dajiro	<b>Mutant of Daitaro*</b>	Hiranasu	*Selected from anther culture	Kochi PARC	2000	2004	1185	Fruit shape; green skin color	3142	
<b>Snapdragon</b>											
<i>Antirrhinum</i> L.											
1	Fuji Sweet Pink 1	<b>A mutant of cv. Touen*</b>	cv. Yukihime	*Colchicine treatment	Nihon Nohyaku Co., Ltd.	1993	1998	6398	Doubled chromosome number	3143	
<b>Job's tear</b>											
<i>Coix ma-yuen</i> Roman.											
1	Hato-yutaka	<b>Tohoku No. 1*</b>	<b>Oou No. 4**</b>	*A progeny of a gamma-ray induced mutant line "TS-kei" from a foreign cultivar; **Gamma ray irradiation to "Okayama-zairai"	NARO(NARC for Tohoku Region)	2004					Kato <i>et al.</i> (2007) <sup>(66)</sup>
<b>Japanese lawn grass</b>											
<i>Zoysia japonica</i> Steud.											
1	FLATZ	<b>COPROS*</b>	An original line	*A descendant (the 1st generation) of a somaclonal mutant through x-ray irradiation to a meristem callus	Kaisui Chemical Industry Co. Ltd.	2006		13648			

Note: Paternal and maternal line written in red are mutant lines or cultivars

<sup>a</sup>Reg. No. is the registration number of The Plant Variety Protection System in Japan

<sup>b</sup>IAEA ID is the variety ID number of the Joint FAO/IAEA Mutant Variety Database (<https://mvd.iaea.org/#Home>)

**Abbreviations:**

ARC: Agricultural Research Center  
 ARI: Agricultural Research Institute  
 JIRCAS: Japan International Research Center for Agricultural Sciences  
 NAES: National Agricultural Experiment Station  
 NARC: National Agriculture Research Center  
 NARO: National Agriculture and Food Research Organization  
 PAES: Prefectural Agriculture Experiment Station  
 PARC: Prefectural Agriculture Research Center

**Literature cited**

- 1) Abe, A., H. Sugawara, M. Takakusagi, and T. Sasaki (2009) Breeding of a new rice cultivar "Tsubuyutaka" with a high yielding ability. *Tohoku Agric. Res.* 62: 7-8 (in Japanese).
- 2) Akama, Y., T. Morimoto, K. Tanabe, M. Inoue, A. Shumiya, K. Fujii, Y. Ito, Y. Asai, T. Koide, S. Kudo, T. Ito, and A. Inagaki (1985) A new rice variety "Ibukiwase". *Res. Bull. Aichi Agric. Res. Ctr.* 17: 38-49 (in Japanese with English summary).
- 3) Akao, S. and H. Kouchi (1992) A supermodulating mutant isolated from soybean cultivar Enrei. *Soil Sci. Plant Nutr.* 38: 183-187.
- 4) Ando, I., H. Araki, H. Shimizu, M. Kuroki, K. Miura, K. Nagano, and K. Konno (2007) "Oborozuki", a new rice cultivar with low amylose content and superior eating quality. *Res. Bull. Natl. Agric. Res. Cent. Hokkaido Reg.* 186: 31-46 (in Japanese with English summary).
- 5) Ando, I., H. Nemoto, H. Kato, H. Ohta, H. Hirabayashi, Y. Takeuchi, H. Sato, T. Ishii, H. Maeda, T. Imbe, M. Hirayama, O. Ideta, M. Sakai, K. Tamura, and N. Aoki (2011) A new rice cultivar "Akidawara" with high yield, good grain appearance, and good eating quality. *Breeding Research* 13: 35-41 (in Japanese).
- 6) Arakawa, M., T. Minoda, H. Ishii, N. Ooka, Y. Takei, Y. Hirano, T. Ueno, K. Saito, K. Tokura, O. Shigematsu, M. Kozasu, K. Sekiguchi, Y. Okada, K. Watanabe, M. Arai, N. Arai, and S. Noda (2008) A new stripe and panicle blast resistant isogenic line "Kinuhikari Saitama SBL". *Bull. Saitama Pref. Agric. Res. Cent.* 8: 33-42 (in Japanese with English summary).
- 7) Arakawa, M., Y. Takei, K. Tokura, K. Yagasaki, K. Kozasu, T. Minoda, H. Ishii, Y. Okada, T. Sekiguchi, N. Ooka, K. Watanabe, K. Otsuka, and N. Arai (2003) New rice varieties, "Sainokagayaki" and "Sainokiribiyaka", with multiple pest resistance. *Bull. Saitama Pref. Agric. Forest. Res. Cent.* 3: 23-41 (in Japanese with English summary).
- 8) Araki, H., K. Konno, K. Miura, K. Nagano, S. Saito, M. Kobayashi, M. Nishimura, and K. Kariya (2002) A new rice cultivar with low amylose content "Hanabusa". *Res. Bull. Natl. Agric. Res. Cent. Hokkaido Reg.* 174: 69-81 (in Japanese with English summary).
- 9) Chuba, M., K. Yuki, H. Sakurada, K. Sato, T. Sano, R. Chuba, N. Yokoo, T. Honma, K. Watanabe, M. Mitobe, S. Sato, H. Miyano, H. Sato, and H. Goto (2007) Breeding of a new glutinous rice cultivar "Koyukimochi" (Yamagatamochi 87). *Issu. Yamagata Pref. Agric. Res.* 39: 1-25 (in Japanese with English summary).
- 10) Ebitani, T., T. Takarada, M. Omoteno, Y. Kojima, Y. Kidani, N. Mukaino, T. Yamaguchi, M. Doi, M. Morikawa, T. Matsushima, H. Ozaki, and M. Funane (2009) Breeding of rice variety "Tomi-no-kaori" suitable for Japanese sake-brewing. *Bull. Toyama Pref. Agric. Res. Cent.* 1: 1-10 (in Japanese with English summary).
- 11) Fujii, H., K. Toriyama, H. Shinoda, R. Yamamoto, and K. Sekizawa (1981) Breeding and characteristics of a new rice cultivar "Fujihikari". *Bull. Chugoku Agric. Exp. Stn. (Series A)* 29: 1-18 (in Japanese with English summary).
- 12) Fujita, M., C. Kiribuchi-Otobe, H. Matsunaka, M. Seki, T. Yoshioka, H. Yoshida, T. Nagamine, and I. Yamaguchi (2007) A new waxy wheat cultivar, "Uramochi". *Bull. Natl. Inst. Crop Sci.* 8: 109-129 (in Japanese with English summary).
- 13) Fujita, M., C. Kiribuchi-Otobe, T. Yoshioka, H. Matsunaka, T. Yanagisawa, H. Yoshida, I. Yamaguchi, T. Ushiyama, T. Nagamine, H. Seko, Y. Amano, and S. Oda (2004) A new hard white wheat cultivar "Tamaizumi". *Bull. Natl. Inst. Crop Sci.* 5: 1-17 (in Japanese with English summary).
- 14) Fushimizu, K., Y. Koushi, Y. Takamori, Y. Nakashima, K. Imure, and K. Fuchikami (2015) Breeding and characteristics of a new rush cultivar "Suzukaze". *Bull. Kumamoto Pref. Agric. Res. Ctr.* 22: 17-28 (in Japanese with English summary).
- 15) Hajima, M., K. Inoue, T. Yoshinaga, T. Morioka, H. Murayama, D. Watanabe, K. Kaneko, and Y. Nakatani (2012) Breeding of a new complex disease resistant rice variety "Akimaturi" with excellent apparent grain quality and good eating quality. *Bull. Yamaguchi Agri. Forest. Tec. Cent.* 3: 16-24 (in Japanese with English abstract).
- 16) Hajima, M., K. Kaneko, K. Inoue, G. Yamae, T. Yoshinaga, Y. Nakatani, T. Morioka, H. Murayama, and D. Watanabe (2010) Study on the new rice cultivar "Setononiji". *Bull. Yamaguchi Agri. Forest. Tec. Cent.* 1: 46-55 (in Japanese with English abstract).
- 17) Hamachi, Y., S. Imabayashi, K. F. Oosato, Y. Kawamura, H. Nishiyama, and M. Yoshino (1998) A new rice cultivar "Hayatsukushi". *Bull. Fukuoka Agric. Res. Cent.* 17: 15-22 (in Japanese).
- 18) Hamachi, Y., S. Imabayashi, K. F. Oosato, H. Nishiyama, M. Yoshino, Y. Kawamura, and Y. Matsue (1998) A new rice cultivar "Tsukushiwase". *Bull. Fukuoka Agric. Res. Cent.* 17: 1-8 (in Japanese).
- 19) Hamachi, Y., K. F. Oosato, Y. Kawamura, S. Imabayashi, H. Nishiyama, T. Wada, M. Yoshino, and T. Yasunaga (2003) A new rice cultivar "Tsukushiroman". *Bull. Fukuoka Pref. Agric. Exp. Stn.* 22: 11-18 (in Japanese with English summary).
- 20) Hamachi, Y., K. F. Oosato, Y. Kawamura, T. Wada, M. Tsubone, S. Imabayashi, T. Yasunaga, H. Nishiyama, and M. Yoshino (2004) A new rice cultivar for sake brewery "Yumeikkon". *Bull. Fukuoka Pref. Agric. Exp. Stn.* 23: 26-31 (in Japanese with English abstract).
- 21) Higashi, T., S. Saito, R. Ikeda, Y. Sunohara, S. Matsumoto, M. Inoue, Z. Oyamada, M. Yamaguchi, H. Kowata, and M. Yokoo (1994) Breeding of a high yielding rice cultivar "Fukuhibiki". *Bull. Tohoku Natl. Agric. Exp. Stn.* 88: 15-38 (in Japanese with English summary).
- 22) Higashi, T., S. Saito, T. Takita, M. Yamaguchi, Y. Sunohara, N. Yokogami, R. Ikeda, Y. Tamura, Z. Oyamada, H. Kowata, M. Inoue, S. Matsumoto, and T. Kataoka (1998) Development of a new rice cultivar with low amylose contents "Snow pearl". *Tohoku Agric. Res.* 51: 1-2 (in Japanese).
- 23) Higashi, T., M. Yamaguchi, Y. Sunohara, Z. Oyamada, H. Kowata, Y. Tamura, N. Yokogami, S. Saito, R. Ikeda, M. Inoue, and S. Matsumoto (1997) Breeding of a new rice cultivar "Okimiri". *Bull. Tohoku Natl. Agric. Exp. Stn.* 92: 15-33 (in Japanese with English abstract).
- 24) Higashi, T., M. Yamaguchi, Z. Oyamada, Y. Sunohara, H. Kowata, Y. Tamura, N. Yokogami, T. Sasaki, S. Abe, K. Matsunaga, E. Okamoto, A. Kano, H. Ikehashi, and H. Araki (1997) Breeding of a purple grain glutinous rice cultivar "Asamurasaki". *Bull. Tohoku Natl. Agric. Exp. Stn.* 92: 1-13 (in Japanese with English abstract).
- 25) Hirano, T., S. Seno, and K. Okubo (2004) A new rice cultivar "Fukuronomoko". *Bull. Agric. Exp. Stn. Okayama Pref. General Agric. Cent.* 22: 1-11 (in Japanese).
- 26) Hirose, T., S. Watanabe, Y. Akama, K. Shindo, N. Horisue, K. Matsumoto, H. Uchiyama, and S. Matsumoto (1977) On the new rice variety "Hayahikari". *Bull. Tohoku Natl. Agric. Exp. Stn.* 55: 1-30 (in Japanese with English summary).
- 27) Hirose, T., Y. Yamamoto, Y. Kawai, K. Sugiura, Y. Kagami, K. Miyata, T. Kobayashi, and H. Hisada (2010) Breeding process and characteristics of a new rice cultivar "Kinofurimochi". *Bull. Gifu Pref. Res. Inst. Agric. Sci. in Hilly and Mountainous Areas* 6: 1-6 (in Japanese with English summary).
- 28) Hirota, Y., S. Matsuyuki, K. Matsumoto, H. Yokoh, S. Tokuda, and T. Kinoshita (2012) A new rice cultivar "Yumeshizuku". *Bull. Saga Pref. Agric. Res. Cent.* 31: 1-21 (in Japanese with English summary).
- 29) Hirota, Y., S. Tokuda, I. Taira, K. Yoshida, T. Kinoshita, and S. Matsuyuki (2012) A new rice cultivar "Sagabiyori". *Bull. Saga Pref. Agric. Res. Cent.* 37: 1-19 (in Japanese).
- 30) Horisue, N., A. Kobayashi, E. Shimura, T. Kudo, Z. Oyamada, N. Nakagawa, Y. Fujimura, T. Mikami, Y. Kawamura, H. Yokohama, H. Tatsuta, M. Suto, M. Takadate, and M. Tateyama (1993) A new rice cultivar "Yukimimochi". *Bull. Aomori Agric. Exp. Stn.* 33: 69-84 (in Japanese with English summary).

- 31) Horisue, N., A. Kobayashi, E. Shimura, T. Kudo, Z. Oyamada, N. Nakagawa, Y. Fujimura, T. Mikami, M. Takadate, Y. Kawamura, H. Tatsuta, and M. Suto (1992) A new rice cultivar "Kitaou". Bull. Aomori Agric. Exp. Stn. 32: 1-16 (in Japanese with English summary).
- 32) Horisue, N., A. Kobayashi, E. Shimura, T. Kudo, Z. Oyamada, N. Nakagawa, Y. Fujimura, T. Mikami, H. Yokohama, Y. Kawamura, H. Tatsuta, M. Suto, M. Takadate, and M. Tateyama (1993) A new rice cultivar "Anekomochi". Bull. Aomori Agric. Exp. Stn. 33: 69-84 (in Japanese with English summary).
- 33) Horisue, N., A. Kobayashi, Y. Sunohara, Y. Uehara, E. Shimura, Y. Kawamura, H. Yokoyama, Y. Fujimura, T. Kudo, Z. Oyamada, T. Mikami, M. Suto, M. Suwa, M. Suto, H. Tatsuta, N. Nakagawa, K. Maeda, and M. Takadate (1997) A new rice cultivar "Ideyumochi". Bull. Aomori Agric. Exp. Stn. 36: 57-72 (in Japanese with English summary).
- 34) Horisue, N., A. Kobayashi, Y. Uehara, T. Kudo, Z. Oyamada, Y. Kawamura, Y. Fujimura, T. Mikami, H. Tatsuta, M. Suto, M. Tateyama, H. Yokoyama, and M. Suwa (1994) A new rice cultivar "Maihime". Bull. Aomori Agric. Exp. Stn. 34: 12-40 (in Japanese with English summary).
- 35) Horiuchi, H., K. Tomita, K. Terada, I. Tanaka, A. Kobayashi, T. Minobe, H. Furuta, A. Yamamoto, H. Shinoyama, I. Ikeda, M. Tanoi, K. Aoki, C. Kagoshima, N. Masaki, T. Minami, and A. Sugimoto (2004) A new rice cultivar "Sakuhikari". Bull. Fukui Pref. Agric. Exp. Stn. 41: 1-14 (in Japanese with English summary).
- 36) Hoshi, T., S. Abe, K. Ishizaki, S. Azuma, K. Kobayashi, K. Higuchi, T. Tamura, A. Harada, M. Ozeki, H. Kanayama, Y. Nagasawa, A. Takeuchi, O. Kawakami, Y. Sasaki, T. Hoshino, Y. Sasaki, M. Koide, and T. Kondo (1998) On the new rice cultivar "Hashirajiri". Bull. Niigata Pref. Agric. Exp. Stn. 43: 1-13 (in Japanese with English summary).
- 37) Hoshi, T., S. Abe, K. Ishizaki, H. Kasaneyama, K. Kobayashi, K. Hirao, T. Matsui, S. Azuma, K. Higuchi, T. Tamura, Y. Asai, K. Nakajima, A. Harada, M. Ozeki, Y. Sasaki, N. Abe, T. Kondo, and H. Kanayama (2002) A new rice cultivar "KOSHIBUKI". Journal of the Niigata Agric. Res. Inst. 5: 21-33 (in Japanese with English summary).
- 38) Ichikawa, Y., K. Yano, Z. Ike, and T. Hoshi (1981) On the new rice variety "NIGATAWASE". Bull. Niigata Pref. Agric. Exp. Stn. 30: 1-9 (in Japanese with English summary).
- 39) Iida, S., O. Iwata, K. Matsushima, Y. Sunohara, H. Nemoto, H. Maeda, T. Ishii, and Y. Tamura (2011) New low-amylose rice cultivar "Himegonomi" with good eating quality and good quality of appearance for the western region of Japan. Bull. Kinki Chugoku Shikoku Agric. Res. Cent. 10: 69-86 (in Japanese with English summary).
- 40) Iida, S., Y. Sunohara, O. Iwata, H. Maeda, H. Nemoto, T. Ishii, T. Yoshida, N. Nakagawa, M. Sakai, S. Furukawa, T. Mizuma, Y. Kiyokawa, and Y. Wakai (2009) New low glutelin brewers' rice cultivar "Mizuhonoka" for the western region. Bull. Kinki Chugoku Shikoku Agric. Res. Cent. 8: 45-62 (in Japanese with English summary).
- 41) Iida, S., Y. Sunohara, H. Maeda, K. Matsushita, H. Nemoto, T. Ishii, T. Yoshida, N. Nakagawa, M. Sakai, and T. Nishio (2004) A new rice cultivar with good eating quality (low amylose) and low glutelin protein, "LGC soft". Bull. Kinki Chugoku Shikoku Natl. Agric. Res. Cent. 3: 57-74 (in Japanese with English summary).
- 42) Imabayashi, S., Y. Hamachi, K. Furuno, H. Nishiyama, Y. Matsue, M. Yoshino, and T. Yoshida (1995) A new rice cultivar "Yumetsukushi". Bull. Fukuoka Agric. Res. Cent. 14: 1-10 (in Japanese with English abstract).
- 43) Inoue, M., Y. Nakajima, N. Saka, A. Syumiyama, S. Kudou, T. Touyama, K. Itou, M. Hayashi, T. Koide, K. Fujii, Y. Akama, T. Morimoto, and K. Tanabe (1993) A new blast-resistant rice variety "Inahikari". Res. Bull. Aichi Agric. Res. Ctr. 25: 1-11 (in Japanese with English summary).
- 44) Ishii, T., F. Ando, and H. Nemoto (2012) "Milky Star", a new rice cultivar with late-planting suitability and low amylose content in endosperm. Bull. Natl. Inst. Crop Sci. 13: 41-59 (in Japanese with English summary).
- 45) Ishii, T., O. Iwata, K. Matsushita, Y. Sunohara, H. Maeda, and S. Iida (2013) "Haigokoro", a new rice cultivar with high-emergence rate, low amylose content and giant embryo. Bull. NARO West. Reg. Agric. Res. Cent. 12: 25-41 (in Japanese with English summary).
- 46) Ishizaki, K. (2007) Studies on practical application of multiline cultivar, Koshihikari, with blast resistant genes in Niigata Prefecture. Journal of Niigata Agric. Res. Inst. 8: 1-37 (in Japanese with English summary).
- 47) Ishizaki, K., T. Matsui, S. Kaneda, K. Kobayashi, H. Kasaneyama, S. Abe, S. Azuma, T. Hoshi, Y. Sasaki, K. Higuchi, K. Hirao, N. Abe, T. Kondo, A. Takeuchi, and M. Koide (2008) A new cultivar "YUKINKOMAI". Journal of Niigata Agric. Res. Inst. 9: 89-98 (in Japanese with English summary).
- 48) Ishizaki, K., T. Matsui, K. Nabata, T. Kanbe, E. Nara, T. Hoshi, Z. Ike, I. Sasaki, T. Hoshino, A. Takeuchi, M. Koide, S. Azuma, T. Abe, and T. Kondou (2014) A new rice cultivar "Niigatajirou". Journal of Niigata Agric. Res. Inst. 12: 121-124 (in Japanese).
- 49) Iwashita, T., Y. Yamakawa, M. Machida, O. Doi, A. Shinya, M. Houman, and K. Moriya (1984) A new rice cultivar "Natsuhikari". Bull. Kagoshima Pref. Agric. Exp. Stn. 12: 1-11 (in Japanese with English summary).
- 50) Izawa, T., A. Shumiya, S. Kudo, T. Kato, K. Fujii, N. Saka, T. Touyama, N. Sugiura, T. Ito, H. Kojima, and Y. Nakajima (2001) A new rice variety "Asahiyume". Res. Bull. Aichi Agric. Res. Ctr. 33: 1-10 (in Japanese with English summary).
- 51) Izawa, T., A. Shumiya, S. Kudo, N. Saka, T. Kato, N. Sugiura, K. Fujii, T. Touyama, Y. Nakajima, T. Tsuji, H. Kojima, T. Ito, and Y. Hamada (2001) A new stripe and panicle blast resistance near isogenic line "Aichinokori SBL". Res. Bull. Aichi Pref. Agric. Res. Ctr. 33: 33-40 (in Japanese with English summary).
- 52) Izawa, T., A. Shumiya, S. Kudo, N. Saka, K. Fujii, T. Touyama, N. Sugiura, Y. Nakajima, T. Tsuji, and H. Kojima (2001) A new rice variety "Daichinokaze" with multiple pests resistance. Res. Bull. Aichi Agric. Res. Ctr. 33: 25-32 (in Japanese with English abstract).
- 53) Izawa, Y., M. Yuzawa, M. Fujii, K. Sotome, K. Oya, S. Kobayashi, T. Okubo, T. Kojima, M. Yamaguchi, H. Ito, K. Kurai, M. Deguchi, K. Tohigi, T. Sotome, and J. Ikeda (2005) A new paddy rice cultivar, "Nasuhikari". Bull. Tohigi Agric. Exp. Stn. 55: 1-14 (in Japanese with English summary).
- 54) Izumi, K., M. Mitsuoka, T. Syodai, T. Matsumoto, and H. Nizeki (1998) Breeding of a new rice cultivar "Yumeizumi". Bull. Kumamoto Pref. Agric. Res. Ctr. 7: 11-20 (in Japanese with English summary).
- 55) Kaji, R., H. Ohta, A. Fukushima, M. Yamaguchi, T. Kataoka, K. Nakagami, T. Endo, H. Kato, S. Ichiba, and K. Tsujiuchi (2013) A new rice variety for milled rice with scutellum "Kimmogumi", with good eating quality and resistance to blast and bacterial blight. Bull. Tohoku Agric. Res. Cent. 115: 1-10 (in Japanese with English summary).
- 56) Kaji, R., M. Okamoto, T. Yagi, H. Hirabayashi, R. Mizobuchi, S. Fukaura, K. Tamura, M. Nishimura, H. Yamashita, and T. Tomimatsu (2006) Fukuizumi, a new rice variety for direct seeding. Bull. Natl. Agric. Res. Cent. Kyushu Okinawa Reg. 47: 63-81 (in Japanese with English summary).
- 57) Kaji, R., M. Sakai, K. Tamura, H. Hirabayashi, M. Okamoto, T. Yagi, R. Mizobuchi, S. Fukaura, M. Nishimura, H. Yamashita, and T. Tomimatsu (2009) A new rice variety "Kinumusume". Bull. Natl. Agric. Res. Cent. Kyushu Okinawa Reg. 52: 79-94 (in Japanese with English summary).
- 58) Kamei, I., T. Suzuki, K. Sato, K. Oota, and S. Goto (1980) A new rice cultivar "Hamahikari". Bull. Yamagata Pref. Agric. Exp. Stn. 14: 19-33 (in Japanese with English summary).
- 59) Kaneda, H., Y. Umada, T. Ebitani, T. Matsushima, T. Ishibashi, Y. Kidani, Y. Yamamoto, M. Doi, and Y. Kozima (1999) New brewer's rice variety "Oyamamishiki". Hokuriku Crop Sci. 34: 4-7 (in Japanese).

- 60) Kaneeda, H., K. Yamamoto, T. Matsushima, T. Ishibashi, T. Ebitani, and Y. Yamamoto (1996) New rice variety "Owarabijin", Hokuriku Crop Sci. 31: 21-29 (in Japanese).
- 61) Kataoka, T., T. Endo, K. Nakagami, T. Takita, N. Yokogami, and H. Kato (2007) Breeding of a new red grain glutinous rice cultivar "Yuyakemochi". Bull. Natl. Agric. Res. Cent. Tohoku Reg. 107: 1-13 (in Japanese with English summary).
- 62) Kataoka, T., M. Yamaguchi, T. Endo, K. Nakagami, T. Takita, N. Yokogami, and H. Kato (2007) Development of a new rice variety "Moeminori" with high eating quality and adaptability to direct seeding. Bull. Natl. Agric. Res. Cent. Tohoku Reg. 107: 15-28 (in Japanese with English summary).
- 63) Kataoka, T., M. Yamaguchi, T. Endo, K. Nakagami, T. Takita, N. Yokogami, and H. Kato (2010) Development of a new purple and very small grain glutinous rice variety "Murasaki-koboshi". Bull. Natl. Agric. Res. Cent. Tohoku Reg. 111: 1-16 (in Japanese with English abstract).
- 64) Kato, H., R. Iba, S. Tsutsumi, H. Mitsuueda, T. Nakahara, H. Takeda, T. Takida, M. Kawaguchi, H. Arasuma, H. Yoshioka, Y. Nagayoshi, K. Wakasugi, H. Yamashita, and S. Ueda (2006) A new rice cultivar "Miyayutaka". Bull. Miyazaki Agric. Exp. Stn. 41: 61-84 (in Japanese with English summary).
- 65) Kato, M., M. Shirota, M. Nakamura, S. Kudo, K. Fujii, T. Tsuji, Y. Hamada, N. Saka, Y. Nakajima, T. Kato, T. Touyama, T. Funao, Y. Sawada, T. Izawa, T. Suzuki, I. Shaku, M. Inoue, A. Shumiyu, and H. Kojima (2008) Breeding of a new multiple resistant rice variety "Aichi 108" with excellent apparent grain quality and good eating quality. Res. Bull. Aichi Agric. Res. Ctr. 40: 83-91 (in Japanese with English Summary).
- 66) Kato, M., M. Yamamori, M. Yui, M. Ishida, I. Chiba, Y. Okuyama, T. Toyama, S. Tanosaki, T. Endo, and M. Shibata (2007) A new Job's tears cultivar "Hatoyutaka". Bull. Natl. Agric. Res. Cent. Tohoku Reg. 107: 43-51 (in Japanese with English summary).
- 67) Kato, S., S. Yumoto, Y. Takada, Y. Kono, S. Shimada, T. Sakai, H. Shimada, K. Takahashi, T. Adachi, K. Tabuchi, and A. Kikuchi (2007) A new soybean cultivar "Kinusayaka" lacking three lipoxigenase isozymes and Group A acetyl saponin. Bull. Natl. Agric. Res. Cent. Tohoku Reg. 107: 29-42 (in Japanese with English summary).
- 68) Kato, T., T. Hatakeyama, S. Masaki, S. Saito, K. Fukuda, M. Ono, K. Shimada, M. Taguchi, and T. Yamamoto (1995) Breeding of new rice cultivars "Kinunohada" and "Tatsukomochi". Bull. Akita Agric. Exp. Stn. 34: 23-48 (in Japanese with English summary).
- 69) Kiribuchi-Otobe, C., M. Seki, H. Matsumaka, T. Yoshida, M. Fujita, T. Yanagisawa, and H. Yoshida (2009) Breeding "Yumeshiho", a new bread wheat cultivar. Bull. Natl. Inst. Crop Sci. 10: 75-88 (in Japanese with English summary).
- 70) Kobayashi, A., N. Horisue, E. Shimura, T. Kudo, Z. Oyamada, N. Nakagawa, Y. Fujimura, T. Mikami, H. Yokoyama, Y. Kawamura, H. Tatsuta, M. Suto, M. Takadate, and M. Tateyama (1993) A new rice cultivar "Yamauta". Bull. Aomori Agric. Exp. Stn. 33: 101-117 (in Japanese with English summary).
- 71) Kobayashi, A., Y. Koga, H. Uchiyamada, S. Samoto, H. Horiuchi, K. Miura, K. Okuno, Y. Fujita, Y. Uehara, S. Ishizaka, M. Nakagahra, T. Yamada, and K. Maruyama (1990) Breeding a new rice variety "Oochikara". Bull. Hokuriku Agric. Exp. Stn. 32: 85-104 (in Japanese with English summary).
- 72) Kobayashi, A., E. Shimamura, N. Horisue, T. Kudo, Z. Oyamada, N. Nakagawa, T. Kanazawa, H. Yokoyama, Y. Fujimura, T. Mikami, Y. Kawamura, M. Suto, M. Takadate, and M. Tateyama (1993) A new rice cultivar "Yamauta". Bull. Aomori Agric. Exp. Stn. 33: 101-117 (in Japanese with English summary).
- 73) Kobayashi, K. (2008) Development of new rice cultivars with unique characteristics in Niigata Prefecture. Journal of Niigata Agric. Res. Inst. 9: 73-79 (in Japanese with English summary).
- 74) Kobayashi, W., M. Suwa, K. Maeda, S. Kanda, Y. Kawamura, and C. Kon (2010) Characteristics of a new ornamental rice variety with white leaf, "Yukiasobi". Tohoku Agric. Res. 63: 3-4 (in Japanese).
- 75) Kodama, I., T. Kawamoto, S. Matsumoto, K. Sato, M. Taguchi, K. Kyoya, T. Kato, T. Hatakeyama, and S. Masaki (2014) Breeding of a large grain rice cultivar exhibit high yields, "Akita63". Bull. Akita Agric. Exp. Stn. 54: 3-28 (in Japanese with English Abstract).
- 76) Kodama, I., S. Matsumoto, T. Kawamoto, S. Masaki, K. Sato, S. Shibata, R. Tamura, T. Kato, and T. Hatakeyama (2010) Breeding of a new rice cultivar, "Awayukikomachi", with a low amylose content. Bull. Akita Agric. Exp. Stn. 50: 3-20 (in Japanese with English abstract).
- 77) Koga, Y., H. Uchiyamada, S. Samoto, S. Ishizaka, Y. Fujita, K. Okuno, Y. Uehara, M. Nakagahra, H. Horiuchi, K. Miura, K. Maruyama, T. Yamada, T. Yagi, and K. Mori (1989) Breeding a new rice cultivar "Kinuhikari". Bull. Hokuriku Agric. Exp. Stn. 30: 1-24 (in Japanese with English summary).
- 78) Koga, Y., H. Uchiyamada, S. Samoto, S. Ishizaka, Y. Uehara, Y. Fujita, K. Okuno, M. Nakagahra, T. Yamada, K. Maruyama, T. Yagi, K. Mori, and K. Miura (1987) Breeding a new rice cultivar "Akichikara". Bull. Hokuriku Agric. Exp. Stn. 29: 23-45 (in Japanese with English summary).
- 79) Kojima, Y., T. Ebitani, H. Kameda, M. Doi, T. Ishibashi, Y. Kidani, M. Mukaino, T. Yamaguchi, M. Omoteno, Y. Yamamoto (2003) Development and utilization of isogenic lines Koshihikari Toyama. I. Development of isogenic lines Koshihikari Toyama BL. Bull. Toyama Agric. Res. Cent. 20: 13-32 (in Japanese with English summary).
- 80) Kon, C., Y. Kawamura, W. Kobayashi, K. Maeda, T. Mikami, H. Yokoyama, M. Takadate, M. Tateyama, and K. Kobayashi (2008) Development of a new rice variety "Nebaryuki" with low amylose content. Tohoku Agric. Res. 61: 1-2 (in Japanese).
- 81) Kono, Y., S. Yumoto, Y. Takada, S. Kato, S. Shimada, T. Sakai, H. Shimada, K. Takahashi, T. Adachi, K. Tabuchi, A. Kikuchi, S. Nakamura, M. Ito, and H. Banba (2006) A new small-seed soybean cultivar, "Suzukaori", with resistance to soybean mosaic virus. Bull. Natl. Agric. Res. Cent. Tohoku Reg. 105: 17-33 (in Japanese with English abstract).
- 82) Koomura, T., T. Ito, A. Shumiya, I. Shaku, M. Takamatsu, T. Kato, K. Fujii, N. Saka, and S. Kudo (1989) A new rice variety "Aichinokoari". Res. Bull. Aichi Agric. Res. Ctr. 19: 1-20 (in Japanese with English summary).
- 83) Koumura, T., M. Takamatsu, I. Shaku, A. Shumiya, T. Haga, M. Taniguchi, S. Kudo, K. Ito, K. Tanabe, Y. Nakazima, and Y. Kutsuna (1983) A new rice variety "Megumimochi". Res. Bull. Aichi Agric. Res. Cent. 15: 14-23 (in Japanese with English summary).
- 84) Kudo, S., T. Kato, I. Endo, K. Sugiura, M. Inoue, N. Saka, K. Ito, T. Otake, T. Touyama, Y. Nakajima, M. Hayashi, and Y. Akama (2000) A new non-glutinous rice variety "Minehikari". Res. Bull. Aichi Agric. Res. Ctr. 32: 1-9 (in Japanese with English abstract).
- 85) Kushibuchi, K. *et al.* (1977) A new rice cultivar "Akihikari". Bull. Aomori Agric. Exp. Stn. 22: 1-20 (in Japanese with English summary).
- 86) Kyoya, K., M. Taguchi, T. Kodama, T. Yamamoto, and Y. Sato (2002) New recommended rice cultivar "Haenuki". Bull. Akita Agric. Exp. Stn. 42: 1-20 (in Japanese with English summary).
- 87) Maeda, C., Y. Kawamura, W. Kobayashi, and C. Kon (2008) Characteristics of a new rice cultivar "Ushiyutaka" for whole-crop silage. Tohoku Journal of Crop Science 51: 21-22 (in Japanese).
- 88) Maeda, H., Y. Sunohara, S. Iida, K. Matsushita, H. Nemoto, T. Ishii, T. Yoshida, N. Kagawa, M. Sakai, T. Hoshino, M. Okamoto, and H. Shinoda (2002) A new rice cultivar for whole crop silage, "Hoshioaba". Bull. Natl. Agric. Res. Ctr. West Reg. 2: 83-98 (in Japanese with English summary).
- 89) Maeda, H., T. Tsuchiya, N. Hiraoka, M. Maeshige, S. Uemoto, and M. Nakayabu (1990) New cold resistant rice varieties "Hirohikari" and "Hirohonami". Bull. Hiroshima Pref. Agric. Exp. Stn. 53: 9-20 (in Japanese with English summary).

- 90) Maeda, M., O. Ito, M. Nakayabu, S. Nakazawa, H. Maeda, T. Hoshima, T. Tsuchiya, and Y. Doi (2000) A new paddy rice variety "Hiroshima 21" (Brand name: "Koimomiji"). Bull. Hiroshima Pref. Agric. Res. Cent. 68: 1-11 (in Japanese with English summary).
- 91) Makino, T., T. Fukuoka, A. Nakane, K. Fukunaga, K. Toya, M. Inagaki, H. Tozawa, H. Sasaki, S. Miyagawa, and Y. Ukai (1992) A new barley cultivar "Masakadomugi". Bull. Natl. Agric. Res. Cent. 24: 97-112 (in Japanese with English summary).
- 92) Masaki, S., T. Kato, T. Hatakeyama, S. Matsumoto, and T. Kawamoto (2004) Breeding of a new rice cultivar "Misatonishiki". Bull. Akita Agric. Exp. Stn. 44: 49-72 (in Japanese with English summary).
- 93) Masaki, S., T. Kato, T. Hatakeyama, S. Matsumoto, T. Kawamoto, T. Yamamoto, S. Dakeishi, S. Saito, K. Fukuda, K. Shimanuki, and N. Ikeda (1999) Breeding of a new rice cultivar "Akinosei". Bull. Akita Pref. Agric. Exp. Stn. 40: 23-43 (in Japanese with English summary).
- 94) Matsumoto, N., Y. Hashimoto, T. Hatakeyama, S. Tabuchi, M. Nishio, T. Nakata, K. Arakawa, and K. Nakamura (1993) On the new rice variety "Ougiwase". Bull. Ishikawa Pref. Agric. Exp. Stn. 17: 1-11 (in Japanese with English summary).
- 95) Matsumoto, S., K. Kokubun, S. Murakami, T. Nagasawa, T. Koyama, S. Nakamura, K. Hashimoto, M. Ishikawa, T. Miyahara, S. Matsumoto, and Z. Komno (1979) A new soybean variety "Nanbushirome". Bull. Tohoku Natl. Agric. Exp. Stn. 59: 87-104 (in Japanese with English summary).
- 96) Matsumoto, S., S. Masaki, T. Kawamoto, I. Kodama, T. Hatakeyama, and T. Kato (2006) Breeding of a new purple grain glutinous rice variety "Komurasaki". Bull. Akita Agric. Exp. Stn. 46: 21-40 (in Japanese with English summary).
- 97) Matsunaga, K., T. Sasaki, K. Nagano, E. Okamoto, S. Abe, K. Uematsu, A. Kano, H. Takizawa, H. Hayasaka, S. Usuki, and T. Kuroda (2002) New rice cultivar "Yumemusubi". Bull. Miyagi Pref. Furukawa Agric. Exp. Stn. 3: 37-51 (in Japanese with English summary).
- 98) Matsunaga, K., T. Sasaki, K. Nagano, E. Okamoto, S. Abe, K. Uematsu, A. Kano, H. Takizawa, H. Hayasaka, S. Usuki, T. Kuroda, and B. Chiba (2002) New rice cultivar "Hatajirushi". Bull. Miyagi Pref. Furukawa Agric. Exp. Stn. 3: 85-99 (in Japanese with English summary).
- 99) Mikami, T., M. Takadate, H. Yokoyama, W. Kobayashi, M. Tateyama, K. Maeda, Y. Kawamura, T. Nakahori, and Z. Oyamada (2004) A new variety "Fuyugeshiki". Bull. Aomori Agric. Forest Res. Ctr. 40: 23-37 (in Japanese with English summary).
- 100) Mikami, T., M. Takadate, W. Kobayashi, M. Tateyama, K. Maeda, H. Yokoyama, R. Kudo, T. Nakahori, Z. Oyamada, and T. Kudo (2000) A new rice cultivar "Yumeakari". Bull. Aomori Agric. Exp. Stn. 37: 31-47 (in Japanese with English summary).
- 101) Mikami, T., M. Takadate, H. Yokoyama, Y. Kawamura, W. Kobayashi, M. Tateyama, K. Maeda, R. Kudo, T. Nakahori, Z. Oyamada, and T. Kudo (2007) A new aromatic rice variety "Koihonoka". Bull. Aomori Agric. Forest Res. Ctr. 45: 45-62 (in Japanese with English summary).
- 102) Mikami, T., M. Takadate, H. Yokoyama, W. Kobayashi, M. Tateyama, K. Maeda, Y. Kawamura, H. Tatsuta, T. Nakahori, T. Kudo, and M. Namioka (2003) A new rice cultivar "Hanaomoi". Bull. Aomori Agric. Forest Res. Ctr. 39: 11-27 (in Japanese with English summary).
- 103) Mitsukawa, M., S. Hatakeyama, K. Kurata, M. Tanaka, S. Araki, T. Matsumoto, H. Syodai, S. Koga, and K. Izumi (2002) Breeding of a new rice cultivar with low amylose content "Akinero". Res. Bull. Kumamoto Pref. Agric. Res. Ctr. 11: 1-11 (in Japanese with English summary).
- 104) Miura, K., Y. Uehara, A. Kobayashi, H. Ohta, H. Shimizu, H. Sasahara, K. Fukui, Y. Komaki, A. Goto, A. Shigemune, and H. Otsuki (2007) A new variety "Ayunohikari". Bull. Natl. Agric. Res. Cent. 9: 1-16 (in Japanese with English summary).
- 105) Miura, K., Y. Uehara, A. Kobayashi, H. Ohta, H. Shimizu, H. Sasahara, K. Fukui, Y. Komaki, H. Otsuki, A. Goto, and A. Shigemune (2006) A new rice variety "Yumeaoba". Bull. Natl. Agric. Res. Cent. 7: 1-23 (in Japanese with English summary).
- 106) Miyata, Y., K. Koike, H. Yamamoto, Y. Kida, N. Kawada, T. Kameyama, and M. Sato (2001) Breeding of a new rice variety "Natsushizuka" and its characteristics. Bull. Shizuoka Agric. Exp. Stn. 46: 1-10 (in Japanese with English summary).
- 107) Mizobuchi, M., Y. Yamasaki, A. Iwasaki, Y. Nakamura, M. Kameshima, and H. Uehigashi (2003) A new rice variety for sake brewing, "Kazentaruko". Bull. Kochi Agric. Res. Cent. 12: 81-90 (in Japanese with English summary).
- 108) Morimoto, T., T. Ito, K. Tanabe, M. Nakamori, M. Taniguchi, A. Inagaki, M. Inoue, Y. Asai, Y. Ito, A. Shumiya, and K. Fujii (1980) A new rice variety "Mineasahi". Res. Bull. Aichi Agric. Res. Cent. 12: 24-36 (in Japanese with English summary).
- 109) Nagano, K., H. Hayasaka, B. Chiba, N. Miyano, K. Sasaki, T. Endo, and K. Wagatsuma (2008) A new rice cultivar "Yukimusubi". Bull. Miyagi Furukawa Agric. Exp. Stn. 7: 19-38 (in Japanese with English summary).
- 110) Nagano, K., K. Matsunaga, T. Sasaki, S. Abe, E. Okamoto, K. Uematsu, H. Takizawa, H. Hayasaka, S. Usuki, T. Kuroda, B. Chiba, and N. Miyano (2005) A new rice cultivar "Koimusubi". Bull. Miyagi Pref. Furukawa Agric. Exp. Stn. 5: 1-14 (in Japanese with English summary).
- 111) Nagano, K., K. Matsunaga, H. Takizawa, H. Hayasaka, S. Usuki, T. Kuroda, B. Chiba, N. Miyano, and K. Sasaki (2005) A new rice cultivar "Takitae". Bull. Miyagi Pref. Furukawa Agric. Exp. Stn. 5: 15-30 (in Japanese with English summary).
- 112) Nagao, I., K. Miura, Y. Uehara, H. Sasahara, H. Ohta, A. Goto, A. Shigemune, and Y. Komaki (2013) "Natsuoba", a new rice cultivar for whole crop silage. Breeding Research 15: 25-31 (in Japanese).
- 113) Nakamura, S., S. Yumoto, K. Takahashi, K. Tabuchi, T. Adachi, A. Kikuchi, M. Kowata, H. Banba, N. Takahashi, A. Okabe, I. Watanabe, T. Nagasawa, S. Murakami, K. Hashimoto, S. Sakai, and K. Igita (1996) A new soybean variety "Ryuhou". Bull. Tohoku Natl. Agric. Exp. Stn. 91: 1-11 (in Japanese with English abstract).
- 114) Nakamura, S., S. Yumoto, K. Takahashi, K. Tabuchi, T. Adachi, A. Kikuchi, M. Kowata, H. Banba, N. Takahashi, A. Okabe, I. Watanabe, T. Nagasawa, S. Murakami, K. Hashimoto, S. Sakai, and K. Igita (1996) A new soybean variety "Suzunone". Bull. Tohoku Natl. Agric. Exp. Stn. 91: 13-23 (in Japanese with English abstract).
- 115) Nakagawa, J., T. Yoshida, K. Teramoto, H. Noda, and S. Taniguchi (2005) A new rice cultivar "Lake 65". Bull. Shiga Agric. Res. Cent. Exp. Stn. 45: 1-12 (in Japanese with English summary).
- 116) Nakagomi, K., M. Yamaguchi, T. Kataoka, T. Endo, T. Takita, T. Higashi, N. Yokogami, H. Kato, and Y. Tamura (2006) Breeding of a new rice cultivar, "Bekooba", for whole-crop silage adapted to direct-seeding cultivation. Bull. Natl. Agric. Res. Cent. Tohoku Reg. 106: 1-14 (in Japanese with English summary).
- 117) Nakagomi, K., M. Yamaguchi, T. Kataoka, T. Endo, T. Takita, N. Yokogami, and H. Kato (2008) Development of a new early maturity rice cultivar, "Bekogonomi", for feed. Bull. Natl. Agric. Res. Cent. Tohoku Reg. 109: 1-13 (in Japanese with English summary).
- 118) Nakamura, Y., M. Kameshima, H. Uga, Y. Morita, and S. Nomura (1993) A new rice extremely early variety "Hatsunanori". Bull. Kochi Agric. Res. Ctr. 2: 45-54 (in Japanese with English summary).
- 119) Nakazawa, Y., T. Tetsuka, K. Iimure, Y. Yuno, and Y. Takamori (1999) Breeding and characteristics of a new rush cultivar "Hinomidori". Res. Bull. Kumamoto Pref. Agric. Res. Ctr. 8: 47-53 (in Japanese with English summary).
- 120) Narizuka, A., S. Orimo, T. Takahashi, M. Osawa, Y. Obuchi, and Y. Saito (1998) A new rice variety with low amylose, "Sawapikari". Bull. Gunma Agric. Exp. Stn. 5: 1-8 (in Japanese with English summary).



- 121) Nemoto, H., S. Iida, H. Maeda, T. Ishii, N. Nakagawa, T. Hoshino, M. Sakai, M. Okamoto, H. Shinoda, and T. Yoshida (2001) A new rice cultivar with giant embryo "Haiminori". Bull. Chugoku Natl. Agric. Exp. Stn. 22: 25-40 (in Japanese with English summary).
- 122) Nihara, S., Y. Tomihisa, S. Okatake, K. Ishida, T. Mizushima, and H. Kimoto (1996) A new variety "Otomegokoro". Bull. Okayama Agric. Exp. Stn. 14: 25-39 (in Japanese).
- 123) Nishikawa, Y., R. Hayashi, K. Wada, T. Nagashima, K. Saito, Y. Koyama, and T. Watanabe (2006) "Chiba 28" a new rice cultivar with medium maturity, high cold resistance, high field resistance to blast, good grain quality and excellent eating quality. Bull. Chiba Agric. Res. Ctr. 5: 75-85 (in Japanese with English summary).
- 124) Nishimura, M., M. Kusaba, K. Miyahara, T. Nishio, S. Iida, T. Imbe, and H. Sato (2005) New rice varieties with low levels of easy-to-digest protein, "LGC-Katsu" and "LGC-Jun". Breed. Sci. 55: 103-105.
- 125) Niwayama, T., K. Suzuki, K. Tokura, K. Morita, H. Shiohara, H. Hasegawa, M. Tamura, S. Ogawa, and N. Minegishi (1987) On the new rice variety Tama-minori. Bull. Saitama Agric. Exp. Stn. 42: 1-15 (in Japanese).
- 126) Niwayama, T., K. Tokura, K. Yagasaki, S. Kato, H. Shiohara, M. Tamura, K. Suzuki, H. Hasegawa, K. Inoue, K. Morita, N. Minegishi, and R. Tateoka (1994) A new rice variety "Yumeminori". Bull. Saitama Agric. Exp. Stn. 47: 1-24 (in Japanese with English summary).
- 127) Niwayama, T., K. Tokura, K. Yagasaki, S. Kato, M. Tamura, K. Suzuki, H. Hasegawa, H. Shiohara, and K. Inoue (1991) A new glutinous rice variety "Heiseimochi". Bull. Saitama Agric. Exp. Stn. 45: 1-17 (in Japanese with English summary).
- 128) Noda, H., K. Teramoto, Y. Kohara, S. Taniguchi, and G. Ueda (1999) A new rice cultivar "Yumeohmi". Bull. Shiga Agric. Exp. Stn. 40: 1-12 (in Japanese with English summary).
- 129) Odanaka, H., Y. Ogi, H. Sugawara, T. Sato, M. Takahashi, Y. Kiuchi, H. Nakano, and A. Nakamishi (2000) Breeding of a new rice variety "Ginginga" suitable for sake brewing. Res. Bull. Iwate Agric. Res. Ctr. 1: 39-50 (in Japanese with English summary).
- 130) Oikawa, T., T. Sasaki, K. Suzuki, S. Abe, K. Matsunaga, S. Wakui, and K. Tanno (1991) A new variety of aromatic rice "Miyakaori". Bull. Miyagi Pref. Furukawa Agric. Exp. Stn. 1: 50-62 (in Japanese with English summary).
- 131) Okamoto, M., H. Hirabayashi, R. Kaji, R. Fukuoka, T. Yagi, H. Nishiyama, M. Nishimura, S. Fukaura, H. Yamashita, T. Takita, and K. Saito (2001) A new rice cultivar "Yawarakomachi". Bull. Natl. Agric. Res. Cent. Kyushu Okinawa Reg. 39: 127-141 (in Japanese with English summary).
- 132) Okamoto, M., R. Kaji, K. Tamura, R. Mizobuchi, H. Hirabayashi, S. Fukaura, M. Nishimura, T. Yagi, H. Yamashita, and T. Tomimatsu (2008) New rice variety "Akiyayaka". Bull. Kyushu Okinawa Agric. Res. Ctr. 49: 33-50 (in Japanese with English summary).
- 133) Ono, K., K. Tanabu, K. Arima, T. Mikami, K. Yamazaki, T. Kanazawa, H. Mimoto, M. Namioka, T. Kudo, and M. Yoshihara (1982) A new rice cultivar "Mutsukomachi". Bull. Aomori Agric. Exp. Stn. 26: 61-79 (in Japanese with English summary).
- 134) Ono, K., K. Tanabu, K. Arima, T. Mikami, K. Yamazaki, T. Kanazawa, H. Mimoto, M. Namioka, T. Kudo, and M. Yoshihara (1982) A new rice cultivar "Mutsukaori". Bull. Aomori Agric. Exp. Stn. 26: 81-102 (in Japanese with English summary).
- 135) Ono, T., S. Kanda, S. Moriyama, and M. Suto (2009) Characteristics of a new rice cultivar "Minayutaka" with cool temperature tolerance and high yield. Tohoku Agric. Res. 62: 5-6 (in Japanese).
- 136) Ooka, N., M. Arakawa, T. Minoda, H. Ishii, Y. Takei, T. Ueno, K. Tokura, K. Saito, O. Shigematsu, M. Kozausu, Y. Okada, S. Noda, K. Watanabe, M. Arai, and N. Arai (2009) Breeding of a new variety, "Sainominori", with multiple pest resistance and high grain quality, excellent eating quality. Bull. Saitama Pref. Agric. Forest. Res. Ctr. 8: 43-54 (in Japanese).
- 137) Saka, N., T. Terashima, S. Kudo, T. Kato, K. Sugura, I. Endo, M. Shiota, M. Inoue, and T. Otake (2007) A new purple grain glutinous rice variety "Minomurasaki" for rice processing. Res. Bull. Aichi Agric. Res. Ctr. 39: 111-120 (in Japanese with English abstract).
- 138) Sakai, M., M. Okamoto, K. Tamura, R. Kaji, R. Mizobuchi, H. Hirabayashi, T. Yagi, M. Nishimura, and S. Fukaura (2006) A new rice variety "Akimasari". Bull. Kyushu Okinawa Agric. Res. Cent. 47: 43-62 (in Japanese with English summary).
- 139) Sakai, M., M. Okamoto, K. Tamura, R. Kaji, R. Mizobuchi, H. Hirabayashi, T. Yagi, M. Nishimura, and S. Fukaura (2009) A new rice variety for whole crop silage "Tachiaoba". Bull. Natl. Agric. Res. Cent. Kyushu Okinawa Reg. 50: 1-20 (in Japanese with English summary).
- 140) Sakai, M., M. Okamoto, K. Tamura, R. Kaji, R. Mizobuchi, H. Hirabayashi, T. Yagi, M. Nishimura, and S. Fukaura (2010) "Nikomaru", high-yielding rice variety with superior eating quality and grain appearance under high temperatures during ripening. Bull. Natl. Agric. Res. Cent. Kyushu Okinawa Reg. 54: 43-61 (in Japanese with English summary).
- 141) Samoto, S., S. Asakuma, K. Hamamura, I. Kato, S. Ishizaka, A. Kobayashi, D. Kanai, T. Yagi, C. Kaneta, H. Ikehashi, K. Maruyama, T. Yamada, Y. Uehara, and K. Mori (1979) On a new rice cultivar "Sachiminori". Bull. Hokuriku Natl. Agric. Exp. Stn. 22: 55-87 (in Japanese with English summary).
- 142) Sasahara, H., K. Miura, H. Shimizu, A. Goto, A. Shigemune, I. Nagaoka, Y. Uehara, A. Kobayashi, H. Ohta, K. Fukui, H. Ohtsuki, M. Yano, and Y. Komaki (2013) "Koshinokaori", a new rice noodle cultivar. Bull. NARO Agric. Res. Cent. 19: 15-29 (in Japanese with English summary).
- 143) Sasaki, T., S. Abe, K. Matsunaga, E. Okamoto, K. Nagano, K. Tanno, Y. Chiba, A. Kano, and K. Uematsu (1994) A new cultivar of aromatic rice "Haginokaori". Bull. Miyagi Pref. Furukawa Agric. Exp. Stn. 2: 18-30 (in Japanese with English summary).
- 144) Sasaki, Y., S. Azuma, T. Kondoh, K. Ishizaki, N. Abe, Y. Sasaki, T. Hoshi, A. Takeuchi, and T. Hoshino (1994) New rice variety "Ipponjime". Hokuriku Crop Science 29: 5-6 (in Japanese).
- 145) Sato, K., H. Sakurada, T. Sano, R. Chiba, N. Yokoo, K. Yuki, S. Sato, A. Yamakawa, H. Sato, and T. Honma (2000) Breeding process and characteristics of a new variety "Yumesayaka". Tohoku Agric. Res. 53: 5-6 (in Japanese with English summary).
- 146) Sato, H., T. Imbe, H. Nemoto, Y. Akama, N. Horisue, H. Ohta, H. Hirabayashi, O. Ikeda, I. Ando, M. Suto, K. Numaguchi, M. Takadate, H. Hirasawa, M. Sakai, K. Tamura, and N. Aoki (2008) "Milky Princess", a new rice cultivar with low-amylose content in endosperm. Bull. Natl. Inst. Crop Sci. 9: 63-79 (in Japanese with English summary).
- 147) Sato, H., S. Saito, M. Oowada, Y. Arai, T. Taira, H. Saito, S. Hanzawa, and Y. Kida (2006) Breeding of a new rice cultivar "Fukumirai". Bull. Fukushima Pref. Agric. Exp. Stn. 37: 40-53 (in Japanese with English abstract).
- 148) Sato, H., S. Saito, M. Oowada, Y. Arai, T. Taira, H. Saito, S. Hanzawa, and Y. Kida, T. Takeda, T. Sato, and M. Takahashi (2003) Breeding of a new rice cultivar "Yumenokaori". Bull. Fukushima Pref. Agric. Exp. Stn. 36: 49-63 (in Japanese with English abstract).
- 149) Sato, S., S. Gotoh, Y. Tanifuji, E. Kikuchi, H. Sakurada, M. Chuba, Y. Kambayashi, T. Uematsu, S. Ohba, E. Ooe, and K. Ohbuchi (1988) On the breeding of the new paddy rice variety "MICHINOKUWASE". Bull. Yamagata Agric. Exp. Stn. 23: 1-15 (in Japanese with English summary).
- 150) Shigemune, A., K. Miura, Y. Uehara, A. Kobayashi, Y. Koga, H. Uchiyamada, S. Samoto, H. Sasahara, A. Goto, H. Ohta, H. Shimizu, Y. Fujita, S. Ishizaka, M. Nakagahara, K. Okuno, T. Yamada, Y. Komaki, H. Horuchi, K. Fukui, H. Otsuki, and K. Maruyama (2011) A new rice variety "Kareimai". Bull. Natl. Agric. Res. Ctr. 16: 17-37 (in Japanese with English summary).

- 151) Shimizu, H., K. Miura, H. Ohta, A. Kobayashi, H. Horiuchi, Y. Koga, K. Okuno, Y. Fujita, and H. Uchiyamada (1993) Breeding process and characteristics of a new rice variety "Minenyokimochi". Jpn. J. Crop Sci. (Hokuriku Branch) 28: 4-6 (in Japanese).
- 152) Shimura, E., A. Kobayashi, H. Sato, N. Horisue, T. Kudo, N. Nakagawa, T. Kanazawa, Z. Oyamada, M. Takadate, H. Yokoyama, Y. Fujimura, T. Mikami, Y. Kawamura, M. Namioka, H. Tatsuta, and M. Suto (1990) A new rice cultivar "Koihime". Bull. Aomori Agric. Exp. Stn. 31: 111-127 (in Japanese with English summary).
- 153) Shimura, E., A. Kobayashi, H. Sato, T. Kudo, N. Nakagawa, T. Kanazawa, Z. Oyamada, M. Takadate, H. Yokoyama, Y. Fujimura, and M. Namioka (1986) A new rice cultivar "Hatsukogane". Bull. Aomori Agric. Exp. Stn. 29: 1-14 (in Japanese with English summary).
- 154) Shiobara, H., T. Niwayama, N. Kasahara, K. Morita, and S. Ogawa (1982) A new rice variety "Musashi-kogane". Bull. Saitama Agric. Exp. Stn. 38: 1-14 (in Japanese).
- 155) Suga, R., Y. Iida, T. Kirihara, K. Yokota, S. Nishimiya, Y. Okutsu, Y. Takagi, and H. Hanawa (2000) Breeding of a new rice variety "Yumehitachi". Bull. Ibaraki Plant Biotech. Inst. 3: 17-29 (in Japanese with English summary).
- 156) Sugawara, H., S. Kawadai, A. Abe, and T. Sasaki (2012) Breeding of a new rice cultivar "Iwate Sake No. 98" good for daiginjo sake. Tohoku Agric. Res. 65: 9-10 (in Japanese).
- 157) Sunohara, Y., M. Sakai, M. Suto, K. Maeda, S. Kanda, H. Yokoyama, Y. Kawamura, M. Tateyama, T. Yashima, and S. Saito (2007) A new rice cultivar "Yukinohana". Bull. Aomori Agric. Forest. Res. Cent. 41: 1-22 (in Japanese with English summary).
- 158) Sunohara, Y., M. Sakai, Y. Uehara, M. Suto, M. Tateyama, S. Kanda, S. Sakamoto, Y. Kawamura, H. Yokoyama, K. Maeda, M. Suta, and T. Yashima (2004) A new rice variety "Komanomai". Bull. Aomori Agric. Forest Res. Cent. 40: 1-21 (in Japanese with English summary).
- 159) Syunura, A., T. Ito, S. Kudo, K. Fujii, T. Kato, N. Saka, T. Touyama, and I. Shaku (1994) A new stripe-resistant rice variety "Matsuribare". Res. Bull. Aichi Agric. Res. Ctr. 26: 1-16 (in Japanese with English abstract).
- 160) Takadate, M., T. Mikami, H. Yokoyama, W. Kobayashi, H. Tatsuta, K. Maeda, R. Kudo, Y. Kawamura, H. Tugawa, M. Tateyama, T. Nakahori, M. Namioka, T. Kudo, and Z. Oyamada (1997) A new rice cultivar "Tsugaruroman". Bull. Aomori Agric. Exp. Stn. 36: 1-17 (in Japanese with English summary).
- 161) Takadate, M., K. Yamazaki, K. Tanabu, T. Mikami, K. Arima, H. Tatsuta, Y. Kawamura, H. Yokoyama, M. Namioka, and T. Kanazawa (1990) A new rice cultivar "Tsugaruotome". Bull. Aomori Agric. Exp. Stn. 31: 95-110 (in Japanese with English summary).
- 162) Takasugi, M., A. Abe, H. Nakano, and Y. Kiuchi (2008) Breeding of a new purple grain rice cultivar "Akenomurasaki". Tohoku Agric. Res. 61: 7-8 (in Japanese).
- 163) Takahashi, H., T. Taguchi, K. Ishikawa, T. Nakata, K. Saito, T. Samuta, K. Iwano, and T. Ishikawa (1999) Breeding process and brewing characteristics of a new rice cultivar, "Gin-no-sei" suitable for sake brewing. Bull. Akita Res. Inst. Food and Brewing 1: 1-7 (in Japanese).
- 164) Takahashi, M., J. Arihara, N. Nakayama, M. Kokubun, S. Shimada, K. Takahashi, and M. Hajika (2003) Breeding of supermodulating soybean cultivar "Sakukei 4". Bull. Natl. Inst. Crop Sci. 4: 17-28 (in Japanese with English summary).
- 165) Takahashi, M., R. Matsumaga, K. Komatsu, M. Hajika, S. Sakai, K. Irita, and Y. Nakazawa (2003) Breeding of a new soybean cultivar "Eru-star" and its characteristics. Bull. Kyushu Okinawa Agric. Res. Cent. 42: 49-65 (in Japanese with English summary).
- 166) Takata, S., M. Mizobuchi, M. Kameshima, A. Iwasaki, and Y. Nakamura (2005) A new extremely early rice variety, "Nangokusodachi". Bull. Kochi Pref. Agric. Res. Ctr. 16. <http://www.nogyo.tosa.pref.kochi.lg.jp/download/?f=LD&id=696&fid=46555> (in Japanese).
- 167) Takita, T., T. Higashi, N. Yokogami, T. Kataoka, H. Kato, M. Yamaguchi, Y. Tamura, H. Kowata, Z. Oyamada, and Y. Sunohara (2002) Breeding of a new rice cultivar with low amylose content "Silky-pearl". Bull. Natl. Agric. Res. Cent. Tohoku Reg. 100: 27-39 (in Japanese with English summary).
- 168) Takita, T., T. Yagi, H. Nishiyama, H. Hidaka, H. Arasuna, A. Todoroki, M. Kobae, M. Kawaguchi, I. Aikou, H. Yoshioka, K. Yoshida, and T. Sonoda (1997) A new cultivar "Hohoemi". Bull. Miyazaki Agric. Exp. Stn. 31: 12-25 (in Japanese with English summary).
- 169) Takita, T., H. Yamashita, Y. Nagayoshi, K. Wakasugi, S. Tsutsumi, M. Kawaguchi, H. Yoshioka, H. Arasuna, T. Sonoda, and H. Hidaka (2000) A new rice cultivar Chihonokaori. Bull. Miyazaki Agric. Exp. Stn. 35: 53-66 (in Japanese with English summary).
- 170) Tamura, K., Y. Kiuchi, H. Nakano, A. Abe, T. Sasaki, K. Ogiuchi, S. Nakajo, Y. Ogi, H. Odanaka, M. Takahashi, S. Ogiya, and Y. Kamiyama (2007) Breeding of a new rice variety "Donpishari". Bull. Iwate Agric. Res. Cent. 7: 13-28 (in Japanese with English abstract).
- 171) Tamura, K., M. Okamoto, R. Kaji, H. Hirabayashi, R. Mizobuchi, T. Yagi, H. Yamashita, H. Nishiyama, H. Motomura, T. Takita, and K. Saito (2007) A new rice variety for whole-crop silage, "Nishiaoba". Bull. Kyushu Okinawa Agric. Res. Cent. 48: 31-48 (in Japanese with English summary).
- 172) Tanabu, K., K. Yamazaki, T. Kudo, M. Takadate, K. Arima, T. Mikami, Y. Kawamura, H. Tatsuta, M. Yoshihara, M. Namioka, T. Kanazawa, K. Ono, and H. Himoto (1987) A new rice cultivar "Hanafubuki". Bull. Aomori Agric. Exp. Stn. 30: 19-30 (in Japanese).
- 173) Tanno, H., Y. Kumihira, Y. Ebe, H. Kikuchi, N. Shinbashi, and K. Sugawara (1997) A new rice variety "Aya". Bull. Hokkaido Pref. Agric. Exp. Stn. 72: 37-53 (in Japanese with English summary).
- 174) The Miyazaki Breeding Group of Rice (2005) Hinohikari, Hohoemi, Karinomai and Akireshiki new rice varieties with superior cooking qualities and taste in Kyushu. Breeding Research 7: 195-200 (in Japanese).
- 175) Tokura, K., K. Yagasaki, M. Arakawa, S. Kato, Y. Takei, R. Tateoka, Y. Okada, K. Watanabe, K. Ohtsuka, K. Inoue, T. Furusawa, and T. Niwayama (2002) Breeding of a new rice variety "Banbanzai". Bull. Saitama Pref. Agric. Forest. Res. Cent. 2: 26-35 (in Japanese with English abstract).
- 176) Tokura, K., K. Yagasaki, M. Arakawa, S. Kato, R. Tateoka, Y. Takei, K. Watanabe, K. Ohtsuka, T. Furusawa, K. Inoue, and T. Niwayama (2000) Breeding of a new rice variety "Sakitamahime", Bull. Saitama Agric. Exp. Stn. 52: 1-12. (in Japanese with English summary).
- 177) Tokura, K., K. Yagasaki, S. Kato, M. Arakawa, T. Niwayama, K. Ohtsuka, R. Tateoka, K. Inoue, T. Furusawa, K. Suzuki (1999) Breeding of a new rice variety "Sainoyume". Bull. Saitama Agric. Exp. Stn. 51: 1-17 (in Japanese with English summary).
- 178) Tomita, M. (2006) A new semi-dwarf Koshihikari-type rice cultivar "Hikari-shinseiki". Breeding Research 8 (Suppl. 1): 238 (in Japanese).
- 179) Tomita, K., H. Horiuchi, A. Kobayashi, I. Tanaka, M. Tanoi, H. Furuta, T. Minobe, K. Terada, A. Yamamoto, K. Kanda, K. Aoki, N. Masaki, T. Minami, A. Sugimoto, C. Kagoshima, and K. Horiuchi (2008) "Mampuku-mochi", a new rice cultivar. Bull. Fukui Pref. Agric. Exp. Stn. 45: 1-23 (in Japanese with English summary).
- 180) Tomita, K., H. Horiuchi, A. Kobayashi, M. Tanoi, I. Tanaka, T. Minobe, K. Kanda, T. Hayashi, K. Terada, A. Sugimoto, C. Kagoshima, and K. Horiuchi (2009) "Akisakar", a new rice cultivar. Bull. Fukui Pref. Agric. Exp. Stn. 46: 1-21 (in Japanese with English summary).

- 181) Tomita, K., H. Horiuchi, A. Kobayashi, I. Tanaka, M. Tanoi, H. Furuta, T. Minobe, K. Terada, A. Yamamoto, K. Kanda, K. Aoki, N. Masaki, T. Minami, A. Sugimoto, C. Kagoshima, and K. Horiuchi (2007) "New-hikari", a new rice cultivar. *Bull. Fukui Pref. Agric. Exp. Stn.* 44: 1-20 (in Japanese).
- 182) Tomita, K., H. Horiuchi, K. Terada, M. Tanoi, A. Kobayashi, I. Tanaka, T. Minobe, H. Furuta, A. Yamamoto, H. Shinoyama, I. Ikeda, K. Aoki, N. Masaki, T. Minami, A. Sugimoto, and C. Kagoshima (2005) "Ikubihikari", a new rice cultivar. *Bull. Fukui Pref. Agric. Exp. Stn.* 42: 1-15 (in Japanese with English summary).
- 183) Tsuchiya, T., M. Maeshige, Y. Doi, S. Otake, S. Umemoto, Z. Katsuba, and Y. Sakai (1995) A new paddy rice variety "Akironan". *Bull. Hiroshima Pref. Agric. Res. Cent.* 62: 12-21 (in Japanese with English summary).
- 184) Tsuru, M., T. Yoshida, A. Sasaki, S. Taya, H. Maeda, T. Kiriyama, and K. Ikeda (1981) A new two-rowed barley cultivar "Kawamizuki". *Bull. Natl. Agric. Exp. Stn.* 21: 489-508 (in Japanese with English summary).
- 185) Uchiyamada, H. *et al.* (1979) A new rice cultivar Miyazaki. *Bull. Miyazaki Agric. Res. Inst.* 13: 1-11 (in Japanese with English summary).
- 186) Uehara, Y., N. Horisue, Y. Sunohara, A. Kobayashi, Y. Kawamura, M. Tateyama, Y. Fujimura, M. Suwa, T. Mikami, H. Yokoyama, H. Tatsuta, M. Suto, T. Kudo, and Z. Oyamada (1997) A new rice cultivar "Yumekogane". *Bull. Aomori Agric. Exp. Stn.* 36: 19-37 (in Japanese with English summary).
- 187) Uehara, Y., N. Horisue, Y. Sunohara, Y. Kawamura, M. Tateyama, M. Suwa, Y. Fujimura, H. Yokoyama, T. Mikami, H. Tatsuta, M. Suto, T. Kudo, and Z. Oyamada (1997) A new rice cultivar "Hamayutaka". *Bull. Aomori Agric. Exp. Stn.* 36: 39-55 (in Japanese with English summary).
- 188) Uehara, Y., A. Kobayashi, Y. Koga, K. Fukui, H. Shimizu, H. Ohta, K. Miura, H. Horiuchi, and Y. Fujita (1995) Breeding of a low amylose rice cultivar "Soft 158". *Bull. Hokuriku Agric. Exp. Stn.* 37: 133-153 (in Japanese with English summary).
- 189) Uehara, Y., A. Kobayashi, Y. Koga, H. Ohta, H. Shimizu, K. Miura, K. Fukui, H. Otsuki, Y. Komaki, H. Sasahara, H. Horiuchi, A. Goto, and K. Okuno (2003) A new rice variety "Mebaumochi". *Bull. Natl. Agric. Res. Cent.* 2: 63-81 (in Japanese with English summary).
- 190) Uehara, Y., A. Kobayashi, Y. Koga, H. Ohta, H. Shimizu, K. Miura, K. Fukui, H. Otsuki, Y. Komaki, H. Sasahara, H. Horiuchi, K. Okuno, Y. Fujita, and A. Goto (2003) A new rice variety "Kusayutaka". *Bull. Natl. Agric. Res. Cent.* 2: 83-105 (in Japanese with English summary).
- 191) Uehara, Y., A. Kobayashi, Y. Koga, H. Ohta, H. Shimizu, K. Miura, K. Fukui, H. Otsuki, H. Horiuchi, K. Okuno, Y. Fujita, Y. Komaki, and H. Sasahara (2002) A new rice variety "Asatsuyu". *Bull. Natl. Agric. Res. Cent.* 1: 23-43 (in Japanese with English summary).
- 192) Uehara, Y., A. Kobayashi, Y. Koga, H. Shimizu, K. Miura, H. Ohta, K. Fukui, K. Okuno, H. Horiuchi, Y. Fujita, H. Otsuki, and S. Ishizaka (1998) A new rice cultivar "Awaminori". *Bull. Hokuriku Agric. Exp. Stn.* 41: 1-18 (in Japanese with English summary).
- 193) Uehara, Y., A. Kobayashi, Y. Koga, H. Uchiyamada, K. Miura, K. Fukui, H. Shimizu, H. Ohta, Y. Fujita, K. Okuno, S. Ishizawa, H. Horiuchi, and M. Nakagahra (1995) Breeding of a new rice cultivar "Dontokoi". *Bull. Hokuriku Agric. Exp. Stn.* 37: 107-131 (in Japanese with English summary).
- 194) Uehara, Y., A. Kobayashi, H. Ohta, H. Shimizu, K. Fukui, K. Miura, H. Otsuki, Y. Komaki, and H. Sasahara (2002) A new rice variety "Shunyu". *Bull. Natl. Agric. Res. Cent.* 1: 1-21 (in Japanese with English summary).
- 195) Uehara, Y., A. Kobayashi, H. Shimizu, H. Ohta, K. Miura, K. Fukui, H. Otsuki, Y. Komaki, and H. Sasahara (2000) A new rice cultivar "Itadaki". *Bull. Hokuriku Agric. Exp. Stn.* 43: 1-23 (in Japanese with English abstract).
- 196) Uehara, Y., Y. Sunohara, N. Horisue, Y. Kawamura, M. Suto, H. Yokoyama, K. Maeda, M. Tateyama, M. Suwa, Y. Fujimura, T. Mikami, H. Tatsuta, T. Yashima, S. Kanda, T. Kudo, and Z. Oyamada (2001) A new rice cultivar "Chiyononochi". *Bull. Aomori Agric. Exp. Stn.* 38: 1-17 (in Japanese with English summary).
- 197) Wada, K., Y. Koyama, Y. Nishikawa, R. Hayashi, T. Watanabe, T. Nagashima, T. Hoshino, H. Nakada, N. Iijima, and S. Kato (2002) Breeding of a new rice cultivar "Fusanomai" suitable for sake brewing. *Bull. Chiba Pref. Agric. Res. Cent.* 1: 79-90 (in Japanese with English summary).
- 198) Wada, T., M. Tsubone, T. Inoue, T. Ogata, Y. Hamachi, Y. Matsue, K. F. Oosato, T. Yasunaga, Y. Kawamura, and A. Ishizuka (2010) "Genkitsukushi" a new rice cultivar with tolerance to high temperature during maturity period and high eating quality. *Bull. Fukuoka Agric. Res. Cent.* 29: 1-9 (in Japanese with English abstract).
- 199) Wakamatsu, K., T. Higashi, T. Kamikado, T. Tanogashira, N. Yoshida, T. Shigemitsu, A. Tanaka, and M. Takemure (2005) Breeding a new rice cultivar "Ayanatsuki". *Bull. Kagoshima Pref. Agric. Res. Cent.* 34: 1-14 (in Japanese with English summary).
- 200) Wakamatsu, K., T. Kamikado, M. Tsuyushige, T. Tanogashira, T. Higashi, A. Tanaka, T. Shigemizu, and M. Takemure (2007) Breeding a new rice cultivar "Satsumayukimochi". *Bull. Kagoshima Pref. Inst. Agric. Dev. Agric. Res.* 1: 11-21 (in Japanese with English summary).
- 201) Watanabe, I., T. Nagasawa, S. Murakami, K. Hashimoto, H. Banba, N. Takahashi, A. Okabe, K. Kokubun, S. Matsumoto, the late T. Koyama, S. Nakamura, S. Sakai, K. Igita, K. Murata, and A. Kikuchi (1990) A new soybean variety "Tomoyutaka". *Bull. Tohoku Natl. Agric. Exp. Stn.* 82: 1-17 (in Japanese with English summary).
- 202) Yamaguchi, I., C. Kiribuchi-Otobe, T. Yamagisawa, T. Nagamine, T. Ushiyama, and H. Yoshida (2003) Breeding of 2 waxy wheat cultivars, "Akebono-mochi" and "Ibuki-mochi", and their main features. *Bull. Natl. Inst. Crop Sci.* 3: 21-33 (in Japanese with English summary).
- 203) Yamaguchi, M., N. Yokogami, T. Kataoka, K. Nakagami, T. Takita, T. Higashi, H. Kato, Y. Tamura, H. Kowata, Z. Oyamada, and Y. Sunohara (2005) Breeding of a new rice cultivar, "Churahikari", with blast resistance and excellent eating quality. *Bull. Natl. Agric. Res. Cent. Tohoku Reg.* 104: 1-16 (in Japanese with English summary).
- 204) Yamaguchi, M., N. Yokogami, T. Kataoka, T. Takita, T. Higashi, H. Kato, Y. Tamura, and H. Kowata (2005) Breeding of a new red grain rice cultivar, "Benigoromo". *Bull. Natl. Agric. Res. Cent. Tohoku Reg.* 103: 13-26 (in Japanese with English abstract).
- 205) Yamakawa, K., H. Yasui, T. Mochizuki, K. Hida, and S. Komochi (1987) Incorporation of disease resistance from *Lycopersicon peruvianum* L. to cultivated tomato. I. Breeding of new varieties "Ryugyokki" etc., having resistance to Fusarium root rot and tobacco mosaic virus inherited from *L. peruvianum*. *Bull. Natl. Res. Inst. Vegetable, Ornamental Plants and Tea, series A* 1: 1-37 (in Japanese with English summary).
- 206) Yamakawa, T., T. Murakami, K. Miyamoto, N. Tachibana, F. Hashidume, N. Tatematsu, and H. Hattoni (2002) A newly developed paddy rice variety "Mienoyume", with medium maturation and high grain quality. *Bull. Mie Pref. Agric. Res. Inst.* 29: 15-23 (in Japanese with English summary).
- 207) Yamakawa, T., N. Tatibana, T. Murakami, F. Hashidume, and N. Tatematsu (2000) Improvement of new paddy rice variety Mienoemi, with early maturity and edible quality. *Bull. Mie Pref. Agric. Res. Inst.* 27: 1-10 (in Japanese with English summary).
- 208) Yamamoto, K. S., Matsushita, T., Teramishi, T., Ishibashi, and T. Matsushima (1989) On new glutinous rice cultivar "MANYO-MOCHI". *Hokuriku Crop Science* 24: 31-33 (in Japanese).

- 209) Yamamoto, A., S. Takahashi, M. Sugiyama, N. Omoso, M. Iwamoto, Y. Abe, T. Hirose, M. Kanda, T. Furuyama, M. Fukuda, K. Fujihara, and Y. Kadowaki (1999) A new rice variety "Kannomai". Bull. Shimane Agric. Exp. Stn. 32: 35-50 (in Japanese with English summary).
- 210) Yamazaki, K., K. Tanabu, M. Takadate, T. Mikami, K. Arima, Y. Kawamura, H. Tatsuta, M. Namioka, T. Kanazawa, K. Ono, and H. Mimoto (1987) A new rice cultivar "Mutsuhomare". Bull. Aomori Agric. Exp. Stn. 30: 1-17 (in Japanese with English summary).
- 211) Yamazaki, S., M. Yuzawa, H. Nagashima, S. Aonuma, M. Miyoshi, A. Shinozaki, Y. Izawa, and M. Yamaguchi (2012) A new paddy rice cultivar "Tochigi-no-hoshi". Bull. Tochigi Agr. Exp. Stn. 68: 1-13 (in Japanese with English summary).
- 212) Yatou, O., T. Nagatani, K. Mori, T. Iwashita, O. Doi, A. Shinya, M. Machida, Y. Yamakawa, M. Homan, K. Moriya, T. Higashi, M. Karube, K. Fukui, and Y. Komaki (1994) A new rice cultivar "Natsunotayori". Bull. Kagoshima Pref. Agric. Exp. Stn. 22: 95-97 (in Japanese with English summary).
- 213) Yokogami, N., T. Takita, T. Higashi, M. Yamaguchi, T. Kataoka, Y. Tamura, Z. Oyamada, Y. Sunohara, and H. Kowata (2000) Development of a new purple-grain non-glutinous rice variety "Okunomurasaki". Tohoku Agric. Res. 53: 7-8 (in Japanese).
- 214) Yoshioka, T., T. Tohno-Oka, N. Kawada, M. Yoshida, K. Matsui, M. Tamio, T. Makino, M. Furusho, T. Fukuoka, and S. Miyagawa (2005) Breeding of a new barley cultivar "Sayakaze". Bull. National Institute of Crop Sci. 7: 1-18 (in Japanese with English summary).
- 215) Yuki, K., H. Sakurada, T. Sano, R. Chuba, K. Sato, N. Yokoo, T. Honma, M. Chuba, S. Sato, H. Miyano, M. Mitobe, H. Sato, K. Watanabe, T. Koseki, and Y. Muraoka (2006) Breeding of a new rice cultivar "Dewanosato" (Yamagatasake 86) for sake brewing. Issu. Yamagata Pref. Agricultural Research 88: 25-39 (in Japanese with English summary).
- 216) Yumoto, S., S. Shimada, Y. Takada, T. Sakai, Y. Kono, H. Shimada, K. Takahashi, T. Adachi, K. Tabuchi, A. Kikuchi, K. Murata, S. Sakai, K. Kitamura, M. Ishimoto, K. Igita, Y. Nakazawa, and M. Hajika (2006) A new Tohoku-district soybean cultivar "Suzusyaka" with three lipoxigenase isozyme deletions. Bull. Natl. Agric. Res. Cent. Tohoku Reg. 105: 35-48 (in Japanese with English abstract).